

PATIENT INFORMATION LEAFLET: INFORMATION FOR THE USER

ACNOTIN 20 mg Soft Capsules

Isotretinoin

Read all of this leaflet carefully before you start using this medicine.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your health care provider.
- This medicine has been prescribed for you. Do not pass it on to others. It may harm them, even if their symptoms are the same as yours.
- If any of the side effects becomes serious, or if you notice any side effects not listed in this leaflet, please tell your health care provider.

In this leaflet:

- a) What ACNOTIN is and what it is used for
- b) Before you take ACNOTIN
- c) How to take ACNOTIN
- d) Possible side effects
- e) How to store ACNOTIN
- f) Further information

a) WHAT ACNOTIN IS AND WHAT IT IS USED FOR

ACNOTIN contains isotretinoin – a substance related to vitamin A, and one of a group of medicines called retinoids (for treatment of acne).

ACNOTIN is used to treat severe types of acne (such as nodular or conglobate acne, or acne that is at risk of causing permanent scarring) in adults and adolescents from 12 years of age only after puberty. You will use ACNOTIN when your acne has not got better with anti-acne treatments, including antibiotics and skin treatments.

ACNOTIN treatment must be supervised by a dermatologist (a doctor specialised in the treatment of skin problems).

b) BEFORE YOU TAKE ACNOTIN

Do not take ACNOTIN

- if you are allergic (hypersensitive) to isotretinoin or any of the other ingredients of ACNOTIN
- if you are pregnant or breast-feeding
- if there is any chance you could become pregnant, you must follow the precautions under “Pregnancy prevention programme”.
- if you have liver disease
- if you have very high levels of blood fats (e.g. high cholesterol or triglycerides)
- if you have very high levels of vitamin A in your body (hypervitaminosis A)
- if you are receiving treatment with tetracyclines (a type of antibiotic) at the same time

If any of these apply to you, go back to your doctor before taking any ACNOTIN.

Take special care with ACNOTIN

Talk to your doctor or pharmacist before taking ACNOTIN:

- If you have ever had any kind of mental health problems. This includes depression, aggressive tendencies or mood changes. It also includes thoughts about hurting yourself or ending your life. This is because your mood may be affected while taking ACNOTIN.

Talk to your doctor if you experience persistent pain in your lower back or buttocks during treatment with ACNOTIN. These symptoms may be signs of sacroiliitis, a type of inflammatory back pain. Your doctor may discontinue treatment with ACNOTIN and refer you to a specialist for treatment of inflammatory back pain. Further evaluation may be needed including imaging modalities such as MRI.

Pregnancy prevention programme

Women who are pregnant must not take ACNOTIN

This medicine can seriously harm an unborn baby (the medicine is said to be ‘teratogenic’) – it can cause serious abnormalities of the unborn baby’s brain, face, ear, eye, heart and certain glands (thymus gland and parathyroid gland). It also makes a miscarriage more likely. This may happen even if ACNOTIN is taken only for a short time during pregnancy.

- You must not take ACNOTIN if you are pregnant or if you think you might be pregnant.
- You must not take ACNOTIN if you are breastfeeding. The medicine is likely to pass into your milk and may harm your baby.
- You must not take ACNOTIN if you could get pregnant during treatment.
- You must not get pregnant for one month after stopping this treatment because some medicine may still be left in your body.

Women who could get pregnant are prescribed ACNOTIN under strict rules. This is because of the risk of serious harm to the unborn baby

These are the rules:

- Your doctor must explain the risk of harm to the unborn baby - you must understand why you must not get pregnant and what you need to do to prevent getting pregnant.
- You must have talked about contraception (birth control) with your doctor. The doctor will give you information on how not to get pregnant. The doctor may send you to a specialist for contraception advice.
- Before you start treatment, your doctor will ask you to take a pregnancy test. The test must show that you are not pregnant when starting treatment with ACNOTIN.

Women must use effective contraception before, during and after taking ACNOTIN

- You must agree to use at least one very reliable method of contraception (for example an intra uterine device or contraceptive implant) or, two effective methods that work in different ways (for example a hormonal contraceptive pill and a condom). Discuss with your doctor which methods would be suitable for you.
- You must use contraception for a month before taking ACNOTIN, during treatment and for a month afterwards.
- You must use contraception even if you do not have periods or you are not sexually active (unless your doctor decides this is not necessary).

Women must agree to pregnancy testing before, during and after taking ACNOTIN

- You must agree to regular follow-up visits, ideally every month.
- You must agree to have regular pregnancy tests, ideally every month during treatment and, because some medicine may still be left in your body, 1 month after stopping ACNOTIN (unless your doctor decides this is not necessary in your case).
- You must agree to extra pregnancy tests if your doctor asks you.
- You must not get pregnant during treatment or for a month afterwards because some medicine may still be left in your body.
- Your doctor will discuss all these points with you, using a checklist and will ask you (or a parent/guardian) to sign it. This form confirms that you have been told about the risks and that you will follow the rules above.

If you get pregnant while taking ACNOTIN, **stop taking the medicine straight away**, and contact your doctor. Your doctor may send you to a specialist for advice.

Also, if you become pregnant within one month after you stop taking ACNOTIN, you should contact your doctor. Your doctor may send you to a specialist for advice.

Your doctor has written information on pregnancy prevention for the users of ACNOTIN which should be given to you.

A new prescription is needed for more treatment. Each prescription is only valid for 7 days.

Advice for men

The levels of oral retinoid in the semen of men taking ACNOTIN are too low to harm their partners' unborn baby. However, you must never share your medication with anyone.

Additional precautions

You should never give this medicinal product to another person. Please take any unused capsules to your pharmacist at the end of treatment.

You should not donate blood during treatment with this medicine and for 1 month after stopping ACNOTIN because an unborn baby could be harmed if a pregnant patient receives your blood.

Mental health problems

You may not notice some changes in your mood and behaviour and so it is very important that you tell your friends and family that you are taking this medicine. They may notice these changes and help you quickly identify any problems that you need to talk to your doctor about.

Advice for all patients

- **Tell your doctor if you have ever had any mental illness** (including depression, suicidal behaviour or psychosis), or if you take medicines for any of these conditions.
- **Severe Skin reactions** (e.g. erythema multiforme (EM), Stevens-Johnson syndrome (SJS) and toxic epidermal necrolysis (TEN)) have been reported with the use of ACNOTIN. The rash may progress to widespread blistering or peeling of the skin. You should also look for ulcers in the mouth, throat, nose, genitals and conjunctivitis (red and swollen eyes).
- **Rarely, ACNOTIN may cause severe allergic reactions** some of which can affect skin in the form of eczema, hives and bruises or red patches on arms and legs. If you develop

an allergic reaction, stop taking ACNOTIN, seek urgent advice from a doctor and tell him that you are taking this medicine.

- **Cut down on intensive exercise and physical activity.** ACNOTIN can cause muscle and joint pain particularly in children and teenagers undertaking vigorous physical activity.
- **ACNOTIN has been associated with inflammatory bowel disease.** Your doctor will take you off ACNOTIN if you have severe bloody diarrhoea without any history of gastrointestinal disorders.
- **ACNOTIN may cause dry eyes, intolerance to contact lenses and visual difficulties including decreased night vision.** Cases of dry eyes not resolving after discontinuation of therapy have been reported. Tell your doctor if you have any of these symptoms. Your doctor may ask you to use lubricating eye ointment or tear replacement therapy. If you use contact lenses and you have developed intolerance to contact lenses, you may be advised to wear glasses during the treatment. Your doctor may refer you to a specialist for advice if you develop visual difficulties and you may be asked to stop taking ACNOTIN.
- **Benign intracranial hypertension has been reported with ACNOTIN use** and in some cases where ACNOTIN was used together with tetracyclines (a type of antibiotic). Stop taking ACNOTIN and seek urgent advice from your doctor if you develop symptoms like headache, nausea, vomiting and visual disturbances. Your doctor may refer you to a specialist to check for swelling of optic disk in the eye (papilloedema).
- **ACNOTIN may increase liver enzyme levels.** Your doctor will do blood tests before, during and after ACNOTIN treatment to check these levels. If they stay high, your doctor may lower your dose or take you off ACNOTIN.
- **ACNOTIN commonly increases blood fats,** such as cholesterol or triglycerides. Your doctor will test these levels before, during and after ACNOTIN treatment. It is best that you do not drink alcoholic drinks or that you at least reduce the amount you usually drink while on treatment. Tell your doctor if you already have high blood fats, diabetes (high blood sugars), are overweight, or an alcoholic. You may need blood tests more often. If your blood fats stay high, your doctor may lower your dose, or take you off ACNOTIN.
- **Tell your doctor if you have any kidney problems.** Your doctor may start you on a lower dose of ACNOTIN and then increase it to the maximum tolerated dose.
- **Tell your doctor if you have problems of fructose intolerance.** Your doctor will not prescribe ACNOTIN if you have intolerance to fructose or sorbitol.
- **ACNOTIN may increase blood sugar levels.** In rare cases, people become diabetic. Your doctor may monitor blood sugar levels during treatment, particularly if you already have diabetes, are overweight, or are an alcoholic.
- **Your skin is likely to get dry.** Use a skin moisturising ointment or cream and a lip balm during treatment. To prevent skin irritation you should avoid using exfoliating or anti-acne products.
- **Avoid too much sun and do not use a sun-lamp or sun-bed.** Your skin may become more sensitive to sunlight. Before you go out in the sun, use a sun-protection product with a high protection factor (SPF 15 or higher).
- **Don't have any cosmetic skin treatments.** ACNOTIN may make your skin more fragile. Don't have any waxing (hair removal), dermabrasion or laser treatments (removing horny skin or scars) during treatment, or for at least 6 months after treatment. They could cause scarring, skin irritation, or rarely, changes in the colour of your skin.

Children and adolescents

The use of ACNOTIN in children under the age of 12 is not recommended. This is because it is not known if it is safe or effective in this age group. **Use in adolescents over**

12 years of age only after puberty.

Taking ACNOTIN with other medicines

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines including herbal and non-prescription products.

- **Do not take vitamin A supplements or tetracyclines** (a type of antibiotic), or use any skin treatments for acne while you are on ACNOTIN. It is fine to use moisturisers and emollients (skin creams or preparations that prevent water loss and have a softening effect on the skin).
- **Avoid the use of topical keratolytic or exfoliative anti-acne agents** while you are on ACNOTIN.

Taking ACNOTIN with food and drink

The capsules should be taken with food once or twice daily.

Pregnancy and breast-feeding

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine. For more information on contraception, pregnancy and breast-feeding, see “Pregnancy prevention programme”.

Driving and using machines

You may not see as well at night during your treatment. This can happen suddenly. In rare cases this has continued after the treatment has stopped. Drowsiness and dizziness have been reported very rarely. If this happens to you, you should not drive or operate machinery.

Important information about some of the ingredients of ACNOTIN

ACNOTIN contains

- **Soya-bean oil:** If you are allergic to peanut or soya, do not take this medicinal product.
- **Sorbitol:** If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicine.

c) HOW TO TAKE ACNOTIN

Always take ACNOTIN exactly as your health care provider has told you. You should check with your health care provider if you are not sure.

The usual starting dose is 0.5 mg per kilogram body weight per day (0.5 mg/kg/day). So if you weigh 60 kg, your dose will usually start at 30 mg a day.

Take the capsules once or twice daily. Take on a full stomach. Swallow them whole, with a drink or a mouthful of food.

After a few weeks your doctor may adjust your dose. This depends on how you are getting on with your medicine. For most patients the dose will be between 0.5 and 1.0 mg/kg/day. If you think that ACNOTIN is too strong or too weak, talk to your doctor or pharmacist.

If you have severe kidney problems, you will usually start on a lower dose (such as 10 mg/day) which will be increased up to the highest dose your body can tolerate. If your body can't tolerate the recommended dose, you may be prescribed a lower dose: that can

mean you are treated for longer and your acne might be more likely to come back.

A course of treatment usually lasts for 16 to 24 weeks. Most patients only need one course. Your acne may continue to improve for up to 8 weeks after treatment. You won't usually start another course until then.

Some people find their acne gets worse during the first weeks of treatment. It usually improves as treatment goes on.

Use in children

The use of ACNOTIN in children under the age of 12 is not recommended. This is because it is not known if it is safe or effective in this age group. Use in adolescents over 12 years of age only after puberty.

If you take more ACNOTIN than you should

If you take too many capsules or someone else accidentally takes your medicine, contact your doctor, pharmacist or nearest hospital immediately.

If you forget to take ACNOTIN

If you miss a dose take it as soon as you can. However, if it is nearly time for your next dose, skip the missed dose and carry on as before. Do not take a double dose (two doses close together).

If you stop taking ACNOTIN

Do not stop taking ACNOTIN without first talking to your doctor or pharmacist.

If you have any further questions on the use of this product, ask your health care provider.

d) POSSIBLE SIDE EFFECTS

Like all medicines, ACNOTIN can cause side effects, although not everybody gets them. Some of the side effects associated with the use of isotretinoin are related to the dose. The side effects are generally reversible after changing the dose or stopping treatment, however some may continue after treatment has stopped. Some side effects can be serious and you must immediately contact your doctor.

Side effects requiring immediate medical attention:

Skin problems

Frequency not known (frequency cannot be estimated from available data)

- Serious skin rashes (erythema multiforme, Stevens- Johnson syndrome, and toxic epidermal necrolysis), which are potentially life-threatening and require immediate medical attention. These appear initially as circular patches often with central blisters usually on arms and hands or legs and feet, more severe rashes may include blistering of the chest and back. Additional symptoms such as infection of the eye (conjunctivitis) or ulcers of the mouth, throat or nose may occur. Severe forms of rash may progress to widespread peeling of the skin which can be life threatening. These serious skin rashes are often preceded by headache, fever, body aches (flu-like symptoms).

If you develop a serious rash or these skin symptoms, stop taking ACNOTIN and contact your doctor immediately.

Mental problems

Rare effects (*may affect up to 1 in every 1000 people*)

- Depression or related disorders. Signs of this include sad or altered mood, anxiety, feelings of emotional discomfort,
- Existing depression getting worse.
- Becoming violent or aggressive.

Very rare effects (*may affect up to 1 in every 10,000 people*)

- Some people have had thoughts about hurting themselves or ending their own lives (suicidal thoughts), have tried to end their own lives (attempted suicide), or have ended their lives (suicide). These people may not appear to be depressed.
- Unusual behaviour.
- Signs of psychosis: a loss of contact with reality, such as hearing voices or seeing things that are not there.

Contact your doctor straight away if you get signs of any of these mental problems. Your doctor may tell you to stop taking ACNOTIN. That may not be enough to stop the effects: you may need more help, and your doctor can arrange this.

Allergic reactions

Rare effects (*may affect up to 1 in every 1000 people*)

- Serious (anaphylactic) reactions: difficulty breathing or swallowing caused by sudden swelling of the throat, face, lips and mouth. Also sudden swelling of the hands, feet and ankles.

Very rare effects (*may affect up to 1 in every 10,000 people*)

- Sudden tight chest, shortness of breath and wheezing, particularly if you have asthma.

If you have a serious reaction, get emergency medical help immediately.

If you have any allergic reaction, stop taking ACNOTIN and contact your doctor.

Bones and muscles

Frequency not known (frequency cannot be estimated from available data)

- Muscle weakness which can be potentially life-threatening, may be associated with trouble moving arms or legs, painful, swollen, bruised areas of the body, dark-coloured urine, reduced or no urine output, confusion or dehydration. These are signs of rhabdomyolysis, a breakdown of muscle tissue which can lead to kidney failure. This may occur if you are doing intensive physical activity while you're on ACNOTIN.

Liver and kidney problems

Very rare effects (*may affect up to 1 in every 10,000 people*)

- Yellow skin or eyes, and feeling tired. These can be signs of hepatitis. **Stop taking ACNOTIN straight away and contact your doctor.**
- Difficulty urinating (passing water), swollen and puffy eyelids, feeling excessively tired. These may be signs of kidney inflammation. **Stop taking ACNOTIN straight away and contact your doctor.**

Nervous system problems

Very rare effects (*may affect up to 1 in every 10,000 people*)

- Lasting headache, along with feeling sick (nausea), being sick (vomiting) and change in your eyesight including blurred vision. These may be signs of benign intracranial

hypertension, especially if ACNOTIN is taken with antibiotics called tetracycline. **Stop taking ACNOTIN straight away and contact your doctor.**

Gut and stomach problems

Very rare effects (*may affect up to 1 in every 10,000 people*)

- Severe abdominal (tummy) pain, with or without severe bloody diarrhoea, feeling sick (nausea) and being sick (vomiting). These can be signs of serious gut conditions. **Stop taking ACNOTIN straight away and contact your doctor.**

Eye disorders

Very rare effects (*may affect up to 1 in every 10,000 people*)

- Blurred vision.

If you get blurred vision, stop taking ACNOTIN straight away and contact your doctor. If your sight is affected in any other way tell a doctor as soon as you can.

Other side effects:

Very common side effects with ACNOTIN: (*may affect more than 1 in 10 people*)

- Dryness of the skin, especially of the lips and face; inflamed skin, chapped and inflamed lips, rash, mild itching and slight peeling. Use a moisturising cream from the start of treatment.
- Skin becomes more fragile and redder than usual, especially the face.
- Back pain; muscle pain; joint pain particularly in children and teenagers.

To avoid making any bone or muscle problems worse, cut down on intensive physical activity while you're on ACNOTIN.

- Inflammation of the eye (conjunctivitis) and eyelid area; eyes feel dry and irritated. Ask a pharmacist for suitable eye drops. If you get dry eyes and wear contact lenses, you may need to wear glasses instead.
- Raised liver enzymes seen in blood tests.
- Changed levels of fats in the blood (including HDL or triglycerides).
- Bruising, bleeding or clotting more easily - if clotting cells are affected.
- Anaemia – weakness, dizziness, pale skin – if red blood cells are affected.

Common side effects with ACNOTIN: (*may affect up to 1 in 10 people*)

- Headache.
- Higher levels of cholesterol in the blood.
- Protein or blood in the urine.
- More liable to get infections - if the white blood cells are affected.
- Inside of the nose becomes dry and crusted, causing mild nosebleeds.
- Sore or inflamed throat and nose.
- Allergic reactions such as rash, itchiness. If you have any allergic reaction, stop taking ACNOTIN and contact your doctor.

Rare side effects with ACNOTIN: (*may affect up to 1 in 1,000 people*)

- Hair loss (alopecia). This is usually only temporary. Your hair should return to normal after the treatment ends.

Very rare side effects with ACNOTIN: (*may affect up to 1 in 10,000 people*)

- You may see less well at night; colour blindness and colour vision gets worse.
- Sensitivity to light may increase; you may find that you need to wear sunglasses to protect your eyes from too bright sunlight.

- Other sight problems including blurred vision, distorted vision, cloudy surface on the eye (corneal opacity, cataracts).
- Excessive thirst; frequent need to urinate; blood tests show an increase in your blood sugar. These can all be signs of diabetes.
- Acne can get worse in the first few weeks, but symptoms should improve with time.
- Skin inflamed, swollen, and darker than usual, especially on the face.
- Excess sweating or itching.
- Arthritis; bone disorders (delayed growth, extra growth and changes to bone density); growing bones may stop growing.
- Calcium deposits in soft tissue, sore tendons, high levels of muscle breakdown products in your blood if you exercise vigorously.
- Increased sensitivity to light.
- Bacterial infections at the base of the nail, changes to nails.
- Swellings, discharging, pus.
- Thickened scarring after surgery.
- Increased body hair.
- Convulsions, drowsiness, dizziness.
- Lymph glands may become swollen.
- Dry throat, hoarseness.
- Hearing difficulties.
- Generally feeling unwell.
- High levels of uric acid in the blood.
- Bacterial infections.
- Inflammation of the blood vessels (sometimes with bruising, red patches).

Unknown frequency: *(frequency cannot be estimated from the available data)*

- Dark or cola-coloured urine
- Problems getting or maintaining an erection
- Lower libido
- Breast swelling with or without tenderness in males
- Vaginal dryness
- Sacroiliitis, a type of inflammatory back pain causing pain in your buttocks or lower back
- Inflammation of the urethra

If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your health care provider.

e) HOW TO STORE ACNOTIN

Keep out of the reach and sight of children.

Store below 30°C in a dry place, away from direct sunlight.

Do not use this medicine after the expiry date (EXP) stated on the pack and blister.

Return left over capsules to your pharmacist. Only keep them if your doctor tells you to.

f) FURTHER INFORMATION

What ACNOTIN contains:

- The active pharmaceutical ingredient is isotretinoin.

The other ingredients as follows;

- Capsule Fill: Beewax White, Butylated Hydroxyanisole, Disodium EDTA, Hydrogenated Vegetable Oil and Soyabean oil.
- Capsule Shell: Gelatin, Glycerin, Sorbitol 76% solution (sorbitol special), Purified Water, Carmosine, Brilliant Blue, Ponceau 4R, Iron Oxide Black and Titanium Dioxide.

What ACNOTIN looks like and contents of the pack:

ACNOTIN 20: Orange colour, oily suspension, filled in 6 minim, oval shaped, opaque purple, opaque soft gelatin shell capsules.

ACNOTIN 20 capsules are available as 10 capsules in a blister, 3 blisters are packed in a printed unit carton with a literature insert.

Marketing Authorization Holder

MEGA LIFESCIENCES Public Company Limited
384 Moo 4, Soi 6, Bangpoo Industrial Estate,
Pattana 3 Road, Phraeksa, Mueang,
Samutprakarn 10280, Thailand.

This leaflet was prepared on {04/2023}.