



Clopi-Denk 75

Film-coated tablet – oral use
Active substance: clopidogrel
Platelet aggregation inhibitor

Package leaflet: Information for the patient

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What Clopi-Denk 75 is and what it is used for
2. What you need to know before you take Clopi-Denk 75
3. How to take Clopi-Denk 75
4. Possible side effects
5. How to store Clopi-Denk 75
6. Contents of the pack and other information

1. What Clopi-Denk 75 is and what it is used for

Clopi-Denk 75 contains clopidogrel and belongs to a group of medicines called antiplatelet medicinal products. Platelets are very small structures in the blood which clump together during blood clotting. By preventing this clumping, antiplatelet medicinal products reduce the chances of blood clots forming (a process called thrombosis).

Clopi-Denk 75 is taken by adults to prevent blood clots (thrombi) forming in hardened blood vessels (arteries), a process known as atherothrombosis, which can lead to atherothrombotic events (such as stroke, heart attack, or death).

You have been prescribed Clopi-Denk 75 to help prevent blood clots and reduce the risk of these severe events because:

- You have a condition of hardening of arteries (also known as atherosclerosis), and
- you have previously experienced a heart attack, stroke or have a condition known as peripheral arterial disease, or
- you have experienced a severe type of chest pain known as “unstable angina” or “myocardial infarction” (heart attack). For the treatment of this condition your doctor may have placed a stent in the blocked or narrowed artery to restore effective blood flow. You should also be given acetylsalicylic acid (a substance present in many medicines used to relieve pain and lower fever as well as to prevent blood clotting) by your doctor.

- You have an irregular heart beat, a condition called “atrial fibrillation”, and you cannot take medicines known as “oral anticoagulants” (vitamin K antagonists) which prevent new clots from forming and prevent existing clots from growing. You should have been told that “oral anticoagulants” are more effective than acetylsalicylic acid or the combined use of Clopi-Denk 75 and acetylsalicylic acid for this condition. Your doctor should have prescribed Clopi-Denk 75 plus acetylsalicylic acid if you cannot take “oral anticoagulants” and you do not have a risk of major bleeding.

2. What you need to know before you take Clopi-Denk 75

- Do not take Clopi-Denk 75**
- if you are allergic to clopidogrel or any of the other ingredients of this medicine (listed in section 6)
 - if you have a medical condition that is currently causing bleeding such as a stomach ulcer or bleeding within the brain
 - if you suffer from severe liver disease.
- If you think any of these apply to you, or if you are in any doubt at all, consult your doctor before taking Clopi-Denk 75.

Warnings and precautions

- If any of the situations mentioned below apply to you, you should tell your doctor before taking Clopi-Denk 75:
- If you have a risk of bleeding such as
 - a medical condition that puts you at risk of internal bleeding (such as a stomach ulcer)
 - a blood disorder that makes you prone to internal bleeding (bleeding inside any tissues, organs or joints of your body)
 - a recent serious injury
 - a recent surgery (including dental)

- a planned surgery (including dental) in the next seven days.
- If you have had a clot in an artery of your brain (ischaemic stroke) which occurred within the last seven days.
- If you have kidney or liver disease.
- If you have had an allergy or reaction to any medicine used to treat your disease.

While you are taking Clopi-Denk 75:

- You should tell your doctor if a surgery (including dental) is planned.
- You should also tell your doctor immediately if you develop a medical condition (also known as Thrombotic Thrombocytopenic Purpura or TTP) that includes fever and bruising under the skin that may appear as red pinpoint dots, with or without unexplained extreme tiredness, confusion, yellowing of the skin or eyes (jaundice) (see section 4).
- If you cut or injure yourself, it may take longer than usual for bleeding to stop. This is linked to the way your medicine works as it prevents the ability of blood clots to form. For minor cuts and injuries e.g., cutting yourself, shaving, this is usually of no concern. However, if you are concerned by your bleeding, you should contact your doctor straightaway (see section 4).

- Your doctor may order blood tests.
- Children and adolescents**
Do not give this medicine to children because it does not work.

Other medicines and Clopi-Denk 75
Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines, including medicines obtained without a prescription.

Some other medicines may influence the use of Clopi-Denk 75 or vice versa.

You should specifically tell your doctor if you take

- medicines that may increase your risk of bleeding such as:
 - oral anticoagulants, medicines used to reduce blood clotting
 - a non-steroidal anti-inflammatory medicine, usually used to treat painful and/or inflammatory conditions of muscle or joints
 - heparin or any other injectable medicine used to reduce blood clotting
 - ticlopidine, other antiplatelet agent
 - a selective serotonin reuptake inhibitor (including but not restricted to fluoxetine or fluvoxamine), medicines usually used to treat depression
- omeprazole or esomeprazole, medicines to treat upset stomach

- fluconazole or voriconazole, medicines to treat fungal infections
- efavirenz, or other anti-retroviral medicines (medicines to treat HIV infections)
- carbamazepine, a medicine to treat some forms of epilepsy
- moclobemide, a medicine to treat depression
- repaglinide, a medicine to treat diabetes
- paclitaxel, a medicine to treat cancer
- opioids: while you are treated with clopidogrel, you should inform your doctor before being prescribed any opioid (used to treat severe pain).

If you have experienced severe chest pain (unstable angina or heart attack), you may be prescribed Clopi-Denk 75 in combination with acetylsalicylic acid, a substance present in many medicines used to relieve pain and lower fever. An occasional use of acetylsalicylic acid (no more than 1,000 mg in any 24 hour period) should generally not cause a problem, but prolonged use in other circumstances should be discussed with your doctor.

Clopi-Denk 75 with food and drink
Clopi-Denk 75 may be taken with or without food.

Pregnancy and breast-feeding
If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

It is preferable not to take this product during pregnancy. If you become pregnant while taking Clopi-Denk 75, consult your doctor immediately as it is recommended not to take clopidogrel while you are pregnant.

You should not breast-feed while taking this medicine. If you are breast-feeding or planning to breast-feed, talk to your doctor before taking this medicine. Ask your doctor or pharmacist for advice before taking any medicine.

Driving and using machines
Clopi-Denk 75 is unlikely to affect your ability to drive or to use machines.

Clopi-Denk 75 contains lactose
If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product.

3. How to take Clopi-Denk 75

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

The recommended dose, including for patients with a condition called ‘atrial fibrillation’ (an irregular heartbeat),

is one tablet of Clopi-Denk 75 per day to be taken orally with or without food, and at the same time each day.

If you have experienced severe chest pain (unstable angina or heart attack), your doctor may give you 300 mg of clopidogrel (4 tablets of Clopi-Denk 75) once at the start of treatment. Then, the recommended dose is one tablet of Clopi-Denk 75 per day to be taken as described above. You must take Clopi-Denk 75 for as long as your doctor continues to prescribe it.

If you take more Clopi-Denk 75 than you should
Contact your doctor or the nearest hospital emergency department because of the increased risk of bleeding.

If you forget to take Clopi-Denk 75
If you forget to take a dose of Clopi-Denk 75, but remember within 12 hours of your usual time, take your tablet straight-away and then take your next tablet at the usual time. If you forget for more than 12 hours, simply take the next single dose at the usual time. Do not take a double dose to make up for a forgotten tablet.

If you stop taking Clopi-Denk 75
Do not stop the treatment unless your doctor tells you so. Contact your doctor or pharmacist before stopping. If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects
Like all medicines, this medicine can cause side effects, although not everybody gets them.

Contact your doctor immediately if you experience:

- Fever, signs of infection or extreme tiredness. These may be due to rare decrease of some blood cells.
- Signs of liver problems such as yellowing of the skin and/or the eyes (jaundice), whether or not associated with bleeding which appears under the skin as red pinpoint dots and/or confusion (see section 2).
- Swelling in the mouth or skin disorders such as rashes and itching, blisters of the skin. These may be the signs of an allergic reaction.

The most common side effect reported with clopidogrel is bleeding.
Bleeding may occur as bleeding in the stomach or bowels, bruising, haematoma (unusual bleeding or bruising under the skin), nose bleed, blood in the urine. In a small number of cases, bleeding in the eye, inside the head, the lung or the joints has also been reported.

5. How to store Clopi-Denk 75
Keep this medicine out of the sight and reach of children. Do not use this medicine after the expiry date which is stated on the carton and blister strip after “Exp.”. The expiry date refers to the last day of that month.

Shelf life: 18 months
Store below 30 °C.
Do not throw away any medicines via wastewater. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

works as it prevents the ability of blood clots to form. For minor cuts and injuries e.g., cutting yourself, shaving, this is usually of no concern. However, if you are concerned by your bleeding, you should contact your doctor straightaway (see section 2).

Other side effects include:
Common side effects (may affect up to 1 in 10 people):
Diarrhoea, abdominal pain, indigestion or heartburn.

Uncommon side effects (may affect up to 1 in 100 people):
Headache, stomach ulcer, vomiting, nausea, constipation, excessive gas in stomach or intestines, rashes, itching, dizziness, sensation of tingling and numbness.

Rare side effects (may affect up to 1 in 1,000 people):
Vertigo, enlarged breasts in males.

Very rare side effects (may affect up to 1 in 10,000 people):
Jaundice, severe abdominal pain with or without back pain, fever, breathing difficulties sometimes associated with cough, generalized allergic reactions (for example, overall sensation of heat with sudden general discomfort until fainting), swelling in the mouth, blisters of the skin, skin allergy, sore throat (stomatitis), decrease in blood pressure, confusion, hallucinations, joint pain, muscular pain, changes in taste or loss of taste of food.

Side effects with frequency not known (frequency cannot be estimated from the available data):
Hypersensitivity reactions with chest or abdominal pain, persistent low blood sugar symptoms.

Reporting of side effects
If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. By reporting side effects you can help provide more information on the safety of this medicine.

6. Contents of the pack and other information
Pharmacodynamic properties
Pharmacotherapeutic group: platelet aggregation inhibitors excl. heparin, ATC Code: B01AC04.

Clopidogrel is a prodrug, one of whose metabolites is an inhibitor of platelet aggregation. Clopidogrel must be metabolised by CYP450 enzymes to produce the active metabolite that inhibits platelet aggregation. The active metabolite of clopidogrel selectively inhibits the binding of adenosine diphosphate (ADP) to its platelet P2Y12 receptor and the subsequent ADP-mediated activation of the glycoprotein GPIIb/IIIa complex, thereby inhibiting platelet aggregation. Due to the irreversible binding, platelets exposed are affected for the remainder of their lifespan (approximately 7 – 10 days) and recovery of normal platelet function occurs at a rate consistent with platelet turnover.

Elimination
Following an oral dose of ¹⁴C-labelled clopidogrel in man, approximately 50% was excreted in the urine and approximately 46% in the faeces in the 120-hour interval after dosing. After a single oral dose of 75 mg, clopidogrel has a half-life of approximately 6 hours. The elimination half-life of the main circulating (inactive) metabolite was 8 hours after single and repeated administration.

What Clopi-Denk 75 contains
The active substance is clopidogrel. Each film-coated tablet contains 75 mg of clopidogrel (as clopidogrel besylate). The other ingredients are microcrystalline cellulose, mannitol, hydroxypropylcellulose, crospovidone (type A), citric acid monohydrate, macrogol 6000, stearic acid, talc, hypromellose, lactose monohydrate, iron oxide red, triacetin, titanium dioxide.

General classification for supply
Medicinal product subject to medical prescription.

What Clopi-Denk 75 looks like and contents of the pack
Clopi-Denk 75 are pink, round, biconvex film-coated tablets. Clopi-Denk 75 is available in aluminium/aluminium blisters. Pack size: 30 film-coated tablets.

Marketing Authorisation Holder
DENK PHARMA GmbH & Co. KG
Prinzregentenstr. 79, 81675 München, Germany

Manufacturer of bulk, packaging and batch release
Artesan Pharma GmbH & Co. KG
Wendlandstr. 1, 29439 Lüchow, Germany

Biotransformation
Clopidogrel is extensively metabolised by the liver. *In vitro* and *in vivo*, clopidogrel is metabolised according to two main metabolic pathways: one mediated by esterases and leading to hydrolysis into its inactive carboxylic acid deriva-

tives (85% of circulating metabolites), and one mediated by multiple cytochromes P450. Clopidogrel is first metabolised to a 2-oxo-clopidogrel intermediate metabolite. Subsequent metabolism of the 2-oxo-clopidogrel intermediate metabolite results in formation of the active metabolite, a thiol derivative of clopidogrel. The active metabolite is formed mostly by CYP2C19 with contributions from several other CYP enzymes, including CYP1A2, CYP2B6 and CYP3A4. The active thiol metabolite which has been isolated *in vitro*, binds rapidly and irreversibly to platelet receptors, thus inhibiting platelet aggregation.

The C_{max} of the active metabolite is twice as high following a single 300 mg clopidogrel loading dose as it is after four days of 75 mg maintenance dose. C_{max} occurs approximately 30 to 60 minutes after dosing.

Elimination
Following an oral dose of ¹⁴C-labelled clopidogrel in man, approximately 50% was excreted in the urine and approximately 46% in the faeces in the 120-hour interval after dosing. After a single oral dose of 75 mg, clopidogrel has a half-life of approximately 6 hours. The elimination half-life of the main circulating (inactive) metabolite was 8 hours after single and repeated administration.

What Clopi-Denk 75 contains
The active substance is clopidogrel. Each film-coated tablet contains 75 mg of clopidogrel (as clopidogrel besylate). The other ingredients are microcrystalline cellulose, mannitol, hydroxypropylcellulose, crospovidone (type A), citric acid monohydrate, macrogol 6000, stearic acid, talc, hypromellose, lactose monohydrate, iron oxide red, triacetin, titanium dioxide.

General classification for supply
Medicinal product subject to medical prescription.

What Clopi-Denk 75 looks like and contents of the pack
Clopi-Denk 75 are pink, round, biconvex film-coated tablets. Clopi-Denk 75 is available in aluminium/aluminium blisters. Pack size: 30 film-coated tablets.

Marketing Authorisation Holder
DENK PHARMA GmbH & Co. KG
Prinzregentenstr. 79, 81675 München, Germany

Manufacturer of bulk, packaging and batch release
Artesan Pharma GmbH & Co. KG
Wendlandstr. 1, 29439 Lüchow, Germany

Biotransformation
Clopidogrel is extensively metabolised by the liver. *In vitro* and *in vivo*, clopidogrel is metabolised according to two main metabolic pathways: one mediated by esterases and leading to hydrolysis into its inactive carboxylic acid deriva-

tives (85% of circulating metabolites), and one mediated by multiple cytochromes P450. Clopidogrel is first metabolised to a 2-oxo-clopidogrel intermediate metabolite. Subsequent metabolism of the 2-oxo-clopidogrel intermediate metabolite results in formation of the active metabolite, a thiol derivative of clopidogrel. The active metabolite is formed mostly by CYP2C19 with contributions from several other CYP enzymes, including CYP1A2, CYP2B6 and CYP3A4. The active thiol metabolite which has been isolated *in vitro*, binds rapidly and irreversibly to platelet receptors, thus inhibiting platelet aggregation.

The C_{max} of the active metabolite is twice as high following a single 300 mg clopidogrel loading dose as it is after four days of 75 mg maintenance dose. C_{max} occurs approximately 30 to 60 minutes after dosing.

Elimination
Following an oral dose of ¹⁴C-labelled clopidogrel in man, approximately 50% was excreted in the urine and approximately 46% in the faeces in the 120-hour interval after dosing. After a single oral dose of 75 mg, clopidogrel has a half-life of approximately 6 hours. The elimination half-life of the main circulating (inactive) metabolite was 8 hours after single and repeated administration.

What Clopi-Denk 75 contains
The active substance is clopidogrel. Each film-coated tablet contains 75 mg of clopidogrel (as clopidogrel besylate). The other ingredients are microcrystalline cellulose, mannitol, hydroxypropylcellulose, crospovidone (type A), citric acid monohydrate, macrogol 6000, stearic acid, talc, hypromellose, lactose monohydrate, iron oxide red, triacetin, titanium dioxide.

General classification for supply
Medicinal product subject to medical prescription.

What Clopi-Denk 75 looks like and contents of the pack
Clopi-Denk 75 are pink, round, biconvex film-coated tablets. Clopi-Denk 75 is available in aluminium/aluminium blisters. Pack size: 30 film-coated tablets.

Marketing Authorisation Holder
DENK PHARMA GmbH & Co. KG
Prinzregentenstr. 79, 81675 München, Germany

Manufacturer of bulk, packaging and batch release
Artesan Pharma GmbH & Co. KG
Wendlandstr. 1, 29439 Lüchow, Germany

Biotransformation
Clopidogrel is extensively metabolised by the liver. *In vitro* and *in vivo*, clopidogrel is metabolised according to two main metabolic pathways: one mediated by esterases and leading to hydrolysis into its inactive carboxylic acid deriva-



Clopi-Denk 75

Comprimé pelliculé – voie orale
Substance active : clopidogrel
Inhibiteur de l'agrégation plaquettaire

Notice : information du patient

Veillez lire attentivement cette notice avant de prendre ce médicament car elle contient des informations importantes pour vous.

- Gardez cette notice. Vous pourriez avoir besoin de la relire.
- Si vous avez d'autres questions, interrogez votre médecin ou votre pharmacien.
- Ce médicament vous a été personnellement prescrit. Ne le donnez pas à d'autres personnes. Il pourrait leur être nocif, même si les signes de leur maladie sont identiques aux vôtres.
- Si vous ressentez un quelconque effet indésirable, parlez-en à votre médecin ou votre pharmacien. Ceci s'applique aussi à tout effet indésirable qui ne serait pas mentionné dans cette notice. Voir rubrique 4.

Que contient cette notice

1. Qu'est-ce que Clopi-Denk 75 et dans quel cas est-il utilisé
2. Quelles sont les informations à connaître avant de prendre Clopi-Denk 75
3. Comment prendre Clopi-Denk 75
4. Quels sont les effets indésirables éventuels
5. Comment conserver Clopi-Denk 75
6. Contenu de l'emballage et autres informations

1. Qu'est-ce que Clopi-Denk 75 et dans quel cas est-il utilisé

Clopi-Denk 75 contient du clopidogrel et appartient à une classe de médicaments appelés « antiagrégants plaquet-taires ». Les plaquettes sont de très petits éléments circulant dans le sang et qui s'agrègent lors de la coagulation

du sang. En empêchant cette agrégation, les antiagrégants plaquet-taires réduisent le risque de formation de caillots sanguins (phénomène appelé « thrombose »).

Clopi-Denk 75 est utilisé chez l'adulte pour éviter la formation de caillots sanguins (thrombus) dans les vaisseaux sanguins (artères) devenus rigides. Cette maladie également appelée « athérombose » peut conduire à la survenue d'événements athérombotiques (tels que l'accident vasculaire cérébral, la crise cardiaque, ou le décès).

On vous a prescrit Clopi-Denk 75 pour empêcher la formation de caillots sanguins et réduire le risque de survenue de tels événements graves car :

- Vous avez des artères qui se sont rigidifiées (aussi connu sous le nom d'athérosclérose), et
- vous avez déjà eu une crise cardiaque, un accident vasculaire cérébral ou vous avez une artériopathie des membres inférieurs, ou
- vous avez une douleur thoracique grave connue sous le nom d'« angor instable » ou d'« infarctus du myocarde » (crise cardiaque). Pour cela, vous avez pu bénéficier d'une pose de stent dans l'artère bouchée ou rétrécie afin de rétablir une circulation sanguine efficace. Votre médecin doit également vous prescrire de l'acide acétylsalicylique (substance présente dans de nombreux médicaments utilisée pour soulager la douleur et faire baisser la fièvre, mais aussi pour prévenir la formation de caillots sanguins).
- Vous avez une fréquence cardiaque irrégulière, maladie appelée fibrillation auriculaire, et vous ne pouvez pas être traité(e) par les médicaments appelés « anticoagulants oraux » (antivitamines K) qui empêchent la formation de nouveaux caillots sanguins et la croissance de caillots sanguins déjà présents. Les « anticoagulants oraux » sont plus efficaces que l'acide acétylsalicylique ou que l'association de Clopi-Denk 75 et d'acide acétylsalicylique pour cette maladie. Votre médecin vous a prescrit Clopi-Denk 75 et de l'acide acétylsalicylique si vous ne pouvez pas être traité(e) par des « anticoagulants oraux » et si vous ne présentez pas de risque majeur de saignement.

2. Quelles sont les informations à connaître avant de prendre Clopi-Denk 75

Ne prenez jamais Clopi-Denk 75

- si vous êtes allergique au clopidogrel ou à l'un des autres composants contenus dans ce médicament (mentionnés dans la rubrique 4)
- si vous avez une maladie actuellement responsable d'un saignement tel qu'un ulcère de l'estomac ou un saignement dans le cerveau

- si vous souffrez d'insuffisance hépatique sévère.
- Si vous pensez être dans l'un de ces cas ou en cas de doute, consultez votre médecin avant de prendre Clopi-Denk 75.

Avertissements et précautions

Si l'une des situations mentionnées ci-dessous s'applique à votre cas, vous devez en avertir votre médecin avant de prendre Clopi-Denk 75 :

- Si vous avez un risque hémorragique tel que :
 - une maladie qui peut provoquer un saignement interne (comme un ulcère de l'estomac)
 - des troubles de la coagulation favorisant des hémorragies internes (saignement au sein d'un tissu, d'un organe ou d'une articulation)
 - une blessure grave récente
 - une intervention chirurgicale récente (y compris dentaire)
 - une intervention chirurgicale (y compris dentaire) prévue dans les 7 jours à venir.
- Si vous avez eu un caillot dans une artère de votre cerveau (accident vasculaire cérébral ischémique) survenu dans les sept derniers jours.
- Si vous présentez une maladie du foie ou des reins.
- Si vous avez des antécédents d'allergie ou de réactions allergiques à tout médicament utilisé pour traiter votre maladie.

Pendant la prise de Clopi-Denk 75 :

- Vous devez avertir votre médecin si une intervention chirurgicale est programmée (y compris dentaire).
- Vous devez aussi avertir votre médecin immédiatement si vous présentez une maladie (appelée « purpura thrombopénique thrombotique » ou « PTT ») incluant fièvre et bleus sous la peau, pouvant apparaître comme des petites têtes d'épingles rouges, accompagné ou non de fatigue extrême inexplicable, confusion, jaunissement de la peau ou des yeux (jaunisse) (voir rubrique 4).
- Si vous vous coupez ou si vous vous blessez, l'arrêt du saignement peut demander plus de temps que d'habitude. Ceci est lié au mode d'action de votre médicament qui empêche la formation de caillots sanguins. Dans le cas de coupures ou blessures superficielles (par exemple, au cours du rasage), vous ne devriez généralement rien constater d'anormal. Cependant, si ce saignement vous préoccupe, vous devez en avertir immédiatement votre médecin (voir rubrique 4).
- Votre médecin pourra vous demander de pratiquer des examens sanguins.

Enfants et adolescents
Ce médicament ne doit pas être utilisé chez les enfants en raison de l'absence d'efficacité.

Autres médicaments et Clopi-Denk 75
Informez votre médecin ou pharmacien si vous prenez, avez récemment pris ou pourriez prendre tout autre médicament, même s'il s'agit d'un médicament obtenu sans ordonnance.

Certains médicaments peuvent exercer une influence sur l'utilisation de Clopi-Denk 75 ou vice-versa. Vous devez informer spécifiquement votre médecin si vous prenez :

- des médicaments qui peuvent augmenter votre risque de saignement tels que :
 - des anticoagulants oraux (médicaments utilisés pour diminuer la coagulation du sang)
 - un anti-inflammatoire non stéroïdien (médicament utilisé habituellement pour traiter la douleur et/ou les maladies inflammatoires des muscles ou des articulations)
 - de l'héparine ou tout autre médicament injectable utilisé pour diminuer la coagulation du sang
 - de la ticlopidine, un autre antiagrégant plaquet-taire
 - un inhibiteur sélectif de la recapture de la sérotonine (incluant notamment la fluoxétine ou la fluvoxamine), médicament utilisé habituellement dans le traitement de la dépression
- de l'oméprazole ou de l'ésoméprazole pour des maux d'estomac
- du fluconazole ou du voriconazole, qui sont des médicaments utilisés dans le traitement d'infections fongiques
- de l'éfavirenz, ou d'autres médicaments antirétroviraux (médicaments pour le traitement des infections par le VIH)
- de la carbamazépine pour le traitement de certaines formes d'épilepsies
- du moclobémide, pour le traitement de la dépression
- du répaglinide, médicament utilisé pour traiter le diabète
- du paclitaxel, médicament utilisé pour traiter un cancer
- des opioïdes : pendant votre traitement par le clopidogrel, vous devez informer votre médecin avant toute prescription d'opioïdes (utilisés pour traiter les douleurs sévères)
- Votre médecin pourra vous demander de pratiquer des examens sanguins.

Si vous avez eu une douleur thoracique grave (angor instable ou crise cardiaque), Clopi-Denk 75 peut vous être prescrit en association avec de l'acide acétylsalicylique, substance présente dans de nombreux médicaments utilisés pour soulager la douleur et faire baisser la fièvre. Une utilisation occasionnelle d'acide acétylsalicylique

SPECIFICATION - DENK PHARMA

product: Clopi-Denk 75 (30°C)

artwork: leaflet

language(s): EN, FR, PT

material number: 6946/002

dimension: 628 x 148 mm

prefold: 2 x pre-folded

pharma code: 7293

printing colours: ♦ PANTONE YELLOW C

♦ Prozess BLACK

revision/date: 1 / 19.04.2020, mt-g

