

**Declot (Warfarin Sodium Tablets USP 5mg)**

**PATIENT INFORMATION LEAFLET: INFORMATION FOR THE USER**

**Declot (Warfarin Sodium Tablet USP 5mg)**

**Category of Distribution:** Prescription Preparation (PP)

**Pharmacological Classification:** Anticoagulant

**ATC Code:** B01AA03

**Read this entire leaflet carefully before you start taking this medicine.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, pharmacist or nurse.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet.

**What is in this leaflet:**

1. What is Warfarin Tablet and what is it used for.
2. What you need to know before you take Warfarin Tablet
3. How to take Warfarin Tablet
4. Possible side effects
5. How to store Warfarin Tablet
6. Contents of the pack and other information

**1. WHAT WARFARIN TABLET IS AND WHAT IT IS USED FOR**

Warfarin is prescription medicine used to treat blood clots and to lower the chance of blood clots forming in your body. Blood clots can cause a stroke, heart attack, or other serious conditions if they form in the legs or lungs.

**2. WHAT YOU NEED TO KNOW BEFORE YOU TAKE WARFARIN TABLET**

**Do not take Warfarin tablet if you**

- your risk of having bleeding problems is higher than the possible benefit of treatment. Your healthcare provider will decide if Warfarin is right for you.

## Declot (Warfarin Sodium Tablets USP 5mg)

### PATIENT INFORMATION LEAFLET: INFORMATION FOR THE USER

- you are pregnant unless you have a mechanical heart valve. Warfarin may cause birth defects, miscarriage, or death of your unborn baby.
- you are allergic to Warfarin or any of the other ingredients in Warfarin.

### **Before taking Warfarin, tell your healthcare provider about all of your medical conditions, including if you:**

- have bleeding problems
- fall often
- have liver problems
- have kidney problems or are undergoing dialysis
- have high blood pressure
- have a heart problem called congestive heart failure
- have diabetes
- plan to have any surgery or a dental procedure
- are pregnant or plan to become pregnant.

- Females who can become pregnant should use effective birth control during treatment, and for at least 1 month after the last dose of Warfarin.

- are breastfeeding. You and your healthcare provider should decide if you will take Warfarin and breastfeed. Check your baby for bruising or bleeding if you take Warfarin and breastfeed.

Tell all of your healthcare providers and dentists that you are taking Warfarin. They should talk to the healthcare provider who prescribed Warfarin for you before you have **any** surgery or dental procedure.

Your Warfarin may need to be stopped for a short time or you may need your dose adjusted.

**Tell your healthcare provider about all the medicines you take**, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

### **Other medicines and Warfarin Tablets**

You may have a higher risk of bleeding if you take Warfarin and other medicines that increase your risk of bleeding, including:

## **Declot (Warfarin Sodium Tablets USP 5mg)**

### **PATIENT INFORMATION LEAFLET: INFORMATION FOR THE USER**

- a medicine that contains heparin
- other medicines to prevent or treat blood clots
- nonsteroidal anti-inflammatory drugs (NSAIDs)

**Tell your healthcare provider if you take any of these medicines. Ask your healthcare provider if you are not sure if your medicine is one listed above.**

Many other medicines can interact with Warfarin and affect the dose you need or increase Warfarin side effects. Do not change or stop any of your medicines or start any new medicines before you talk to your healthcare provider.

Do not take other medicines that contain Warfarin sodium while taking Warfarin.

### **Warfarin Tablets with food and drink**

**Some foods and beverages can interact with Warfarin and affect your treatment and dose.**

Eat a normal, balanced diet. Talk to your healthcare provider before you make any diet changes. Do not eat large amounts of leafy, green vegetables. Leafy, green vegetables contain vitamin K.

Certain vegetable oils also contain large amounts of vitamin K. Too much vitamin K can lower the effect of Warfarin.

## **3. HOW TO TAKE WARFARIN TABLETS**

- **Take Warfarin exactly as prescribed.** Your healthcare provider will adjust your dose from time to time depending on your response to Warfarin.
- **You must have regular blood tests and visits with your healthcare provider to monitor your condition.**
- Call your healthcare provider right away if you:
  - take too much Warfarin
  - are sick with diarrhea, an infection, or have a fever
  - fall or injure yourself, especially if you hit your head. Your healthcare provider may need to check you.

**Declot (Warfarin Sodium Tablets USP 5mg)**

## **PATIENT INFORMATION LEAFLET: INFORMATION FOR THE USER**

### **If you take more Warfarin Tablets than you should**

Do not take more medicine than your doctor tells you to. If you or anyone else has swallowed more tablets than doctor have prescribed, contact your nearest hospital casualty department or doctor immediately. Take the medicine pack with you.

### **If you forget to take Warfarin Tablets**

If you miss a dose of Warfarin, call your healthcare provider. Take the dose as soon as possible on the same day.

**Do not** take a double dose of Warfarin the next day to make up for a missed dose.

## **POSSIBLE SIDE EFFECTS**

Warfarin can cause bleeding which can be serious and sometimes lead to death. This is because Warfarin is a blood thinner medicine that lowers the chance of blood clots forming in your body.

- You may have a higher risk of bleeding if you take Warfarin and:
  - are 65 years of age or older
  - have a history of stomach or intestinal bleeding
  - have high blood pressure (hypertension)
  - have a history of stroke, or “mini-stroke” (transient ischemic attack or TIA)
  - have serious heart disease
  - have a low blood count or cancer
  - have had trauma, such as an accident or surgery
  - have kidney problems
- **Death of skin tissue (skin necrosis or gangrene).** This can happen soon after starting Warfarin. It happens because blood clots form and block blood flow to an area of your body. Call your healthcare provider right away if you have pain, color, or temperature change to any area of your body. You may need medical care right away to prevent death or loss (amputation) of your affected body part.
- **Kidney problems.** Kidney injury may happen in people who take Warfarin. Tell your healthcare provider right away if you develop blood in your urine. Your healthcare

## **Declot (Warfarin Sodium Tablets USP 5mg)**

### **PATIENT INFORMATION LEAFLET: INFORMATION FOR THE USER**

provider may do tests more often during treatment with Warfarin to check for bleeding if you already have kidney problems.

- **“Purple toes syndrome.”** Call your healthcare provider right away if you have pain in your toes and they look purple in color or dark in color.

These are not all of the side effects of Warfarin. For more information, ask your healthcare provider or pharmacist.

### **HOW TO STORE WARFARIN TABLETS**

- ‘Store below 30°C in a dry place Protect from light’.
- Keep Warfarin in a tightly closed container.
- Follow your healthcare provider or pharmacist instructions about the right way to throw away outdated or unused Warfarin.
- Females who are pregnant should not handle crushed or broken Warfarin tablets.

Keep Warfarin and all medicines out of the reach of children.

### **CONTENTS OF THE PACK AND OTHER INFORMATION**

#### **What Warfarin Tablets contain**

The **active** substance is Warfarin sodium

The **other** ingredients are: lactose monohydrate, Pregelatinized starch, Hydroxy propyl cellulose, Isopropyl alcohol and magnesium stearate

#### **What Warfarin Tablets looks like and contents of the pack**

‘Peach colored, round, biconvex, uncoated tablets debossed with ‘M’ and ‘83’ separated by scoreline on one side and plain on another side.’

HDPE container of 30/100 Tablets, Such 1 container is packed in a carton along with pack insert.

**Declot (Warfarin Sodium Tablets USP 5mg)**

**PATIENT INFORMATION LEAFLET: INFORMATION FOR THE USER**

**Supplier**

Macleods Pharmaceuticals Ltd.  
304, Atlanta Arcade,  
Marol Church Road, Andheri (East)  
Mumbai- 400 059, India  
Phone: +91-22-66762800  
Fax: +91-22-2821 6599  
E-mail: [exports@macleodspharma.com](mailto:exports@macleodspharma.com)

**Manufacturer**

Macleods Pharmaceuticals Ltd  
Block N-2, Village: Theda,  
PO: Lodhimajra, Tehsil Baddi,  
Dist: Solan,  
Himachal Pradesh – 174 101, India