

PATIENT INFORMATION LEAFLET

Feverex Toto Suspension (Each 5mL contains: Paracetamol)

Read all of this leaflet carefully before you start using this medicine

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, or pharmacist.
- This medicine has been prescribed for your child. Do not pass it on to others. It may harm them, even if their symptoms are the same as yours.
- If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

In this leaflet:

- a) What Feverex Toto Suspension is and what it is used for
 - b) Before you give Feverex Toto Suspension to your child
 - c) How to give Feverex Toto Suspension to your child
 - d) Possible side effects
 - e) How to store Feverex Toto Suspension
 - f) Further information
- a) **WHAT FEVEREX TOTO SUSPENSION IS AND WHAT IT IS USED FOR**
Feverex Toto Suspension belongs to a group of medicines called pain-killers or analgesics. It is used for the relief of painful or feverish conditions such as teething, headaches, toothache, earache, muscular pains, symptoms of cold and flu.
- b) **BEFORE YOU GIVE FEVEREX TOTO SUSPENSION TO YOUR CHILD**

Do not give Feverex Toto Suspension

- If your child is allergic to paracetamol or any of the other ingredients of Feverex Toto Suspension
- If your child is taking any other paracetamol containing products, other flu, cold or decongestant products, or alcohol

Take special care with Feverex Toto Suspension

- If your child has liver or kidney problems
- If your child is underweight or malnourished
- If your child has an inherited intolerance to fructose or been diagnosed with an intolerance to some other sugars

Taking other medicines

Please tell your doctor or pharmacist if your child is taking or have recently taken any other medicines, including medicines obtained without a prescription.

c) **HOW TO GIVE FEVEREX TOTO SUSPENSION**

Always give Feverex Toto Suspension exactly as your doctor or healthcare provider has told you. You should check with your doctor or pharmacist if you are not sure.

The usual dose is;

Age (Years)	Dosage
3 months to less than 1 year	Half a spoonful (2.5ml)
1 year to less than 3 years	One spoonful (5ml)
3 years to less than 5 years	One and a half spoonful (7.5ml)
5 years to less than 8 years	Two spoonful (10ml)
8 years to less than 12 years	Three spoonful (15ml)
The dosage may be repeated up to four times daily	

If you give your child more Feverex Toto Suspension than you should

Seek immediate medical advice in the event of an overdose, even if your child feels well, because of the risk of delayed, serious liver damage.

If you give your child more suspension at the same time, or you think the child may have swallowed any, contact your nearest hospital casualty department or tell your doctor immediately. Symptoms of an overdose include feeling or being sick, loss of appetite, stomach pain or liver damage.

If you forget to give Feverex Toto Suspension

Give the next dose when needed, provided that the last dose was given at least 4 hours ago. Do not give a double dose to make up for a forgotten dose.

d) POSSIBLE SIDE EFFECTS

Like all medicines, Feverex Toto Suspension can cause side effects, although not everybody gets them. Some possible side effects include; skin rashes and haematological reactions, vomiting, gastrointestinal oedema. Liver damage and renal tubular necrosis may occur following over dosage.

If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

e) HOW TO STORE FEVEREX TOTO SUSPENSION

Keep out of the reach and sight of children.

Do not store above 30 °C. Store in a dry place. Protect from light and from moisture.

Do not use Feverex Toto Suspension after the expiry date which is stated on the label and carton after "EXP". The expiry date refers to the last day of that month.

Medicines should not be disposed of via waste water or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

f) FURTHER INFORMATION

What Feverex Toto Suspension contains

- The active substance is Paracetamol 120mg per 5mL

- The other ingredients are; Xanthan Gum, Sodium Benzoate, Sodium Propyl Parabenzoate, Citric Acid, Raspberry Colour, Raspberry Flavour Oil, Glycerin, Sorbitol Solution, Rectified Spirit and Purified Water

What Feverex Toto suspension looks like and contents of the pack

A red viscous syrupy liquid with a sweet taste and an odour of raspberry packed in amber coloured bottle

Pack size: 60mL

Name and full physical address of Marketing Authorization Holder and Manufacturing Site:-



Beta Healthcare International limited
Plot No.: LR: 209/6554, Mogadishu Road, Industrial Area
P.O Box 42569-00100 Nairobi, Kenya.
Tel: 254 020 2652042/89
Email: info@ke.betashelys.com
Website: www.betacare.co.ke

This leaflet was last approved in 02/2023.