

Sandoz	Business use only	Page 1 of 10
1.3.1 spc-label-pl - common-spc - 3,016 (NL/H/1171/001/IA/016)		20130722
METFORMIN HYDROCHLORIDE 500 MG; 850 MG FILM-COATED TABLET		721-5715.00 721-5716.00

SUMMARY OF PRODUCT CHARACTERISTICS

1 NAME OF THE MEDICINAL PRODUCT

Metformin 500 mg film-coated tablets

Metformin 850 mg film-coated tablets

2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Metformin 500 mg film-coated tablets

Each Metformin 500 mg film-coated tablet contains 500 mg metformin hydrochloride equivalent to 390 mg metformin base.

Metformin 850 mg film-coated tablets

Each Metformin 850 mg film-coated tablet contains 850 mg metformin hydrochloride equivalent to 662.9 mg metformin base.

For the full list of excipients, see section 6.1.

3 PHARMACEUTICAL FORM

Film-coated tablet

Metformin 500 mg film-coated tablets

Round, white, double radius film-coated tablet, embossed “M 500” on one side.

Metformin 850 mg film-coated tablets

Oval, white film-coated tablet with a score on one side and embossed “M 850” on the other side.

4 CLINICAL PARTICULARS

4.1 Therapeutic Indications

Treatment of type 2 diabetes mellitus, particularly in overweight patients, when dietary management and exercise alone does not result in adequate glycaemic control.

- In adults, Metformin 500 mg/ 850 mg may be used as monotherapy or in combination with other oral antidiabetic agents, or with insulin.
- In children from 10 years of age and adolescents, Metformin 500 mg/ 850 mg may be used as monotherapy or in combination with insulin.

A reduction of diabetic complications has been shown in overweight type 2 diabetic adult patients treated with metformin as first-line therapy after diet failure (see section 5.1).

4.2 Posology and Method of Administration

Adults

Monotherapy and combination with other antidiabetic agents

- The usual starting dose is 500 mg or 850 mg metformin hydrochloride 2 or 3 times daily given during or after meals. After 10 to 15 days the dose should be adjusted on the basis of blood

Sandoz	Business use only	Page 2 of 10
1.3.1 spc-label-pl - common-spc - 3,016 (NL/H/1171/001/IA/016)		20130722
METFORMIN HYDROCHLORIDE 500 MG; 850 MG FILM-COATED TABLET		721-5715.00 721-5716.00

glucose measurements. A slow increase of dose may improve gastrointestinal tolerability. The maximum recommended dose of metformin hydrochloride is 3 g daily.

- If transfer from another oral antidiabetic agent is intended: discontinue the other agent and initiate metformin at the dose indicated above.

Combination with insulin

Metformin and insulin may be used in combination therapy to achieve better blood glucose control. Metformin hydrochloride is given at the usual starting dose of 500 mg or 850 mg 2-3 times daily while insulin dosage is adjusted on the basis of blood glucose measurements.

Elderly

Due to the potential for decreased renal function in elderly subjects, the metformin dosage should be adjusted based on renal function. Regular assessment of renal function is necessary (see section 4.4).

Children and adolescents

Monotherapy and combination with insulin

Metformin can be used in children from 10 years of age and adolescents

- The usual starting dose is one tablet of 500 mg or 850 mg once daily, given during meals or after meals
- After 10 to 15 days the dose should be adjusted on the basis of blood glucose measurements. A slow increase of dose may improve gastrointestinal tolerability. The maximum recommended dose of Metformin hydrochloride is 2 g daily, taken as 2 or 3 divided doses.

4.3 Contraindications

- Hypersensitivity to metformin or to any of the excipients.
- Diabetic ketoacidosis, diabetic pre-coma.
- Renal failure or renal dysfunction (creatinine clearance < 60 mL/min) (see section 4.4).
- Acute conditions with the potential to alter renal function such as:
 - dehydration,
 - severe infection,
 - shock
- Acute or chronic disease which may cause tissue hypoxia such as:
 - cardiac or respiratory failure,
 - recent myocardial infarction,
 - shock
- Hepatic insufficiency, acute alcohol intoxication, alcoholism.

4.4 Special Warnings and Special Precautions for Use

Lactic acidosis

Lactic acidosis is a rare, but serious (high mortality in the absence of prompt treatment), metabolic complication that can occur due to metformin accumulation. Reported cases of lactic acidosis in patients on metformin have occurred primarily in diabetic patients with significant renal failure. The incidence of lactic acidosis can and should be reduced by assessing also other associated risk factors, such as poorly controlled diabetes, ketosis, prolonged fasting, excessive alcohol intake, hepatic insufficiency and any condition associated with hypoxia.

Sandoz	Business use only	Page 3 of 10
1.3.1 spc-label-pl - common-spc - 3,016 (NL/H/1171/001/IA/016)		20130722
METFORMIN HYDROCHLORIDE 500 MG; 850 MG FILM-COATED TABLET		721-5715.00 721-5716.00

Diagnosis:

The risk of lactic acidosis must be considered in the event of non-specific signs such as muscle cramps with digestive disorders as abdominal pain and severe asthenia

Lactic acidosis is characterised by acidotic dyspnea, abdominal pain and hypothermia followed by coma. Diagnostic laboratory findings are decreased blood pH, plasma lactate levels above 5 mmol/L, and an increased anion gap and lactate/pyruvate ratio.

If metabolic acidosis is suspected, metformin should be discontinued and the patient should be hospitalised immediately (see section 4.9).

Renal function

As metformin is excreted by the kidney, creatinine clearance (this can be estimated from serum creatinine levels by using the Cockcroft-Gault formula) should be determined before initiating treatment and regularly thereafter:

- * at least annually in patients with normal renal function,
- * at least two to four times a year in patients with creatinine clearance level at the lower limit of normal and in elderly subjects.

Decreased renal function in elderly subjects is frequent and asymptomatic. Special caution should be exercised in situations where renal function may become impaired, for example when initiating antihypertensive therapy, diuretic therapy or when starting therapy with an NSAID (see section 4.3).

Administration of iodinated contrast agent

As the intravascular administration of iodinated contrast agents in radiologic studies can lead to renal failure metformin must be discontinued prior to, or at the time of the test and not be reinstituted until 48 hours afterwards, and only after renal function has been re-evaluated and found to be normal (see section 4.5).

Surgery

Metformin must be discontinued 48 hours prior to elective surgeries under general, spinal or peridural anaesthesia. Therapy may be restarted no earlier than 48 hours following surgery or resumption of oral nutrition and only if normal renal function has been established.

Other precautions:

- All patients should continue their diet with a regular distribution of carbohydrate intake during the day. Overweight patients should continue their energy-restricted diet.
- The usual laboratory tests for diabetes monitoring should be performed regularly.
- Metformin alone does not cause hypoglycaemia, although caution is advised when it is used in combination with insulin or other oral antidiabetics (e.g sulfonylureas or meglitinides).

Children and adolescents

The diagnosis of type 2 diabetes mellitus should be confirmed before treatment with metformin is initiated.

No effect of metformin on growth and puberty has been detected during controlled clinical studies of one-year duration but no long-term data on these specific points are available. Therefore, a careful

Sandoz	Business use only	Page 4 of 10
1.3.1 spc-label-pl - common-spc - 3,016 (NL/H/1171/001/IA/016)		20130722
METFORMIN HYDROCHLORIDE 500 MG; 850 MG FILM-COATED TABLET		721-5715.00 721-5716.00

follow-up of the effect of metformin on these parameters in metformin-treated children, especially pre-pubescent children, is recommended.

Children aged between 10 and 12 years

Only 15 subjects aged between 10 and 12 years were included in the controlled clinical studies conducted in children and adolescents. Although efficacy and safety of metformin in these children did not differ from efficacy and safety in older children and adolescents, particular caution is recommended when prescribing to children aged between 10 and 12 years.

4.5 Interaction with Other Medicinal Products and Other Forms of Interaction

Concomitant use is not recommended:

Alcohol

Acute alcohol intoxication is associated with an increased risk of lactic acidosis, particularly in case of:

- fasting or malnutrition
- hepatic insufficiency

Avoid consumption of alcohol and alcohol-containing medicinal product.

Iodinated contrast agents

Intravascular administration of iodinated contrast agents may lead to renal failure, resulting in metformin accumulation and an increased risk of lactic acidosis (see section 4.4).

Metformin must be discontinued prior to or at the time of the test and not be reinstituted until 48 hours afterwards and only after renal function has been re-evaluated and found to be normal (see section 4.4).

Combinations requiring precautions for use

- *Medicinal products with intrinsic hyperglycaemic activity* as glucocorticoids (systemic or by local route) and sympathomimetics. More frequent blood glucose monitoring may be required, especially at the beginning of treatment. If necessary, adjust the metformin dosage during the therapy with the respective medicinal products.
- *ACE-inhibitors* may decrease the blood glucose levels. If necessary, adjust the dosage of the antidiabetic medicinal product during therapy with the other medicinal product and upon its discontinuation.
- *Diuretics especially loop diuretics*, may increase the risk of lactic acidosis due to their potential to decrease renal function.

4.6 Pregnancy and Lactation

Pregnancy

Uncontrolled diabetes during pregnancy (gestational or permanent) is associated with increased risk of congenital abnormalities and perinatal mortality.

A limited amount of data from the use of metformin in pregnant women does not indicate an increased risk of congenital abnormalities. Animal studies do not indicate harmful effects with respect

Sandoz	Business use only	Page 5 of 10
1.3.1 spc-label-pl - common-spc - 3,016 (NL/H/1171/001/IA/016)		20130722
METFORMIN HYDROCHLORIDE 500 MG; 850 MG FILM-COATED TABLET		721-5715.00 721-5716.00

to pregnancy, embryonal or foetal development, parturition or postnatal development (see also section 5.3).

When the patient plans to become pregnant and during pregnancy, diabetes should not be treated with metformin but insulin should be used to maintain blood glucose levels as close to normal as possible in order to lower the risk of foetal malformations associated with abnormal blood glucose levels.

Lactation

Metformin is excreted into human breast milk. No adverse effects were observed in breastfed newborns/infants. However, as only limited data is available, breastfeeding is not recommended during metformin treatment. A decision on whether to discontinue breast-feeding should be made, taken into account the benefit of breast-feeding and the potential risk to adverse effects on the child.

Fertility

Fertility of male or female rats was unaffected by metformin when administered at doses as high as 600 mg/kg/day, which is approximately three times the maximum recommended human daily dose based on body surface area comparisons.

4.7 Effects on Ability to Drive and Use Machines

Metformin monotherapy does not cause hypoglycaemia and therefore has no effect on the ability to drive or to use machines. However, patients should be alerted to the risk of hypoglycaemia when metformin is used in combination with other antidiabetic agents (sulfonylureas, insulin or meglitinides).

4.8 Undesirable Effects

The following undesirable effects may occur under treatment with metformin. Frequencies are defined as follows:

very common: >1/10;
common: > 1/100; <1/10
uncommon: > 1/1,000; < 1/100;
rare: > 1/10,000, < 1/1,000
very rare: < 1/10,000, not known (cannot be estimated from the available data)

Metabolism and nutrition disorders

Very rare:

- Decrease of vitamin B12 absorption with decrease of serum levels during long-term use of metformin. Consideration of such aetiology is recommended if a patient presents with megaloblastic anaemia.
- Lactic acidosis (see section 4.4)

Nervous system disorders

Common:

- Taste disturbance

Gastrointestinal disorders:

Very common:

- Gastrointestinal disorders such as nausea, vomiting, diarrhoea, abdominal pain and loss of appetite. These undesirable effects occur most frequently during initiation of therapy and resolve spontaneously in most cases. To prevent them, it is recommended that metformin be

Sandoz	Business use only	Page 6 of 10
1.3.1 spc-label-pl - common-spc - 3,016 (NL/H/1171/001/IA/016)		20130722
METFORMIN HYDROCHLORIDE 500 MG; 850 MG FILM-COATED TABLET		721-5715.00 721-5716.00

taken in 2 or 3 daily doses during or after meals. A slow increase of the dose may also improve gastrointestinal tolerability.

Hepatobiliary disorders

Very rare:

- Isolated reports of liver function tests abnormalities or hepatitis resolving upon metformin discontinuation.

Skin and subcutaneous tissue disorders:

Very rare:

- Skin reactions such as erythema, pruritus, urticaria

Children and adolescents

In published and post marketing data and in controlled clinical studies in a limited paediatric population aged 10-16 years treated during 1 year, adverse event reporting was similar in nature and severity to that reported in adults.

4.9 Overdose

Hypoglycaemia has not been seen with metformin doses of up to 85 g, although lactic acidosis has occurred in such circumstances. High overdose of metformin or concomitant risks may lead to lactic acidosis. Lactic acidosis is a medical emergency and must be treated in hospital. The most effective method to remove lactate and metformin is haemodialysis.

5 PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic Properties

Pharmacotheapeutic group: Blood glucose lowering drugs, excl. insulins, Biguanides, ATC code: A10BA02

Metformin is a biguanide with antihyperglycaemic effects, lowering both basal and postprandial plasma glucose. It does not stimulate insulin secretion and therefore does not produce hypoglycaemia.

Metformin may act via 3 mechanisms:

1. reduction of hepatic glucose production by inhibiting gluconeogenesis and glycogenolysis
2. in muscle, by increasing insulin sensitivity, improving peripheral glucose uptake and utilisation
3. and delay of intestinal glucose absorption.

Metformin stimulates intracellular glycogen synthesis by acting on glycogen synthase. Metformin increases the transport capacity of all types of membrane glucose transporters (GLUTs) known to date.

In humans, independently of its action on glycaemia, metformin has favourable effects on lipid metabolism. This has been shown at therapeutic doses in controlled, medium-term or long-term clinical studies: metformin reduces total cholesterol, LDL cholesterol and triglyceride levels.

Clinical efficacy

The prospective randomised (UKPDS) study has established the long-term benefit of intensive blood glucose control in adult patients with type 2 diabetes.

Analysis of the results for overweight patients treated with metformin after failure of diet alone showed:

Sandoz	Business use only	Page 7 of 10
1.3.1 spc-label-pl - common-spc - 3,016 (NL/H/1171/001/IA/016)		20130722
METFORMIN HYDROCHLORIDE 500 MG; 850 MG FILM-COATED TABLET		721-5715.00 721-5716.00

- a significant reduction of the absolute risk of any diabetes-related complication in the metformin group (29.8 events/1000 patient-years) versus diet alone (43.3 events/1000 patient-years), $p=0.0023$, and versus the combined sulphonylurea and insulin monotherapy groups (40.1 events/1000 patient-years), $p=0.0034$.
- a significant reduction of the absolute risk of diabetes-related mortality: metformin 7.5 events/1000 patient-years, diet alone 12.7 events/1000 patient-years, $p=0.017$;
- a significant reduction of the absolute risk of overall mortality: metformin 13.5 events/1000 patient-years versus diet alone 20.6 events/1000 patient-years ($p=0.011$), and versus the combined sulphonylurea and insulin monotherapy groups 18.9 events/1000 patient-years ($p=0.021$);
- a significant reduction in the absolute risk of myocardial infarction: metformin 11 events/1000 patient-years, diet alone 18 events/1000 patient-years ($p=0.01$)

For metformin used as second-line therapy, in combination with a sulphonylurea, benefit regarding clinical outcome has not been shown.

In type 1 diabetes, the combination of metformin and insulin has been used in selected patients, but the clinical benefit of this combination has not been formally established.

Controlled clinical studies in a limited paediatric population aged 10-16 years with type 2 diabetes treated during one year demonstrated a similar response in glycaemic control to that seen in adults.

5.2 Pharmacokinetic Properties

Absorption:

After an oral dose of metformin, T_{max} is reached in 2.5 hours. Absolute bioavailability of a 500 mg or 850 mg metformin hydrochloride tablet is approximately 50-60% in healthy subjects. After an oral dose, the non-absorbed fraction recovered in faeces was 20-30%.

After oral administration, metformin absorption is saturable and incomplete. It is assumed that the pharmacokinetics of metformin absorption are non-linear.

At the usual metformin doses and dosing schedules, steady state plasma concentrations are reached within 24 to 48 hours and are generally less than 1 microgram/ml. In controlled clinical trials, maximum metformin plasma levels (C_{max}) did not exceed 4micrograms/ml, even at maximum doses. Food decreases the extent and slightly delays the absorption of metformin. Following administration of a dose of 850 mg, a 40% lower plasma peak concentration, a 25% decrease in AUC (area under the curve) and a 35 minute prolongation of time to peak plasma concentration were observed. The clinical relevance of these decreases is unknown.

Distribution:

Plasma protein binding is negligible. Metformin partitions into erythrocytes. The blood peak is lower than the plasma peak and appears at approximately the same time. The red blood cells most likely represent a secondary compartment of distribution. The mean V_d ranged between 63-276 l.

Metabolism:

Metformin is excreted unchanged in the urine. No metabolites have been identified in humans.

Elimination:

Renal clearance of metformin is > 400 ml/min, indicating that metformin is eliminated by glomerular filtration and tubular secretion. Following an oral dose, the apparent terminal elimination half-life is approximately 6.5 hours.

When renal function is impaired, renal clearance is decreased in proportion to that of creatinine and thus the elimination half-life is prolonged, leading to increased levels of metformin in plasma.

Sandoz	Business use only	Page 8 of 10
1.3.1 spc-label-pl - common-spc - 3,016 (NL/H/1171/001/IA/016)		20130722
METFORMIN HYDROCHLORIDE 500 MG; 850 MG FILM-COATED TABLET		721-5715.00 721-5716.00

Children and adolescents

Single dose study: After single doses of metformin hydrochloride 500 mg, paediatric patients have shown a similar pharmacokinetic profile to that observed in healthy adults.

Multiple dose study: Data are restricted to one study: After repeated doses of 500 mg BID for 7 days in paediatric patients the peak plasma concentration (C_{max}) and systemic exposure (AUC_{0-t}) were reduced by approximately 33% and 40%, respectively compared to diabetic adults who received repeated doses of 500 mg BID for 14 days. As the dose is individually titrated based on glycaemic control, this is of limited clinical relevance.

5.3 Preclinical Safety Data

Preclinical data reveal no special hazard for humans based on conventional studies on safety pharmacology, repeated dose toxicity, genotoxicity, carcinogenic potential and reproductive toxicity.

6 PHARMACEUTICAL PARTICULARS

6.1 List of Excipients

Tablet core:

Povidone K 90

Magnesium stearate

Film coating:

Hypromellose

Macrogol 4000

Titanium dioxide

6.2 Incompatibilities

Not applicable.

6.3 Shelf life

3 years

6.4 Special Precautions for Storage

No special storage conditions

Sandoz	Business use only	Page 9 of 10
1.3.1 spc-label-pl - common-spc - 3,016 (NL/H/1171/001/IA/016)		20130722
METFORMIN HYDROCHLORIDE 500 MG; 850 MG FILM-COATED TABLET		721-5715.00 721-5716.00

6.5 Nature and Contents of Container

NL/H/1170/01-02 + NL/H/1172/01-02

Metformin 500 mg:

HDPE bottles with LDPE caps with 30, 60, 100, 250, 400, 500 film-coated tablets.

PVC/ aluminium blister with 28, 30, 50, 56, 60, 84, 90, 100, 120, 180, 250, 270 film-coated tablets.

Metformin 850 mg:

HDPE bottles with LDPE caps with 30, 60, 100, 200, 250, 500 film-coated tablets.

PVC/ aluminium blister with 20, 28, 30, 40, 50, 56, 60, 84, 90, 100, 120, 180, 250, 300 film-coated tablets.

NL/H/1171/01-02

Metformin 500 mg:

HDPE bottles with LDPE caps with 30, 60, 90, 100, 250, 400 film-coated tablets.

PVC/ aluminium blister with 30, 50, 60, 84, 90, 100, 250 film-coated tablets.

Metformin 850 mg:

HDPE bottles with LDPE caps with 30, 100, 200, 250 film-coated tablets.

PVC/ aluminium blister with 30, 40, 56, 60, 100, 250 film-coated tablets.

Not all pack sizes may be marketed.

Sandoz	Business use only	Page 10 of 10
1.3.1 spc-label-pl - common-spc - 3,016 (NL/H/1171/001/IA/016)		20130722
METFORMIN HYDROCHLORIDE 500 MG; 850 MG FILM-COATED TABLET		721-5715.00 721-5716.00

6.6 Instructions for Use and Handling

No special requirements

Any unused product or waste material should be disposed of in accordance with local requirements.

7. MARKETING AUTHORIZATION HOLDER AND MANUFACTURER

Name: Sandoz GmbH, Kundl

Address: Biochemiestrasse 106250.

Country: Austria.

Manufacturer

Name: Lek S.A.

Company name: Lek S.A..

Address: ul. Podlipie 16, 95-010 Stryków

Country: Poland

8. MARKETING AUTHORIZATION NUMBER

TO BE ALLOCATED

9. DATE OF FIRST AUTHORIZATION / RENEWAL OF AUTHORIZATION

TO BE ALLOCATED

10. DATE OF REVISION OF THE TEXT

04/2014