

## **SUMMARY OF PRODUCT CHARACTERISTICS**

## **1 NAME OF THE MEDICINAL PRODUCT**

Nexium 10 mg gastro-resistant granules for oral suspension, sachet

## **2 QUALITATIVE AND QUANTITATIVE COMPOSITION**

Each sachet contains: 10 mg esomeprazole (as magnesium trihydrate).

### Excipients with known effect

Each sachet contains 6.8 mg sucrose and 2.8 g glucose

For the full list of excipients, see section 6.1.

## **3 PHARMACEUTICAL FORM**

Gastro-resistant granules for oral suspension, sachet.

Pale yellow fine granules. Brownish granules may be visible.

## **4 CLINICAL PARTICULARS**

### **4.1 Therapeutic indications**

Nexium oral suspension is primarily indicated for:

#### Paediatric population

##### Children 1-11 years old

#### *Gastroesophageal Reflux Disease (GERD)*

- treatment of endoscopically proven erosive reflux esophagitis
- symptomatic treatment of gastroesophageal reflux disease (GERD)

#### Adults and adolescents from the age of 12 years

For indications in patients from the age of 12 years reference is made to the Nexium gastro-resistant tablet SmPC.

Nexium oral suspension may also be used by patients having difficulty swallowing dispersed Nexium gastro-resistant tablets.

### **4.2 Posology and method of administration**

#### Posology

#### Paediatric population

### *Children 1 – 11 years with a bodyweight of $\geq 10$ kg*

#### *Gastroesophageal Reflux Disease (GERD)*

##### *- Treatment of endoscopically proven erosive reflux esophagitis*

- Weight  $\geq 10$  -  $< 20$  kg: 10 mg once daily for 8 weeks.
- Weight  $\geq 20$  kg: 10 mg or 20 mg once daily for 8 weeks.

##### *- Symptomatic treatment of gastroesophageal reflux disease (GERD)*

- 10 mg once daily for up to 8 weeks.

Doses over 1 mg/kg/day have not been studied.

### *Children below the age of 1 year*

The experience of treatment with esomeprazole in infants  $< 1$  year is limited and treatment is therefore not recommended (see section 5.1).

### *Adults and adolescents from the age of 12 years*

For posology in patients from the age of 12 years reference is made to the Nexium gastro-resistant tablet SmPC.

### *Special populations*

#### *Renal impairment*

Dose adjustment is not required in patients with impaired renal function. Due to limited experience in patients with severe renal insufficiency, such patients should be treated with caution (see section 5.2).

#### *Hepatic impairment*

Dose adjustment is not required in patients with mild to moderate liver impairment. For patients  $\geq 12$  years with severe liver impairment, a maximum dose of 20 mg Nexium should not be exceeded. For children 1-11 years with severe liver impairment, a maximum dose of 10 mg should not be exceeded (see section 5.2).

#### *Elderly*

Dose adjustment is not required in the elderly.

### *Method of administration*

For a 10 mg dose empty the contents of a 10 mg sachet into a glass containing 15 ml water. For a 20 mg dose empty the contents of two 10 mg sachets into a glass containing 30 ml water. Do not use carbonated water. Stir the contents until the granulate has been dispersed and leave for a few minutes to thicken. Stir again and drink within 30 minutes. The granules must not be chewed or crushed. Rinse with 15 ml water to obtain all granules.

For patients who have a nasogastric or gastric tube in place: see section 6.6 for preparation and administration instructions.

### 4.3 Contraindications

Hypersensitivity to the active substance, to substituted benzimidazoles or to any of the excipients listed in section 6.1.

Esomeprazole should not be used concomitantly with nelfinavir (see section 4.5).

### 4.4 Special warnings and precautions for use

In the presence of any alarm symptom (e.g. significant unintentional weight loss, recurrent vomiting, dysphagia, haematemesis or melaena) and when gastric ulcer is suspected or present, malignancy should be excluded, as treatment with Nexium may alleviate symptoms and delay diagnosis.

#### Long term use

Patients on long-term treatment (particularly those treated for more than a year) should be kept under regular surveillance. Long-term treatment is indicated in adults and adolescents (12 years and older, see section 4.1).

#### On demand treatment

Patients on on-demand treatment should be instructed to contact their physician if their symptoms change in character. On demand treatment has not been investigated in children and is therefore not recommended in this patient group.

#### Gastrointestinal infections

Treatment with proton pump inhibitors may lead to a slightly increased risk of gastrointestinal infections such as *Salmonella* and *Campylobacter* (see section 5.1).

#### Absorption of vitamin B12

Esomeprazole, as all acid-blocking medicines, may reduce the absorption of vitamin B12 (cyanocobalamin) due to hypo- or achlorhydria. This should be considered in patients with reduced body stores or risk factors for reduced vitamin B12 absorption on long-term therapy.

#### Hypomagnesaemia

Severe hypomagnesaemia has been reported in patients treated with proton pump inhibitors (PPIs) like esomeprazole for at least three months, and in most cases for a year. Serious manifestations of hypomagnesaemia such as fatigue, tetany, delirium, convulsions, dizziness and ventricular arrhythmia can occur but they may begin insidiously and be overlooked. In most affected patients, hypomagnesaemia improved after magnesium replacement and discontinuation of the PPI.

For patients expected to be on prolonged treatment or who take PPIs with digoxin or medicinal products that may cause hypomagnesaemia (e.g. diuretics), healthcare professionals should consider measuring magnesium levels before starting PPI treatment and periodically during treatment.

#### Risk of fractures

Proton pump inhibitors, especially if used in high doses and over long durations (>1 year), may modestly increase the risk of hip, wrist and spine fracture,

predominantly in the elderly or in presence of other recognised risk factors. Observational studies suggest that proton pump inhibitors may increase the overall risk of fracture by 10-40%. Some of this increase may be due to other risk factors. Patients at risk of osteoporosis should receive care according to current clinical guidelines and they should have an adequate intake of vitamin D and calcium.

#### Subacute cutaneous lupus erythematosus (SCLE)

Proton pump inhibitors are associated with very infrequent cases of SCLE. If lesions occur, especially in sun-exposed areas of the skin, and if accompanied by arthralgia, the patient should seek medical help promptly and the health care professional should consider stopping Nexium. SCLE after previous treatment with a proton pump inhibitor may increase the risk of SCLE with other proton pump inhibitors.

#### Combination with other medicinal products

Co-administration of esomeprazole with atazanavir is not recommended (see section 4.5). If the combination of atazanavir with a proton pump inhibitor is judged unavoidable, close clinical monitoring is recommended in combination with an increase in the dose of atazanavir to 400 mg with 100 mg of ritonavir; esomeprazole 20 mg should not be exceeded.

Esomeprazole is a CYP2C19 inhibitor. When starting or ending treatment with esomeprazole, the potential for interactions with medicinal products metabolised through CYP2C19 should be considered. An interaction is observed between clopidogrel and esomeprazole (see section 4.5). The clinical relevance of this interaction is uncertain. As a precaution, concomitant use of esomeprazole and clopidogrel should be discouraged.

When prescribing esomeprazole for on demand therapy, the implications for interactions with other pharmaceuticals, due to fluctuating plasma concentrations of esomeprazole should be considered (see section 4.5).

#### Serious cutaneous adverse reactions (SCARs)

Serious cutaneous adverse reactions (SCARs) such as erythema multiforme (EM), Stevens-Johnson syndrome (SJS), toxic epidermal necrolysis (TEN) and drug reaction with eosinophilia and systemic symptoms (DRESS), which can be life-threatening, have been reported very rarely in association with esomeprazole treatment.

Patients should be advised of the signs and symptoms of the severe skin reaction EM/SJS/TEN/DRESS and should seek medical advice from their physician immediately when observing any indicative signs or symptoms.

Esomeprazole should be discontinued immediately upon signs and symptoms of severe skin reactions and additional medical care/close monitoring should be provided as needed.

Re-challenge should not be undertaken in patients with EM/SJS/TEN/DRESS.

#### Sucrose and glucose

This medicinal product contains sucrose and glucose. Patients with rare hereditary problems of fructose intolerance, glucose-galactose malabsorption or sucrase-isomaltase insufficiency should not take this medicine.

#### Interference with laboratory tests

Increased Chromogranin A (CgA) level may interfere with investigations for neuroendocrine tumours. To avoid this interference, esomeprazole treatment should be stopped for at least 5 days before CgA measurements (see section 5.1). If CgA and gastrin levels have not returned to reference range after initial measurement, measurements should be repeated 14 days after cessation of proton pump inhibitor treatment.

### **4.5 Interaction with other medicinal products and other forms of interaction**

#### Effects of esomeprazole on the pharmacokinetics of other medicinal products

##### *Protease inhibitors*

Omeprazole has been reported to interact with some protease inhibitors. The clinical importance and the mechanisms behind these reported interactions are not always known. Increased gastric pH during omeprazole treatment may change the absorption of the protease inhibitors. Other possible interaction mechanisms are via inhibition of CYP2C19.

For atazanavir and nelfinavir, decreased serum levels have been reported when given together with omeprazole and concomitant administration is not recommended. Co-administration of omeprazole (40 mg once daily) with atazanavir 300 mg/ritonavir 100 mg to healthy volunteers resulted in a substantial reduction in atazanavir exposure (approximately 75% decrease in AUC,  $C_{\max}$  and  $C_{\min}$ ). Increasing the atazanavir dose to 400 mg did not compensate for the impact of omeprazole on atazanavir exposure. The co-administration of omeprazole (20 mg qd) with atazanavir 400 mg/ritonavir 100 mg to healthy volunteers resulted in a decrease of approximately 30% in the atazanavir exposure as compared with the exposure observed with atazanavir 300 mg/ritonavir 100 mg qd without omeprazole 20 mg qd. Co-administration of omeprazole (40 mg qd) reduced mean nelfinavir AUC,  $C_{\max}$  and  $C_{\min}$  by 36-39% and mean AUC,  $C_{\max}$  and  $C_{\min}$  for the pharmacologically active metabolite M8 was reduced by 75-92%. Due to the similar pharmacodynamic effects and pharmacokinetic properties of omeprazole and esomeprazole, concomitant administration with esomeprazole and atazanavir is not recommended (see section 4.4) and concomitant administration with esomeprazole and nelfinavir is contraindicated (see section 4.3).

For saquinavir (with concomitant ritonavir), increased serum levels (80-100%) have been reported during concomitant omeprazole treatment (40 mg qd). Treatment with omeprazole 20 mg qd had no effect on the exposure of darunavir (with concomitant ritonavir) and amprenavir (with concomitant ritonavir). Treatment with esomeprazole 20 mg qd had no effect on the exposure of amprenavir (with and without concomitant ritonavir). Treatment with omeprazole 40 mg qd had no effect on the exposure of lopinavir (with concomitant ritonavir)

##### *Methotrexate*

When given together with PPIs, methotrexate levels have been reported to increase in some patients. In high-dose methotrexate administration a temporary withdrawal of esomeprazole may need to be considered.

#### *Tacrolimus*

Concomitant administration of esomeprazole has been reported to increase the serum levels of tacrolimus.

#### *Medicinal products with pH dependent absorption*

Gastric acid suppression during treatment with esomeprazole and other PPIs might decrease or increase the absorption of medicinal products with a gastric pH dependent absorption. As with other medicinal products that decrease intragastric acidity, the absorption of medicinal products such as ketoconazole, itraconazole and erlotinib can decrease and the absorption of digoxin can increase during treatment with esomeprazole. Concomitant treatment with omeprazole (20 mg daily) and digoxin in healthy subjects increased the bioavailability of digoxin by 10% (up to 30% in two out of ten subjects). Digoxin toxicity has been rarely reported. However, caution should be exercised when esomeprazole is given at high doses in elderly patients. Therapeutic drug monitoring of digoxin should then be reinforced.

#### *Medicinal products metabolised by CYP2C19*

Esomeprazole inhibits CYP2C19, the major esomeprazole metabolising enzyme. Thus, when esomeprazole is combined with medicinal products metabolised by CYP2C19, such as diazepam, citalopram, imipramine, clomipramine, phenytoin etc., the plasma concentrations of these medicinal products may be increased and a dose reduction could be needed. This should be considered especially when prescribing esomeprazole for on demand therapy.

#### *Diazepam*

Concomitant administration of 30 mg esomeprazole resulted in a 45% decrease in clearance of the CYP2C19 substrate diazepam.

#### *Phenytoin*

Concomitant administration of 40 mg esomeprazole resulted in a 13% increase in trough plasma levels of phenytoin in epileptic patients. It is recommended to monitor the plasma concentrations of phenytoin when treatment with esomeprazole is introduced or withdrawn.

#### *Voriconazole*

Omeprazole (40 mg once daily) increased voriconazole (a CYP2C19 substrate)  $C_{max}$  and  $AUC_{\tau}$  by 15% and 41%, respectively.

#### *Cilostazol*

Omeprazole as well as esomeprazole act as inhibitors of CYP2C19. Omeprazole, given in doses of 40 mg to healthy subjects in a cross-over study, increased  $C_{max}$  and  $AUC$  for cilostazol by 18% and 26% respectively, and one of its active metabolites by 29% and 69% respectively.

#### *Cisapride*

In healthy volunteers, concomitant administration of 40 mg esomeprazole and cisapride resulted in a 32% increase in area under the plasma concentration-time curve (AUC) and a 31% prolongation of elimination half-life( $t_{1/2}$ ) but no significant increase in peak plasma levels of cisapride. The slightly prolonged QTc interval observed after administration of cisapride alone, was not further prolonged when cisapride was given in combination with esomeprazole (see section 4.4).

#### *Warfarin*

Concomitant administration of 40 mg esomeprazole to warfarin-treated patients in a clinical trial showed that coagulation times were within the accepted range. However, post-marketing, a few isolated cases of elevated INR of clinical significance have been reported during concomitant treatment. Monitoring is recommended when initiating and ending concomitant esomeprazole treatment, during treatment with warfarin or other coumarine derivatives.

#### *Clopidogrel*

Results from studies in healthy subjects have shown a pharmacokinetic (PK)/pharmacodynamic (PD) interaction between clopidogrel (300 mg loading dose/75 mg daily maintenance dose) and esomeprazole (40 mg p.o.daily) resulting in decreased exposure to the active metabolite of clopidogrel by an average of 40% and resulting in decreased maximum inhibition of (ADP induced) platelet aggregation by an average of 14%.

When clopidogrel was given together with a fixed dose combination of esomeprazole 20 mg +ASA 81 mg compared to clopidogrel alone in a study in healthy subjects there was a decreased exposure by almost 40% of the active metabolite of clopidogrel. However, the maximum levels of inhibition of (ADP induced) platelet aggregation in these subjects were the same in the clopidogrel and the clopidogrel + the combined (esomeprazole + ASA) product groups.

Inconsistent data on the clinical implications of a PK/PD interaction of esomeprazole in terms of major cardiovascular events have been reported from both observational and clinical studies. As a precaution concomitant use of clopidogrel should be discouraged.

#### *Investigated medicinal products with no clinically relevant interaction*

##### *Amoxicillin and quinidine*

Esomeprazole has been shown to have no clinically relevant effects on the pharmacokinetics of amoxicillin or quinidine.

##### *Naproxen or rofecoxib*

Studies evaluating concomitant administration of esomeprazole and either naproxen or rofecoxib did not identify any clinically relevant pharmacokinetic interactions during short-term studies.

#### Effects of other medicinal products on the pharmacokinetics of esomeprazole

##### *Medicinal products which inhibit CYP2C19 and/or CYP3A4*

Esomeprazole is metabolised by CYP2C19 and CYP3A4. Concomitant administration of esomeprazole and a CYP3A4 inhibitor, clarithromycin (500 mg b.i.d.), resulted in a



doubling of the exposure (AUC) to esomeprazole. Concomitant administration of esomeprazole and a combined inhibitor of CYP2C19 and CYP 3A4 may result in more than doubling of the esomeprazole exposure. The CYP2C19 and CYP3A4 inhibitor voriconazole increased omeprazole AUC<sub>t</sub> by 280%. A dose adjustment of esomeprazole is not regularly required in either of these situations. However, dose adjustment should be considered in patients with severe hepatic impairment and if long-term treatment is indicated. Long-term treatment is indicated in adults and adolescents (12 years and older, see section 4.1).

#### *Medicinal products which induce CYP2C19 and/or CYP3A4*

Medicinal products known to induce CYP2C19 or CYP3A4 or both (such as rifampicin and St. John's wort) may lead to decreased esomeprazole serum levels by increasing the esomeprazole metabolism.

#### Paediatric population

Interaction studies have only been performed in adults.

### **4.6 Fertility, pregnancy and lactation**

#### Pregnancy

Clinical data on exposed pregnancies with Nexium are insufficient. With the racemic mixture omeprazole, data on a larger number of exposed pregnancies from epidemiological studies indicate no malformative nor foetotoxic effect. Animal studies with esomeprazole do not indicate direct or indirect harmful effects with respect to embryonal/foetal development. Animal studies with the racemic mixture do not indicate direct or indirect harmful effects with respect to pregnancy, parturition or postnatal development. Caution should be exercised when prescribing to pregnant women.

#### Breast-feeding

It is not known whether esomeprazole is excreted in human breast milk. There is insufficient information on the effects of esomeprazole in newborns/infants. Esomeprazole should not be used during breast-feeding.

### **4.7 Effects on ability to drive and use machines**

Esomeprazole has minor influence on the ability to drive and use machines. Adverse reactions such as dizziness (uncommon) and blurred vision (rare) has been reported (see section 4.8). If affected patients should not drive or use machines.

### **4.8 Undesirable effects**

#### Summary of the safety profile

Headache, abdominal pain, diarrhoea and nausea are among those adverse reactions that have been most commonly reported in clinical trials (and also from post-marketing use). In addition, the safety profile is similar for different formulations, treatment indications, age groups and patient populations. No dose-related adverse reactions have been identified.

### Tabulated list of adverse reactions

The following adverse drug reactions have been identified or suspected in the clinical trials programme for esomeprazole and post-marketing. None was found to be dose-related. The reactions are classified according to frequency: very common  $\geq 1/10$ ; common  $\geq 1/100$  to  $< 1/10$ ; uncommon  $\geq 1/1,000$  to  $< 1/100$ ; rare  $\geq 1/10,000$  to  $< 1/1,000$ ; very rare  $< 1/10,000$ ; not known (cannot be estimated from the available data).

<b>System Organ Class</b>	<b>Frequency</b>	<b>Undesirable Effect</b>
Blood and lymphatic system disorders	Rare	Leukopenia, thrombocytopenia
	Very rare	Agranulocytosis, pancytopenia
Immune system disorders	Rare	Hypersensitivity reactions e.g. fever, angioedema and anaphylactic reaction/shock
Metabolism and nutrition disorders	Uncommon	Peripheral oedema
	Rare	Hyponatraemia
	Not known	Hypomagnesaemia (see section 4.4); severe hypomagnesaemia can correlate with hypocalcaemia. Hypomagnesaemia may also be associated with hypokalaemia.
Psychiatric disorders	Uncommon	Insomnia
	Rare	Agitation, confusion, depression
	Very rare	Aggression, hallucinations
Nervous system disorders	Common	Headache
	Uncommon	Dizziness, paraesthesia, somnolence
	Rare	Taste disturbance
Eye disorders	Rare	Blurred vision
Ear and labyrinth disorders	Uncommon	Vertigo
Respiratory, thoracic and mediastinal disorders	Rare	Bronchospasm
Gastrointestinal disorders	Common	Abdominal pain, constipation, diarrhoea, flatulence, nausea/vomiting, fundic gland polyps (benign)
	Uncommon	Dry mouth
	Rare	Stomatitis, gastrointestinal candidiasis
	Not known	Microscopic colitis
Hepatobiliary disorders	Uncommon	Increased liver enzymes
	Rare	Hepatitis with or without jaundice

System Organ Class	Frequency	Undesirable Effect
	Very rare	Hepatic failure, encephalopathy in patients with pre-existing liver disease
Skin and subcutaneous tissue disorders	Uncommon	Dermatitis, pruritus, rash, urticaria
	Rare	Alopecia, photosensitivity
	Very rare	Erythema multiforme, Stevens-Johnson syndrome, toxic epidermal necrolysis (TEN), drug reaction with eosinophilia and systemic symptoms (DRESS)
	Not known	Subacute cutaneous lupus erythematosus (see section 4.4)
Musculoskeletal and connective tissue disorders	Uncommon	Fracture of the hip, wrist or spine (see section 4.4)
	Rare	Arthralgia, myalgia
	Very rare	Muscular weakness
Renal and urinary disorders	Very rare	Interstitial nephritis; in some patients renal failure has been reported concomitantly.
Reproductive system and breast disorders	Very rare	Gynaecomastia
General disorders and administration site conditions	Rare	Malaise, increased sweating

#### Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product.

## **4.9 Overdose**

There is very limited experience to date with deliberate overdose. The symptoms described in connection with 280 mg were gastrointestinal symptoms and weakness. Single doses of 80 mg esomeprazole were uneventful. No specific antidote is known. Esomeprazole is extensively plasma protein bound and is therefore not readily dialyzable. As in any case of overdose, treatment should be symptomatic and general supportive measures should be utilised.

## **5 PHARMACOLOGICAL PROPERTIES**

### **5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: Drugs for acid-related disorders, proton pump inhibitor  
ATC code: A02B C05

Esomeprazole is the S-isomer of omeprazole and reduces gastric acid secretion through a specific targeted mechanism of action. It is a specific inhibitor of the acid pump in the parietal cell. Both the R- and S-isomer of omeprazole have similar pharmacodynamic activity.

#### Mechanism of action

Esomeprazole is a weak base and is concentrated and converted to the active form in the highly acidic environment of the secretory canaliculi of the parietal cell, where it inhibits the enzyme  $H^+K^+-ATPase$  – the acid pump and inhibits both basal and stimulated acid secretion.

#### Pharmacodynamic effects

After oral dosing with esomeprazole 20 mg and 40 mg the onset of effect occurs within one hour. After repeated administration with 20 mg esomeprazole once daily for five days, mean peak acid output after pentagastrin stimulation is decreased 90% when measured 6 – 7 hours after dosing on day five.

After five days of oral dosing with 20 mg and 40 mg of esomeprazole, intragastric pH above 4 was maintained for a mean time of 13 hours and 17 hours, respectively over 24 hours in symptomatic GERD patients. The proportion of patients maintaining an intragastric pH above 4 for at least 8, 12 and 16 hours respectively were for esomeprazole 20 mg 76%, 54% and 24%. Corresponding proportions for esomeprazole 40 mg were 97%, 92% and 56%.

Using AUC as a surrogate parameter for plasma concentration, a relationship between inhibition of acid secretion and exposure has been shown.

Healing of reflux esophagitis with esomeprazole 40 mg occurs in approximately 78% of patients after four weeks, and in 93% after eight weeks.

During treatment with antisecretory medicinal products, serum gastrin increases in response to the decreased acid secretion. Also CgA increases due to decreased gastric acidity. The increased CgA level may interfere with investigations for neuroendocrine tumours. Available published evidence suggests that proton pump inhibitors should be discontinued between 5 days and 2 weeks prior to CgA measurements. This is to allow CgA levels that might be spuriously elevated following PPI treatment to return to reference range.

An increased number of ECL cells possibly related to the increased serum gastrin levels, have been observed in both children and adults during long term treatment with esomeprazole. The findings are considered to be of no clinical significance.

During long-term treatment with antisecretory medicinal products gastric glandular cysts have been reported to occur at a somewhat increased frequency. These changes are a physiological consequence of pronounced inhibition of acid secretion, are benign and appear to be reversible.

Decreased gastric acidity due to any means including proton pump inhibitors, increases gastric count of bacteria normally present in the gastrointestinal tract.

Treatment with proton pump inhibitors may lead to a slightly increased risk of gastrointestinal infections such as *Salmonella* and *Campylobacter* and, in hospitalised patients, possibly also *Clostridium difficile*.

### Paediatric population

#### *GERD – 1 to 11 Years of Age*

In a multicentre, parallel-group study, 109 paediatric patients with endoscopically proven GERD (1 to 11 years of age) were treated with Nexium once daily for up to 8 weeks to evaluate safety and tolerability. Dosing by patient weight was as follows:

Weight <20 kg: once daily treatment with esomeprazole 5 mg or 10 mg  
Weight ≥20 kg: once daily treatment with esomeprazole 10 mg or 20 mg

Patients were endoscopically characterised as to the presence or absence of erosive esophagitis. Fifty-three patients had erosive esophagitis at baseline. Of the 45 patients who had follow-up endoscopy, 42 (93.3%) of these patients had their erosive esophagitis resolved (88.9%) or improved (4.4%) after 8 weeks of treatment.

#### *GERD – 0 to 11 months of age*

In a placebo-controlled study (98 patients aged 1-11 months) efficacy and safety in patients with signs and symptoms of GERD were evaluated. Esomeprazole 1 mg/kg once daily was given for 2 weeks (open-label phase) and 80 patients were included for an additional 4 weeks (double blind, treatment-withdrawal phase). There was no significant difference between esomeprazole and placebo for the primary endpoint time to discontinuation due to symptom worsening.

In a placebo-controlled study (52 patients aged <1 month) efficacy and safety in patients with symptoms of GERD were evaluated. Esomeprazole 0.5 mg/kg once daily was given for a minimum of 10 days. There was no significant difference between esomeprazole and placebo in the primary endpoint, change from baseline of number of occurrences of symptoms of GERD.

Results from the paediatric studies further show that 0.5 mg/kg and 1.0 mg/kg esomeprazole in <1 month old and 1 to 11 month old infants, respectively, reduced the mean percentage of time with intra-esophageal pH < 4.

The safety profile appeared to be similar to that seen in adults.

In a study in paediatric GERD patients (<1 to 17 years of age) receiving long-term PPI treatment, 61% of the children developed minor degrees of ECL cell hyperplasia with no known clinical significance and with no development of atrophic gastritis or carcinoid tumours.

## **5.2 Pharmacokinetic properties**

### Absorption

Esomeprazole is acid labile and is administered orally as enteric-coated granules. In vivo conversion to the R-isomer is negligible. Absorption of esomeprazole is rapid, with peak plasma levels occurring approximately 1-2 hours after dose. The absolute

bioavailability is 64% after a single dose of 40 mg and increases to 89% after repeated once-daily administration. For 20 mg esomeprazole the corresponding values are 50% and 68%, respectively. Food intake both delays and decreases the absorption of esomeprazole although this has no significant influence on the effect of esomeprazole on intragastric acidity.

#### Distribution

The apparent volume of distribution at steady state in healthy subjects is approximately 0.22 l/kg body weight. Esomeprazole is 97% plasma protein bound.

#### Biotransformation

Esomeprazole is completely metabolised by the cytochrome P450 system (CYP). The major part of the metabolism of esomeprazole is dependent on the polymorphic CYP2C19, responsible for the formation of the hydroxy- and desmethyl metabolites of esomeprazole. The remaining part is dependent on another specific isoform, CYP3A4, responsible for the formation of esomeprazole sulphone, the main metabolite in plasma.

#### Elimination

The parameters below reflect mainly the pharmacokinetics in individuals with a functional CYP2C19 enzyme, extensive metabolisers.

Total plasma clearance is about 17 l/h after a single dose and about 9 l/h after repeated administration. The plasma elimination half-life is about 1.3 hours after repeated once-daily dosing. Esomeprazole is completely eliminated from plasma between doses with no tendency for accumulation during once-daily administration.

The major metabolites of esomeprazole have no effect on gastric acid secretion. Almost 80% of an oral dose of esomeprazole is excreted as metabolites in the urine, the remainder in the faeces. Less than 1% of the parent drug is found in urine.

#### Linearity/non linearity

The pharmacokinetics of esomeprazole has been studied in doses up to 40 mg b.i.d. The area under the plasma concentration-time curve increases with repeated administration of esomeprazole. This increase is dose-dependent and results in a more than dose proportional increase in AUC after repeated administration. This time - and dose-dependency is due to a decrease of first pass metabolism and systemic clearance probably caused by an inhibition of the CYP2C19 enzyme by esomeprazole and/or its sulphone metabolite.

#### Special patient populations

##### Poor metabolisers

Approximately  $2.9 \pm 1.5\%$  of the population lack a functional CYP2C19 enzyme and are called poor metabolisers. In these individuals the metabolism of esomeprazole is probably mainly catalysed by CYP3A4. After repeated once-daily administration of 40 mg esomeprazole, the mean area under the plasma concentration-time curve was approximately 100% higher in poor metabolisers than in subjects having a functional CYP2C19 enzyme (extensive metabolisers). Mean peak plasma concentrations were increased by about 60%. These findings have no implications for the posology of esomeprazole.

### Gender

Following a single dose of 40 mg esomeprazole the mean area under the plasma concentration-time curve is approximately 30% higher in females than in males. No gender difference is seen after repeated once-daily administration. These findings have no implications for the posology of esomeprazole.

### Hepatic impairment

The metabolism of esomeprazole in patients with mild to moderate liver dysfunction may be impaired. The metabolic rate is decreased in patients with severe liver dysfunction resulting in a doubling of the area under the plasma concentration-time curve of esomeprazole. Therefore, a maximum of 20 mg should not be exceeded in patients with severe dysfunction. Esomeprazole or its major metabolites do not show any tendency to accumulate with once-daily dosing.

### Renal impairment

No studies have been performed in patients with decreased renal function. Since the kidney is responsible for the excretion of the metabolites of esomeprazole but not for the elimination of the parent compound, the metabolism of esomeprazole is not expected to be changed in patients with impaired renal function.

### Elderly

The metabolism of esomeprazole is not significantly changed in elderly subjects (71–80 years of age).

### Paediatric population

#### *Adolescents 12-18 years:*

Following repeated dose administration of 20 mg and 40 mg esomeprazole in adolescents 12-18 years of age, the total exposure (AUC) and the time to reach maximum plasma drug concentration ( $t_{max}$ ) was similar to that in adults.

#### *Children 1 – 11 years:*

Following repeated dose administration of 10 mg esomeprazole, the total exposure (AUC) was similar within the age range 1 to 11 years and the exposure was similar to the exposure seen with the 20 mg dose in adolescents and adults. Following repeated dose administration of 20 mg esomeprazole, the total exposure (AUC) was higher in 6 to 11 year-olds compared to the same dose in adolescents and adults.

## **6 PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

#### *Esomeprazole granules:*

Glycerol monostearate 40-55,

Hydroxypropyl cellulose

Hypromellose

Magnesium stearate

Methacrylic acid –ethyl acrylate copolymer (1:1) dispersion 30%

Polysorbate 80

Sugar spheres (sucrose and maize starch)  
Talc  
Triethyl citrate

*Excipient granules:*

Citric acid anhydrous (for pH adjustment)  
Crospovidone  
Glucose  
Hydroxypropyl cellulose  
Yellow iron oxide (E172)  
Xanthan gum

## **6.2 Incompatibilities**

Not applicable

## **6.3 Shelf life**

3 years  
To be used within 30 minutes after reconstitution.

## **6.4 Special precautions for disposal and other handling**

Do not store above 30°C.

## **6.5 Nature and contents of container**

Carton containing 28 or 30 sachets. Not all pack sizes may be marketed.

Sachets (containing granules): Laminate consisting of three layers: polyethylene terephthalate (PET), aluminium, low density polyethylene (LDPE) which protects the granules against moisture.

## **6.6 Special precautions for disposal and other handling**

No special requirements for disposal.

### For patients who have a nasogastric or gastric tube in place

1. For a 10 mg dose, add the contents of a 10 mg sachet into 15 ml of water.
2. For a 20 mg dose add the contents of two 10 mg sachets into 30 ml of water.
3. Stir.
4. Leave for a few minutes to thicken.
5. Stir again.
6. Draw the suspension into a syringe.
7. Inject through the enteric tube, French size 6 or larger, into the stomach within 30 minutes after reconstitution.
8. Refill the syringe with 15 ml water for a 10 mg dose and 30 ml for a 20 mg dose.
9. Shake and flush any remaining contents from the enteric tube into the stomach.
10. Any unused suspension should be discarded.



**7      MARKETING AUTHORISATION HOLDER**

AstraZeneca UK Limited  
Silk Road Business Park, Macclesfield, Cheshire SK10 2NA  
United Kingdom

**8      MARKETING AUTHORISATION NUMBER(S)**

Rwanda FDA-HMP-MA-0287

**9      DATE OF FIRST AUTHORISATION/RENEWAL OF THE  
AUTHORISATION**

Date of first authorisation: 12 May 2023

**10     DATE OF REVISION OF THE TEXT**

17 December 2021