



## LEAFLET

# AGOCIM-200

(CIMETIDINE TABLETS BP 200 MG)

### Composition :

Each film coated tablet contains :  
Cimetidine BP 200 mg

**Category:** Histamine H<sub>2</sub> Receptor Antagonist

### Clinical Pharmacology :

It competitively inhibits histamine induced gastric secretion. All phases (basal, psychic, neurogenic, gastric) of secretion are suppressed whereas it has no effect on gastric and esophageal motility.

### Indications :

Duodenal, gastric, recurrent and stomal ulceration. Oesophageal reflux diseases and other conditions where a reduction of gastric acid is beneficial including persistent dyspeptic symptoms. Zollinger-Ellison syndrome. Stress ulcers and gastritis.

### Dosage and Administration (directions for use) :

Benign gastric ulcer: 400 mg twice at breakfast bedtime.

Acute duodenal ulcer: 800 mg at bed time.

Prophylaxis of recurrent duodenal ulcer maintain dose of 400 mg at bedtime

Peptic oesophagitis : 0.8-1.6 mg daily.-1.6 mg daily..

Persistent dyspeptic symptoms: 200 mg 4 time daily. Max . period of treatment should be 4 weeks. 800-1600 mg/day depending on severity of symptoms, has to be given in 4 divided doses. Maintain 400 mg at bedtime.

### Precautions & Warnings :

Exclude presence of gastric malignancy. Pregnancy, lactation, Impaired renal function i.v. Injection by slow infusion

**Contraindications :** Hypersensitivity

### Adverse reactions / Side effects :

Headache, dizziness, bowel upset ash diarrhoea, confusion hallucinations, delirium, convulsion, Gynaecomastia, Sodium and water retention. Impotence.