



## Literature

# AGOS CAB APPLICATION

( Benzyl Benzoate Application BP 25 % w/v )

### Composition :

Each ml contains:

Benzyl Benzoate BP 250 mg

Flavoured Aqueous base q.s.

It is an oily liquid with a faint aromatic smell and is a drug of choice for scabies. It is available as an emulsion which is applied all over except face & neck after cleansing bath. For pediculosis the emulsion is applied at the site of infestation taking care not to get into eyes. It is washed off with soap and water after 24 hours.

### Indication:

- Scabies is a skin infestation caused by a microscopic mite. The mite burrows under the skin and causes severe itching.
- Pediculosis is an infestation of the head, body or pubic area by lice.
- Agoscab application kills both mites and lice.

### Dosage & Administration:

#### Adults:

Scabies (Mites)

After a hot bath and drying, Shake the bottle apply the emulsion to the whole body except the face and head. You may find it easier to apply using a paintbrush. Special attention should be paid to:

- Wrists, ankles, armpits, buttocks and groin area
- Webs of the fingers and toes and under nails

Then allow the emulsion to dry, which usually takes about 10 minutes in a warm room.

It is desirable to put on clean underclothing and to have clean night garments and bed linen. Discarded clothes and linen are effectively disinfected by the ordinary process of washing and ironing.

If the application is thorough, one treatment is normally enough but a second application within five days will reduce the chance of any infestation remaining. Alternatively, Agoscab application can be applied to the whole body, except the head and face, on three occasions at 12 hourly intervals.

You should have a hot bath 12 hours after the last application and change to clean clothes and bed linen. Discarded clothes and bed linen should be washed.

#### Pediculosis (Lice)

The affected area should be coated with the emulsion, as above. Wash off after 24 hours with soap and water. In severe cases this procedure may need to be repeated two or three times.

A further examination should be carried out a week after the last application to confirm that the lice have been killed.

#### Children and babies:

Dilute the emulsion before application with an equal quantity of water for older children and with 3 parts of water for babies

**Contraindications:** Irritated skin, neonates



**AGOG Pharma Ltd.**



(WHO - GMP CERTIFIED - GOVT RECOGNISED EXPORT HOUSE)

CIN: 11-09015

Regd. Office & Factory : Plot No. 33, Sector II, The Vasai Taluka Industrial Co-op. Estate Ltd. Gauraipada, Vasai (E), Dist. Thane - 401 208. INDIA.  
Tel. : 95250 - 2455801 / 2452714 / 2453525 • Fax : 95250 - 2452074 (0091 - 250 - 2452074) • Email : agog@vsnl.net & agogpharma@rediffmail.com

**Special Precaution :**

Wash hands after use. Avoid use on broken skin, the eyes, nose & lips, breast-feeding (suspend feeding until product has been washed off).

**Drug Interaction:**

Although certain medicines should not be used together at all, in other cases two different medicines may be used together even if an interaction might occur. In these cases, physician may want to change the dose, or other precautions may be necessary.

**Other Interactions:**

Certain medicines should not be used at or around the time of eating food or eating certain types of food since interactions may occur. Using alcohol or tobacco with certain medicines may also cause interactions to occur.

**Side effects:**

Agoscab application may cause mild skin irritation or a short lived burning sensation.

This side effect may be more severe in sensitive patients. If the effect is severe, the emulsion should be washed off immediately with soap and water.

Excessive application of the emulsion has been reported to cause:

- Difficulty in passing urine
- Fits or convulsions in infants

If either of these occur, wash off the emulsion and contact your physician immediately. All medicines may have unwanted effects which are not mentioned in the product leaflet. If notice any other changes in health whilst taking this medicine, consult the physician immediately .

**Directions:**

As directed by the physician.

Protect from light.

Keep out of reach of children .

Shake well before use.

**Storage :**

Store under normal storage conditions (15° C to 30°C) .

**Shelf life :** 36 months

**Presentation :**

Pet bottle of 100 ml

I 552/01/01



Manufactured in India by:

**AGOG PHARMA LTD.**

Plot No. 33, Sector II, The Vasai Taluka Indl. Co-op. Estate Ltd., Vasai (E), Dist. Thane. INDIA.