



AGOG Pharma Ltd.



(WHO - GMP CERTIFIED - GOVT RECOGNISED EXPORT HOUSE)

Regd. Office & Factory : Plot No. 33, Sector II, The Vasai Taluka Industrial Co-op. Estate Ltd. Gauraipada, Vasai (E), Dist. Thane - 401 208. INDIA.
Tel. : 95250 - 2455801 / 2452714 / 2453525 • Fax : 95250 - 2452074 (0091 - 250 - 2452074) • Email : agog@vsnl.net & agogpharma@rediffmail.com

LEAFLET

Agovir
Aciclovir Tablets BP

Appearance : White round flat uncoated tablet having embossed A/200 on one side and 'AGOG' on other side.

Composition :
Each uncoated tablet contains :
Aciclovir BP 200 mg

Clinical Pharmacology : It is deoxiguanosine analogue and requires a virus specific enzyme for conversion to the active metabolite that inhibits DNA synthesis and viral replication.

Indications : Genital herpes simplex, Musocutaneous, H. Simplex, H. simplex keratitis, Herpes zoster, chicken pox.

Dosage and Administration (direction for use) :
Adults : Herpes simplex: 200 mg 5 times daily at 4 hrs interval for 5 days. Herpes zoster: 800 mg 5 times daily at 4 hrs. interval for 7 days. Prophylaxis or suppression; 200 mg 4 times daily.
Children : below 2 yrs : Herpes simplex: 100 mg 5 times at 4 hrs. interval. Children above 2 yrs dose same as adult, Prophylaxis : for children below 2 yrs: 100 mg 4 times daily at 6 hrs intervals and over 2 yrs: same as adult.



AGOG Pharma Ltd.



(WHO - GMP CERTIFIED - GOVT RECOGNISED EXPORT HOUSE)

Regd. Office & Factory : Plot No. 33, Sector II, The Vasai Taluka Industrial Co-op. Estate Ltd. Gaurai-pada, Vasai (E), Dist. Thane - 401 208. INDIA.
Tel. : 95250 - 2455801 / 2452714 / 2453525 • Fax : 95250 - 2452074 (0091 - 250 - 2452074) • Email : agog@vsnl.net & agogpharma@rediffmail.com

Precautions & Warnings :

Renal Impairment, necessitates dose reduction, Maintain adequate hydration.

Paediatrics : Reduced dose necessary. Not recommended in age below 2 years.

Pregnancy : Contraindicated orally and parentally.

Lactation : may reduce lactation.

Contraindications : Hypersensitivity, glaucoma, psychiatric diseases depression

Adverse reactions / Side effects : Orally causes headache, nausea, CNS effects. Intravenously causes sweating, emesis and fall in BP, dose dependent decrease in G. F.R. reversible neurological manifestations.

Drug Interactions : Probenecid decreases the renal tubular secretion of aciclovir leading to increased concentration in serum, prolonged half life and increased toxicity.

Nephrotoxic agents like paracetamol, aminoglycosides amphotericin B, lithium, methotrexate and methoxyflurane may increase the potential for nephrotoxicity of aciclovir when concurrently administered.

Presentation : 10 x 10 Tablets Blister in a printed carton.
1 x 10 Tablets Blister in a printed carton
100 Tablets in a HDPE jar,
1000 Tablets in a HDPE jar.



Manufactured in India by :
AGOG PHARMA LTD.
Plot No.33, Sector II, The Vasai Taluka Indl.
Co-op.Estate Ltd., Vasai (E), Dist. Thane, INDIA.