


SAME SIZE ARTWORK
LEAFLET SIZE: 140 mm x 88 mm



syphilitic skin lesions and in cases of hypersensitivity to any of the components.

Storage:
Store below 30°C. Protect from light.

For the use only of Registered Medical Practitioner or a Hospital or a Laboratory

candido-B cream

SPACE FOR
PHARMACODE

Presentation
CANDID-B Cream
Available in tube of 15g
Category of Distribution - POM

MANUFACTURED BY :

glenmark

PHARMACEUTICALS LTD.
PLOT NO. E-37, 39, D-ROAD,
M.I.D.C. INDUSTRIAL AREA,
SATPUR, NASHIK - 422 007,
MAHARASHTRA, (INDIA).
© TRADE MARK

PE00000 RW

Composition
Clotrimazole USP 1%
Anhydrous Beclomethasone Dipropionate BP 0.025%

Action
CANDID-B Cream offers effective therapy for dermatomycoses complicated by inflammation, pruritus, erythema or other eczematous features. Clotrimazole has broad spectrum fungicidal activity against Dermatophytes (Trichophyton, Epidermophyton and Microsporum spp.), Candida and Aspergillus spp., Malassezia furfur and other fungi. It also inhibits some strains of gram-positive and gram-negative bacteria.

Beclomethasone Dipropionate, the potent and fast acting topical corticosteroid, exerts marked anti-inflammatory, antipruritic, anti-allergic and vaso-constrictive effects. On account of these properties, it gives prompt relief in inflammation, pruritis, erythema etc.

Indications
CANDID-B Cream will be of benefit in the following fungal infections of the skin, particularly when accompanied by eczematous features :

- Tinea Capitis
- Tinea Pedis
- Tinea Comorisi
- Tinea Versicolor
- Tinea Barbae
- Erythrasma
- Tinea Cruris
- Vulvo-vaginal Candidiasis
- Tinea Manum
- Candidal Nappy Rash
- Tinea Unguium
- Candidal Paronychia

CANDID-B Cream will be particularly useful in large lesions and for hairy and intertriginous skin areas.

Application
The affected area is washed and dried. CANDID-B Cream is then rubbed in. In hairy sites the hair should be parted before application to allow direct contact with the lesions. The frequency of application is generally 2 to 3 times daily. After controlling the eczematous symptoms, a switchover to CANDID - B Cream may be advised till the primary fungal infection is completely eradicated.

times daily. After controlling the eczematous symptoms, a switchover to CANDID - B Cream may be advised till the primary fungal infection is completely eradicated.

symptoms, a switchover to CANDID - B Cream may be advised till the primary fungal infection is completely eradicated.


Tolerance
CANDID-B Cream are normally well tolerated. If irritation develops, the preparation should be discontinued.

Side Effects and Precautions
The use of steroids topically over large skin areas, with or without occlusive dressing, may result in systemic absorption of the steroid and suitable precautions should be taken by the physician. When occlusive dressings are put, miliana, folliculitis or pyoderma may sometimes develop beneath the occlusive material. Localised atrophy and striae have been reported with topical steroids.

Though topical steroids have not been reported to cause adverse effect on pregnancy, their safety in pregnancy has not been firmly established. They should, therefore, not be used in large amounts or for long periods of time during pregnancy. These formulations should not be used in the eyes or in the ear if the eardrum is perforated.


Contraindications
These formulations should not be used in tuberculosis of the skin, herpes simplex, chicken pox, measles, vaccinia,

PE00000 RW



SAME SIZE ARTWORK
LEAFLET SIZE: 140 mm x 88 mm

ICONOGRAPHICS CODE:

PANTONE SHADE
 PANTONE PROCESS C
Supersedes Artwork Code

PHARMACODE: