



LEAFLET

(Chlorpromazine Tablets BP 25 mg)

CHLOZINE-25

COMPOSITION :

Each film coated tablet contains:

Chlorpromazine Hydrochloride BP ... 25 mg

Excipients q.s.

Approved colour used

CLINICAL PHARMACOLOGY :

Chlorpromazine is a phenothiazine antipsychotic. It has a wide range of activity arising from its depressant action on the CNS and its alpha-adrenergic blocking and antimuscarinic activities. Chlorpromazine is a dopaminic inhibitor; the turnover of dopamine in the brain is also increased. The antagonism of central dopaminergic function, especially at the D₂-dopaminergic receptor, is related to therapeutic effect in psychotic conditions. It has antiemetic, serotonin-blocking and weak antimuscarinic properties and slight ganglion-blocking activity. Chlorpromazine can relax skeletal muscle.

USES:

- Acute and chronic shizophrenia in adults and children
- Acute mania as in bilopular disorder
- Autistic children
- In the alleviation of intractable hiccup

DOSAGE AND ADMINISTRATION :

1) By mouth, schizophrenia and other psychoses, mania, short-term adjunctive management of severe anxiety, psychomotor Agitation, excitement and violent or dangerously impulsive behavior initially 25 mg 3 times daily, adjusted according to response to usual maintenance dose of 75-300 mg daily (but up to 1 gm daily may be required in psychoses).



AGOG Pharma Ltd.



(WHO - GMP CERTIFIED - GOVT RECOGNISED EXPORT HOUSE)

Regd. Office & Factory : Plot No. 33, Sector II, The Vasai Taluka Industrial Co-op. Estate Ltd. Gauripada, Vasai (E), Dist. Thane - 401 208. INDIA.
Tel. : 95250 - 2455801 / 2452714 / 2453525 • Fax : 95250 - 2452074 (0091 - 250 - 2452074) • Email : agog@vsnl.net & agogpharma@rediffmail.com

- 2) Children up to 5 yrs. 0.5-1 mg/Kg body weight thrice daily.
Over 5 yrs. 1/3 of adult dose.
- 3) Intractable hiccup, 25-50 mg 3-4 times daily.

Elderly: third to half adult daily.

SPECIAL PRECAUTIONS :

It is contraindicated in patients with pre-existing CNS depression or coma, bone marrow suppression or phaeochromocytoma. It should be used with caution or not at all in patients with impaired Liver, Kidney, Cardiovascular, Cerebrovascular. It is contraindicated in Epilepsy, Parkinsonism.

ADVERSE EFFECT :

Dry mouth, constipation, difficulty with micturition, blurred vision, mydriasis, Hypersensitivity reaction various haematological disorders has been reported.

CONTRAINDICATIONS :

Liver impairment, Coma, bone marrow depression, Low leukocyte counts.

INTERACTIONS :

Additive depressant effective with sedatives, hypnotics, antihistamine, opioids and alcohols. Potentiation of anticholinergic effects of antiparkinson agents.

STORAGE : Store under normal storage conditions (15°C to 30°C).
Protect from light.

Keep all Medicines out of reach of children.

PRESENTATION : Blister pack of 10 x 10 Tablets
Jar pack of 1000 Tablets



Manufactured in India by:

AGOG PHARMA LTD.

Plot No. 33, Sector II, The Vasai Taluka Indl.
Co-op. Estate Ltd., Vasai (E), Dist. Thane. INDIA.