

150mm

150mm

220mm

Black	Height: 220mm	Color: 15-03-18
Note: For Printing match the colour with the sample.		
PSJ solutions India	Client: UNIVERSAL CORPORATION LTD.	Approved: _____
Sp: _____		

CLOTRINE
CLOTRIMAZOLE PESSARIES USP

A Product of

UNIVERSAL
CORPORATION LTD.

Club Road, Plot No. 13777
P.O. Box 1748-00902, Kikuyu, Kenya
www.ucl.co.ke

16/03/18

INDICATIONS: Vaginal candidiasis, Mixed Vaginal Infections, bacterial Vaginitis and Invasive Leishmaniasis.

APPLICATION AND DOSAGE: CLOTRINE Vaginal Pessaries should be inserted daily, preferably on empty stomach at night for 6 consecutive nights (for 100mg pessaries) or 3 consecutive nights (for 200mg pessaries) until the symptoms have completely subsided. The pessaries should be inserted as far as possible into the vagina. This is best achieved with the applicator when lying on the back with the legs pulled in close towards the body (see special directions).

The treatment should not be carried out during menstruation but should be completed before it begins. For complete relief, it is advisable to apply the pessaries during the first 2 days of menstruation. For the prevention of reinfection, the partner should be treated locally with CLOTRINE Cream. CLOTRINE Vaginal Pessaries do not stain the underwear.

TOLERANCE: There is virtually no absorption of the active ingredient through the vaginal mucosa. Therefore no systemic effects are expected. CLOTRINE Vaginal Pessaries are well tolerated locally. Administration of CLOTRINE Vaginal Pessaries during pregnancy especially in the first trimester is only under SPECIAL DIRECTIONS FOR USING THE APPLICATOR.

1. To remove the pessary first break the foil and gently push the pessary out.
2. Pull out plunger A until it stops. Place one vaginal pessary into the applicator B.
3. Insert applicator containing the pessary carefully as deep as possible into the vagina (This is done best by lying on the back).
4. Push plunger A until it stops, thereby depositing the pessary into the vagina.
5. Withdraw the applicator.
6. After use, remove plunger A completely by pulling it out of the applicator B. Thereafter, wash both in warm (not boiling) soapy water and dry thoroughly.

POSSIBLE SIDE EFFECTS: Like all medicines, CLOTRINE Vaginal Pessaries can cause side effects, although not everyone gets them. As with all medicines, some people may be allergic to the pessary. If you are allergic, a reaction will occur soon after you have used the medicine. If you experience an allergic reaction or redness, burning, pain, itching or swelling gets worse, stop using this product and tell your doctor straight away or contact the Accident and Emergency Department of your nearest hospital. Signs of an allergic reaction may include:

- Swelling of your lips, throat or tongue
- Swallowing or breathing problems
- Weakness, feeling dizzy or faint
- Nausea

Rarely, immediately after you insert the pessary you might experience:

- Mild burning or irritation

Other potential side effects have been reported:

- Itching and redness in the vagina, swelling, burning, stinging, pain in the genital area
- Irritation, itching or pain in the lower stomach, burning, stinging, pain in the genital area

If you experience any of the above effects or react badly to the pessary in any other way not listed in this leaflet, tell your doctor or pharmacist immediately.

IMPORTANT: Pregnant women should strictly abide by the instruction of their Physicians.

PRESENTATION: 100mg: Pack of 6 pessaries with applicator.
200mg: Pack of 3 pessaries with applicator.
300mg: Pack of 1 pessary with applicator.

STORAGE CONDITIONS: Do not store above 30°C. Store in a dry place. Protect from direct light. Keep out of reach of children. Last revision date: 15 March 2018.

Manufactured by:
UNIVERSAL
CORPORATION LTD.
P.O. Box 1748-00902, Kikuyu, Kenya

1036278



