

Contraindications :

1. Hypersensitivity to Carboprost Tromethamine.
2. Acute pelvic inflammatory disease.
3. Patients with active cardiac, pulmonary, renal or hepatic disease.

Precautions :

ENDOPROST should be administered cautiously in patients with a history of asthma, hypo or hypertension, cardiovascular, renal or hepatic disease, anemia, jaundice, diabetes, or epilepsy.

ENDOPROST is not recommended in patients having chorioamnionitis during labour as it may have an inhibitory effect on **ENDOPROST** activity.

ENDOPROST should be used with caution in patients with compromised or scarred uterus.

Adverse Effects :

The most frequent adverse effects observed are due to its contractile effect on smooth muscle, which are transient and reversible on cessation of therapy. These include vomiting, diarrhoea, hyperpyrexia and flushing. The incidences of these common side effects during high dose treatment of **ENDOPROST** can be minimized by pretreatment or co-administration of antiemetic and antidiarrheal drugs.

Dosage & Administration :

1. As a prophylactic treatment for the control of post partum haemorrhage :
ENDOPROST at a dose equivalent to 125mcg - 250mcg of Carboprost should be given intramuscularly at the delivery of the anterior shoulder of fetus.

2. For Refractory Post Partum Haemorrhage :
ENDOPROST at a dose equivalent to 250mcg of Carboprost is given by deep intramuscular injection at interval of about 90 minutes. The interval may be reduced if necessary but should not be less than 15 minutes. The total dose should not exceed 2000mcg (8 vials of 250 mcg).

Presentation :

ENDOPROST is available in pack sizes of 0.5ml and 1ml.

ENDOPROST - 0.5ml :

Each 0.5ml containing : Carboprost Tromethamine U.S.P. equivalent to 125mcg of Carboprost.

ENDOPROST - 1ml :

Each 1ml containing : Carboprost Tromethamine U.S.P. equivalent to 250mcg of Carboprost.

Storage :

ENDOPROST should be stored in a refrigerator (2°C - 8°C). Do not freeze.

Keep out of reach of children.

Manufactured in India by :
BHARAT SERUMS AND VACCINES LIMITED
Plot No. K-27, Additional M.I.D.C., Ambernath (E) - 421 501

IN90278E1AC

For the use only of a Registered Medical Practitioner or a Hospital or a Laboratory.



Carboprost Tromethamine Injection U.S.P.
ENDOPROST
125mcg / 250mcg

For Intramuscular use only

Description :

ENDOPROST is a clear colourless sterile aqueous solution of Carboprost Tromethamine for Intramuscular Injection. Carboprost is a synthetic 15 methyl analogue of naturally occurring prostaglandin F2a.

Carboprost Tromethamine is chemically a salt of (5Z,13E)-(8R,9S,11R,15S)-9,11,15-trihydroxy-15-methyl-5,13-prostadien-1-oic acid with 2-amino-2-hydroxymethyl-1,3-propanediol. The molecular formula is C₂₇H₄₆O₆N and has a molecular weight of 489.64. It is a white to slightly off-white crystalline powder and is soluble in water.

Each ml of ENDOPROST contains :

Carboprost Tromethamine U.S.P. equivalent to 125mcg & 250mcg of Carboprost as an aqueous solution containing Benzyl Alcohol U.S.P. and Sodium Chloride U.S.P. pH is adjusted with Sodium Hydroxide U.S.P. and Hydrochloric acid U.S.P.

Clinical Pharmacology :

Carboprost is a Uterine stimulant and is used for the control of post partum haemorrhage (PPH). Carboprost Tromethamine when administered as an intramuscular injection arrests intractable atonic post partum hemorrhage by inducing myometrial contractions. This haemostasis produced by Carboprost Tromethamine at the site of placental attachment minimizes the third stage blood loss and thereby reduces the maternal mortality and morbidity. Also, Carboprost tromethamine when administered prophylactically at the time of labour, because of its uterine stimulant activity it complements the physiological processes during labour, resulting in reduction of the duration of third stage and hence minimize post partum blood loss.

Indications & Usage :

ENDOPROST is indicated for the treatment of post partum haemorrhage, especially due to uterine atony which has not responded to conventional methods of management.

ENDOPROST (125mcg - 250mcg) may be used prophylactically by intramuscular injection at the birth of anterior shoulder in patients with following risks.

- Grand multipara.
- Pre-eclampsia.
- Rapid or prolonged labour.
- Placental abnormalities.
- Previous uterine surgery.
- History of PPH.
- History of retained placenta.
- Over distention of uterus.

Size : L x H = 90 x 120 mm.

Size : L x H = 90 x 120 mm.

■ Pantone Process Black C ■ 50% Pantone Process Black C ■ 25% Pantone Process Black C

Paper : 40 gsm, ITC Print Paper
Outline & Cutting marks not to print
Artwork Code No. : IN90278E1AC
Back to Back Printing
2 Horizontal Folds