



AGOG Pharma Ltd.



(WHO - GMP CERTIFIED - GOVT RECOGNISED EXPORT HOUSE)

Regd. Office & Factory : Plot No. 33, Sector II, The Vasai Taluka Industrial Co-op. Estate Ltd. Gaurai-pada, Vasai (E), Dist. Thane - 401 208, INDIA.
Tel. : 95250 - 2455801 / 2452714 / 2453525 • Fax : 95250 - 2452074 (0091 - 250 - 2452074) • Email : agog@vsnl.net & agogpharma@rediffmail.com

LEAFLET

ERO-125

DRY SYRUP

Erythromycin Estolate for Oral Suspension USP

Appearance : Dry powder after reconstitution forms pink coloured suspension.

Label Claim: Each 5 ml. of reconstituted suspension contains: Erythromycin Estolate USP Equivalent to Erythromycin 125 mg.

Clinical Pharmacology : Bacteriostatic, highly active against *Str. Pyogenes* and *C. lostricia*, *C. diphtherias*, *Listeria*, *Campylobacter*, *Legionella* and *Mycoplasma*. Moderately sensitive are *H. influenzae*, *H. dureyi*, *B. pertusis*, *Chlamydia trachomatis* and *Rickettsiae*. Erythromycin is used as an alternative to penicillins in patients allergic to the latter group and an alternative to tetracyclines in chlamydial infections and some other conditions.

Indication (s) : Upper respiratory tract infections. Acute pneumonia. Vaginitis due to *chlamydia trachomatis*. Impetigo. Wound and burn infections. Infected eczema. Acne vulgaris. Sycosis vulgaris.

Direction of use : Add cool boiled water upto the mark on the bottle and shake vigorously. Prepared suspension to be used within 7 days after the reconstitution.

Dosage and Administration: 250 mg. 6 hrly by oral, the dose may be doubled for severe infections. Gonorrhoea: 500 mg. 6 hrly for 6-7 days. Syphilis: 1 g 8 hrly. for 10-15 days (30 days in late Syphilis). Prophylaxis against streptococci: 250 mg. 12 hrly. Children: 7.5 - 25 mg/kg. body wt. hrly or 15 to 50 mg/kg. body wt. 12 hrly. The dose can be doubled if required in severe cases or as directed by the physician.

Precautions & Warnings : Hepatitis with Cholestatic jaundice, resembling viral hepatitis or extra hepatic biliary obstruction and reversible abnormalities in LFT's may be associated with prolonged or repeated therapy. Interacts with Theophylline, digoxin, Carbamazepine, oral anticoagulants.

Paediatrics: Reduced dose necessary. **Pregnancy:** Safe. **Lactation:** Contraindicated. **Elderly:** No special problem.

Contraindications : Hypersensitivity to Erythromycin.

Adverse reactions/Side effects
Mild to severe epigastric pain, Diarrhoea and other allergic reactions.

Drug interactions : Many potentiate action of carbamazepine, cyclosporin, Theophylline warfarin. Increases serum digoxin levels, reduces therapeutic effect of penicillins tarfenadine and astemizole increase the risk the adverse effect on the heart.

Storage : Store under normal storage conditions 15°C to 30°C.
Protect From sunlight.
Keep all medicines out of reach of children.

Presentation: Dry powder in 100ml amber pet bottles.



Manufactured in India by:
AGOG PHARMA LTD.
Plot No. 33, Sector - II, The Vasai Taluka Indl.
Co-op. Estate Ltd., Vasai (East), Dist. Thane, INDIA.



AGOG Pharma Ltd.



(WHO - GMP CERTIFIED - GOVT RECOGNISED EXPORT HOUSE)

Regd. Office & Factory : Plot No. 33, Sector II, The Vasai Taluka Industrial Co-op. Estate Ltd. Gaurai-pada, Vasai (E), Dist. Thane - 401 208, INDIA.
Tel. : 95250 - 2455801 / 2452714 / 2453525 • Fax : 95250 - 2452074 (0091 - 250 - 2452074) • Email : agog@vsnl.net & agogpharma@rediffmail.com

1.5.3 Patient information leaflet (PIL)

ERO-125

DRY SYRUP

Erythromycin Estolate for Oral Suspension USP

Appearance : Dry powder after reconstitution forms pink coloured suspension.

Label Claim : Each 5 ml. of reconstituted suspension contains: Erythromycin Estolate USP Equivalent to Erythromycin 125 mg.

Clinical Pharmacology : Bacteriostatic, highly active against *Str. Pyogenes* and *C. lostridia*, *C. diphtheriae*, *Listeria*, *Campylobacter*, *Legionella* and *Mycoplasma*. Moderately sensitive are *H. influenzae*, *H. ducreyi*, *B. pertussis*, *Chlamydia trachomatis* and *Rickettsiae*. Erythromycin is used as an alternative to penicillins in patients allergic to the latter group and an alternative to tetracyclines in chlamydial infections and some other conditions.

Indication (s) : Upper respiratory tract infections. Acute pneumonia. Vaginitis due to *Chlamydia trachomatis*. Impetigo. Wound and burn infections. Infected eczema. Acne vulgaris. Sycosis vulgaris.

Direction of use : Add cool boiled water upto the mark on the bottle and shake vigorously. Prepared suspension to be used within 7 days after the reconstitution.

Dosage and Administration : 250 mg. 6 hrly by oral, the dose may be doubled for severe infections. Gonorrhoea: 500 mg. 6 hrly for 6-7 days. Syphilis: 1 g 8 hrly. for 10-15 days (30 days in late Syphilis). Prophylaxis against streptococci: 250 mg. 12 hrly. Children: 7.5 - 25 mg/kg. body wt. hrly or 15 to 50 mg/kg. body wt. 12 hrly. The dose can be doubled if required in severe cases or as directed by the physician.

Precautions & Warnings : Hepatitis with Cholestatic jaundice, resembling viral hepatitis or extra hepatic biliary obstruction and reversible abnormalities in LFT's may be associated with prolonged or repeated therapy. Interacts with Theophylline, digoxin, Carbamazepine, oral anticoagulants.

Paediatrics : Reduced dose necessary. **Pregnancy :** Safe. **Lactation :** Contraindicated. **Elderly :** No special problem.

Contraindications : Hypersensitivity to Erythromycin.

Adverse reactions/Side effects
Mild to severe epigastric pain, Diarrhoea and other allergic reactions.

Drug interactions : Many potentiate action of carbamazepine, cyclosporin, Theophylline warfarin. Increases serum digoxin levels, reduces therapeutic effect of penicillins tarfenadine and astemizole increase the risk the adverse effect on the heart.

Storage : Store under normal storage conditions 15°C to 30°C.
Protect From sunlight.
Keep all medicines out of reach of children.

Presentation : Dry powder in 100ml amber pet bottles.



Manufactured in India by:
AGOG PHARMA LTD.
Plot No. 33, Sector - II, The Vasai Taluka Indl.
Co-op. Estate Ltd., Vasai (East), Dist. Thane, INDIA.