



AGOG Pharma Ltd.



(WHO - GMP CERTIFIED - GOVT RECOGNISED EXPORT HOUSE)

Regd. Office & Factory : Plot No. 33, Sector II, The Vasai Taluka Industrial Co-op. Estate Ltd. Gaurapada, Vasai (E), Dist. Thane - 401 208. INDIA.
Tel. : 95250 - 2455801 / 2452714 / 2453525 • Fax : 95250 - 2452074 (0091 - 250 - 2452074) • Email : agog@vsnl.net & agogpharma@rediffmail.com

LEAFLET

GLIBENCLAMIDE TABLETS BP

APPEARANCE :

A white coloured, oval shape, uncoated tablets having embossed 'AGOG' on one side and breakline on other side.

LABEL CLAIM :

Each uncoated tablet contains :
Glibenclamide BP 5 mg.

CLINICAL PHARMACOLOGY :

Glibenclamide dose provokes a brisk release of insulin from pancreas. It acts on beta cell membrane leading to enhanced calcium flux across it hence degranulation. After chronic admin, the insulinemic action of Glibenclamide declines, but improvement in glucose tolerance is maintained.

Thus an oral anti-diabetic preparation with an efficient hypoglycemia action.

INDICATIONS (S) :

Non-insulin dependent Diabetes Mellitus. Patients who have failed to respond to other oral antidiabetic drugs.

DOSAGES :

Initially 2.5 mg daily as single dose at breakfast increasing gradually by another 2.5 mg (only after repeating laboratory examinations).
Max 15 mg daily.

CONTRA-INDICATIONS :

Insulin-dependent Diabetes Mellitus. Diabetic coma, severe metabolic decompensation (Keep acidosis) severe renal impairment, hypersensitivity, pregnancy.



AGOG Pharma Ltd.



(WHO - GMP CERTIFIED - GOVT RECOGNISED EXPORT HOUSE)

Regd. Office & Factory : Plot No. 33, Sector II, The Vasai Taluka Industrial Co-op. Estate Ltd. Gaurai-pada, Vasai (E), Dist. Thane - 401 208. INDIA.
Tel. : 95250 - 2455801 / 2452714 / 2453525 • Fax : 95250 - 2452074 (0091 - 250 - 2452074) • Email : agog@vsnl.net & agogpharma@rediffmail.com

SPECIAL PRECAUTIONS :

In underweight, unusual physical exercise or irregular meals hypoglycemia may occur. Symptoms of hypoglycemia are headache, irritability, emotional dejection, rest lessees, insomnia, profuse sweating, impairment of performance & alertness. Glibenclamide should always be taken immediately before or along with substantial meals :

PAEDIATRICS :

Not recommended.

PREGNANCY & LACTATION :

Should not be used.

SIDE - EFFECTS :

Nausea, vomiting, anorexia or a feeling of pressure or fullness, Pruritus, urticaria, leuco-Penia, thrombocytopenia.

DRUG INTERACTION :

Warfarin, salicylate, sulphonamides and alcohol potentiate hypoglycemia effect. Glucocorticoid, diuretics and oestrogen reduce hypoglycemia effect. Synergistic hypoglycemia action with metformin.

STORAGE :

Store under normal storage conditions (15°C - 30°C).

Protect from light.

Keep all medicines out of reach of children.

PRESENTATION :

Bulk pack 1000's. Blister pack of 10 x 10, 100 x 10 Tablets.



Manufactured in India by :

AGOG PHARMA LTD.

Plot No.33, Sector II, The Vasai Taluka Indl.

Co-op.Estate Ltd., Vasai (E), Dist. Thane, INDIA.