

SAME SIZE ARTWORK  
LEAFLET SIZE : 120 mm x 170 mm  
AFTER FOLDING SIZE: 30 mm x 170 mm

30 mm

20 mm

SPACE FOR PHARMACODE

SPACE FOR PHARMACODE

For the use only of a Registered Medical Practitioner  
or a Hospital or a Laboratory

# Mumfer<sup>TM</sup> SYRUP

**Composition**  
Each 5 ml contains  
Iron (III) hydroxide polymaltose  
complex equivalent to Elemental Iron 50 mg  
In a syrup base.

**Description**  
**Mumfer** is a haematinic syrup, available in a delicious flavour.

**Indications**  
Treatment of latent iron deficiency and iron deficiency anaemia and prevention of iron deficiency before, during and after pregnancy (during lactation).  
For prophylactic therapy of iron deficiency to cover the recommended daily dietary allowances (RDA) during pregnancy & lactation.  
For prophylactic therapy of iron deficiency to cover the recommended daily dietary allowances (RDA) during childhood.

**Pharmacodynamics**  
Iron (III) hydroxide polymaltose complex is non-toxic, due to the fact that an active transport of iron or a rate determining competitive ligand exchange instead of passive diffusion process is assumed to occur. In contrast, the absorption of Ferrous salts is not controlled by a respective feedback mechanism, even in normal or iron-loaded subjects, and therefore, adverse drug reactions may occur especially when administering several doses a day.

**Dosage & Administration**  
Dosage and duration of therapy are dependent upon the extent of iron deficiency.  
The daily dose can be divided into separate doses or can be taken at one time. **Mumfer** syrup should be taken during or immediately after a meal.  
In cases of manifest iron deficiency the therapy takes 3-5 months until a normalisation of the haemoglobin value is achieved. Afterwards the therapy should be continued for several weeks with a dosage such as described for latent iron deficiency to replenish the iron stores.

PE00000 RW

SAME SIZE ARTWORK  
LEAFLET SIZE : 120 mm x 170 mm  
AFTER FOLDING SIZE: 30 mm x 170 mm

| Mumfer  | Manifestation of iron deficiency | Latent iron deficiency | Prophylactic therapy (RDA) |
|---|----------------------------------|------------------------|----------------------------|
| Infants (upto 1 year)                         | 2.5-5 ml daily                   |                        |                            |
| Children (1-12 years)                         | 5-10 ml daily                    | 2.5-5 ml daily         |                            |
| Children (>12 years adults and nursing women) | 10-20 ml daily                   | 5-10 ml daily          |                            |
| Pregnant women                                | 20-30 ml daily                   | 10 ml daily            | 5-10 ml daily              |

**Contraindications**  
Hypersensitivity to any ingredient of the formulation.

**Interactions**  
No interactions have been observed. Since, the iron is complex bound, ionic interactions with foodstuff components (phytin, oxalates, tannin, etc.) and concomitant administration of medicaments (tetracyclines, antacids) are unlikely to occur. **Mumfer** does not cause teeth staining. The haemoccult-test (selective for haemoglobin) for the detection of occult blood is not impaired and therefore iron therapy must not be interrupted.

**Effects on ability to drive and use machines**  
None.

**Overdose**  
In case of overdose neither intoxication nor iron overload have been reported.

**Incompatibilities**  
None known.

**Pharmaceutical precautions**  
Store below 30°C. Protect from light.


**Presentation**  
A bottle of 150 ml.  
Reg. No.

MANUFACTURED BY:  
**glenmark PHARMACEUTICALS LTD.**  
B/2, MAHALAXMI CHAMBERS, 22, BHULABHAI DESAI ROAD,  
MUMBAI-400 026, (INDIA).  
AT: PLOT NO. E-37, 39, D-ROAD, MIDC, SATPUR,  
NASHIK - 422 007, MAHARASHTRA STATE, INDIA.  
TM Trade Mark

PE00000 RW

ICONGRAPHICS CODE:

PANTONE SHADE



PANTONE  
BLACK  
PROCESS C

**Supersedes  
Artwork Code:  
PE52959**

**PHARMACODE:**