

## 1. NAME OF THE MEDICINAL PRODUCT

NovoRapid 100 units/ml solution for injection in vial  
NovoRapid Penfill 100 units/ml solution for injection in cartridge  
NovoRapid FlexPen 100 units/ml solution for injection in pre-filled pen  
NovoRapid InnoLet 100 units/ml solution for injection in pre-filled pen  
NovoRapid FlexTouch 100 units/ml solution for injection in pre-filled pen  
NovoRapid PumpCart 100 units/ml solution for injection in cartridge

## 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

### NovoRapid vial

1 ml solution contains 100 units insulin aspart\* (equivalent to 3.5 mg). 1 vial contains 10 ml equivalent to 1,000 units.

### NovoRapid Penfill

1 ml solution contains 100 units insulin aspart\* (equivalent to 3.5 mg). 1 cartridge contains 3 ml equivalent to 300 units.

### NovoRapid FlexPen/NovoRapid InnoLet/NovoRapid FlexTouch

1 ml solution contains 100 units insulin aspart\* (equivalent to 3.5 mg). 1 pre-filled pen contains 3 ml equivalent to 300 units.

### NovoRapid PumpCart

1 ml solution contains 100 units insulin aspart\* (equivalent to 3.5 mg). 1 cartridge contains 1.6 ml equivalent to 160 units.

\*Insulin aspart is produced in *Saccharomyces cerevisiae* by recombinant DNA technology.

For the full list of excipients, see section 6.1.

## 3. PHARMACEUTICAL FORM

Solution for injection.

The solution is clear, colourless and aqueous.

## 4. CLINICAL PARTICULARS

### 4.1 Therapeutic indications

NovoRapid is indicated for treatment of diabetes mellitus in adults, adolescents and children aged 1 year and above.

### 4.2 Posology and method of administration

#### Posology

The potency of insulin analogues, including insulin aspart, is expressed in units, whereas the potency of human insulin is expressed in international units.

NovoRapid dosing is individual and determined in accordance with the needs of the patient. It should normally be used in combination with intermediate-acting or long-acting insulin.

Moreover NovoRapid can be used for continuous subcutaneous insulin infusion (CSII) in pump systems.

It can also be administered intravenously by physicians or other healthcare staff if applicable.

Blood glucose monitoring and insulin dose adjustments are recommended to achieve optimal glycaemic control.

The individual insulin requirement in adults and children is usually between 0.5 and 1.0 unit/kg/day. In a basal-bolus treatment regimen 50-70% of this requirement may be provided by NovoRapid and the remainder by intermediate-acting or long-acting insulin.

Adjustment of dose may be necessary if patients undertake increased physical activity, change their usual diet or during concomitant illness.

### **Special populations**

#### *Elderly (≥ 65 years old)*

NovoRapid can be used in elderly patients.

In elderly patients, glucose monitoring should be intensified and the insulin aspart dose adjusted on an individual basis.

#### *Renal and hepatic impairment*

Renal or hepatic impairment may reduce the patient's insulin requirements.

In patients with renal or hepatic impairment, glucose monitoring should be intensified and the insulin aspart dose adjusted on an individual basis.

#### *Paediatric population*

NovoRapid can be used in children and adolescents aged 1 year and above in preference to soluble human insulin when a rapid onset of action might be beneficial, for example, in the timing of the injections in relation to meals (see sections 5.1 and 5.2).

The safety and efficacy of NovoRapid in children below 1 year of age have not been established. No data are available.

### **Transfer from other insulin medicinal products**

When transferring from other insulin medicinal products, adjustment of the NovoRapid dose and the dose of the basal insulin may be necessary. NovoRapid has a faster onset and a shorter duration of action than soluble human insulin. When injected subcutaneously into the abdominal wall, the onset of action will occur within 10-20 minutes of injection. The maximum effect is exerted between 1 and 3 hours after the injection. The duration of action is 3 to 5 hours.

Close glucose monitoring is recommended during the transfer and in the initial weeks thereafter (see section 4.4).

### **Method of administration**

NovoRapid is a rapid-acting insulin analogue.

NovoRapid is administered subcutaneously by injection in the abdominal wall, the thigh, the upper arm, the deltoid region or the gluteal region. Injection sites should always be rotated within the same region in order to reduce the risk of lipodystrophy. Subcutaneous injection in the abdominal wall ensures a faster absorption than other injection sites. Compared to soluble human insulin the faster onset of action of NovoRapid is maintained regardless of the injection site. The duration of action will vary according to the dose, injection site, blood flow, temperature and level of physical activity.

Due to the faster onset of action, NovoRapid should generally be given immediately before a meal. When necessary NovoRapid can be given soon after a meal.

#### *Continuous Subcutaneous Insulin Infusion (CSII)*

NovoRapid may be used for CSII in pump systems suitable for insulin infusion. CSII should be administered in the abdominal wall. Infusion sites should be rotated.

When used with an insulin infusion pump, NovoRapid should not be mixed with any other insulin medicinal products.

Patients using CSII should be comprehensively instructed in the use of the pump system and use the correct reservoir and tubing for the pump (see section 6.6). The infusion set (tubing and cannula) should be changed in accordance with the instructions in the product information supplied with the infusion set.

Patients administering NovoRapid by CSII must have an alternative insulin delivery method available in case of pump system failure.

#### *Intravenous use*

If necessary, NovoRapid can be administered intravenously which should be carried out by physicians or other healthcare staff.

For intravenous use, infusion systems with NovoRapid 100 units/ml at concentrations from 0.05 unit/ml to 1.0 unit/ml insulin aspart in the infusion fluids 0.9% sodium chloride, 5% dextrose or 10% dextrose including 40 mmol/l potassium chloride using polypropylene infusion bags, are stable at room temperature for 24 hours.

Although stable over time, a certain amount of insulin will be initially adsorbed to the material of the infusion bag. Monitoring of blood glucose is necessary during insulin infusion.

For detailed user instructions, please refer to the package leaflet.

#### NovoRapid vial

##### *Administration with a syringe*

NovoRapid vials are for use with insulin syringes with the corresponding unit scale. See also section 6.2.

#### NovoRapid Penfill

##### *Administration with an insulin delivery system*

NovoRapid Penfill is designed to be used with Novo Nordisk insulin delivery systems and NovoFine or NovoTwist needles.

#### NovoRapid FlexPen

##### *Administration with FlexPen*

NovoRapid FlexPen is a pre-filled pen (colour-coded) designed to be used with NovoFine or NovoTwist disposable needles up to a length of 8 mm. FlexPen delivers 1-60 units in increments of 1 unit.

#### NovoRapid InnoLet

##### *Administration with InnoLet*

NovoRapid InnoLet is a pre-filled pen designed to be used with NovoFine or NovoTwist disposable needles up to a length of 8 mm. InnoLet delivers 1-50 units in increments of 1 unit.

#### NovoRapid FlexTouch

##### *Administration with FlexTouch*

NovoRapid FlexTouch is a pre-filled pen (colour-coded) designed to be used with NovoFine or NovoTwist disposable needles up to a length of 8 mm. FlexTouch delivers 1-80 units in increments of 1 unit.

#### NovoRapid PumpCart

##### *Administration via Continuous Subcutaneous Insulin Infusion (CSII)*

NovoRapid PumpCart is only for use with an insulin infusion pump system designed to be used with this cartridge, such as the Accu-Chek Insight and YpsoPump insulin pumps.

CSII should be administered in the abdominal wall. Infusion sites should be rotated.

### **4.3 Contraindications**

Hypersensitivity to the active substance or to any of the excipients (see section 6.1).

### **4.4 Special warnings and precautions for use**

Before travelling between different time zones, the patient should seek the doctor's advice since this may mean that the patient has to take the insulin and meals at different times.

#### NovoRapid PumpCart

##### *Misuse of NovoRapid PumpCart*

NovoRapid PumpCart is only for use with an insulin infusion pump system designed to be used with this cartridge, such as the Accu-Chek Insight and YpsoPump insulin pumps. It must not be used with other devices not designed for NovoRapid PumpCart, as this may result in incorrect insulin dosing and subsequent hyper- or hypoglycaemia.

### **Hyperglycaemia**

Inadequate dosing or discontinuation of treatment, especially in type 1 diabetes, may lead to hyperglycaemia and diabetic ketoacidosis. Usually the first symptoms of hyperglycaemia develop gradually over a period of hours or days. They include thirst, increased frequency of urination, nausea, vomiting, drowsiness, flushed dry skin, dry mouth, loss of appetite as well as acetone odour of breath. In type 1 diabetes, untreated hyperglycaemic events eventually lead to diabetic ketoacidosis, which is potentially lethal.

### **Hypoglycaemia**

Omission of a meal or unplanned, strenuous physical exercise may lead to hypoglycaemia. Especially in children, care should be taken to match insulin doses (especially in basal-bolus regimens) with food intake, physical activities and current blood glucose level in order to minimise the risk of hypoglycaemia.

Hypoglycaemia may occur if the insulin dose is too high in relation to the insulin requirement. In case of hypoglycaemia or if hypoglycaemia is suspected NovoRapid must not be injected. After stabilisation of patient's blood glucose adjustment of the dose should be considered (see sections 4.8 and 4.9).

Patients whose blood glucose control is greatly improved, e.g. by intensified insulin therapy, may experience a change in their usual warning symptoms of hypoglycaemia, and should be advised accordingly. Usual warning symptoms may disappear in patients with longstanding diabetes.

A consequence of the pharmacodynamics of rapid-acting insulin analogues is that if hypoglycaemia occurs, it may occur earlier after an injection when compared with soluble human insulin.

Since NovoRapid should be administered in immediate relation to a meal, the rapid onset of action should be considered in patients with concomitant diseases or treatment where a delayed absorption of food might be expected.

Concomitant illness, especially infections and feverish conditions, usually increases the patient's insulin requirements. Concomitant diseases in the kidney, liver or affecting the adrenal, pituitary or thyroid gland can require changes in the insulin dose.

When patients are transferred between different types of insulin medicinal products, the early warning symptoms of hypoglycaemia may change or become less pronounced than those experienced with their previous insulin.

### **Transfer from other insulin medicinal products**

Transferring a patient to another type or brand of insulin should be done under strict medical supervision. Changes in strength, brand (manufacturer), type, origin (animal, human insulin or human insulin analogue) and/or method of manufacture (recombinant DNA versus animal source insulin) may result in the need for a change in dose. Patients transferred to NovoRapid from another type of insulin may require an increased number of daily injections or a change in dose from that used with their usual insulin medicinal products. If an adjustment is needed, it may occur with the first dose or during the first few weeks or months.

### **Injection site reactions**

As with any insulin therapy, injection site reactions may occur and include pain, redness, hives, inflammation, bruising, swelling and itching. Continuous rotation of the injection site within a given area reduces the risk of developing these reactions. Reactions usually resolve in a few days to a few weeks. On rare occasions, injection site reactions may require discontinuation of NovoRapid.

### **Combination of NovoRapid with pioglitazone**

Cases of cardiac failure have been reported when pioglitazone was used in combination with insulin, especially in patients with risk factors for development of cardiac heart failure. This should be kept in mind if treatment with the combination of pioglitazone and NovoRapid is considered. If the combination is used, patients should be observed for signs and symptoms of heart failure, weight gain and oedema. Pioglitazone should be discontinued if any deterioration in cardiac symptoms occurs.

### **Avoidance of accidental mix-ups/medication errors**

Patients must be instructed to always check the insulin label before each injection to avoid accidental mix-ups between NovoRapid and other insulin products.

### **Insulin antibodies**

Insulin administration may cause insulin antibodies to form. In rare cases, the presence of such insulin antibodies may necessitate adjustment of the insulin dose in order to correct a tendency to hyper- or hypoglycaemia.

## **4.5 Interaction with other medicinal products and other forms of interaction**

A number of medicinal products are known to interact with the glucose metabolism.

The following substances may reduce the patient's insulin requirements:

Oral antidiabetic medicinal products, monoamine oxidase inhibitors (MAOI), beta-blockers, angiotensin converting enzyme (ACE) inhibitors, salicylates, anabolic steroids and sulphonamides.

The following substances may increase the patient's insulin requirements:

Oral contraceptives, thiazides, glucocorticoids, thyroid hormones, sympathomimetics, growth hormone and danazol.

Beta-blockers may mask the symptoms of hypoglycaemia.

Octreotide/lanreotide may either increase or decrease the insulin requirement.

Alcohol may intensify or reduce the hypoglycaemic effect of insulin.

#### **4.6 Fertility, pregnancy and lactation**

##### **Pregnancy**

NovoRapid (insulin aspart) can be used in pregnancy. Data from two randomised controlled clinical trials (322 and 27 exposed pregnancies) do not indicate any adverse effect of insulin aspart on pregnancy or on the health of the foetus/newborn when compared to human insulin (see section 5.1).

Intensified blood glucose control and monitoring of pregnant women with diabetes (type 1 diabetes, type 2 diabetes or gestational diabetes) are recommended throughout pregnancy and when contemplating pregnancy. Insulin requirements usually fall in the first trimester and increase subsequently during the second and third trimester. After delivery, insulin requirements normally return rapidly to pre-pregnancy values.

##### **Breast-feeding**

There are no restrictions on treatment with NovoRapid during breast-feeding. Insulin treatment of the nursing mother presents no risk to the baby. However, the NovoRapid dose may need to be adjusted.

##### **Fertility**

Animal reproduction studies have not revealed any differences between insulin aspart and human insulin regarding fertility.

#### **4.7 Effects on ability to drive and use machines**

The patient's ability to concentrate and react may be impaired as a result of hypoglycaemia. This may constitute a risk in situations where these abilities are of special importance (e.g. driving a car or operating machinery).

Patients should be advised to take precautions to avoid hypoglycaemia while driving. This is particularly important in those who have reduced or absent awareness of the warning signs of hypoglycaemia or have frequent episodes of hypoglycaemia. The advisability of driving should be considered in these circumstances.

#### **4.8 Undesirable effects**

##### **Summary of the safety profile**

Adverse reactions observed in patients using NovoRapid are mainly due to the pharmacologic effect of insulin.

The most frequently reported adverse reaction during treatment is hypoglycaemia. The frequencies of hypoglycaemia vary with patient population, dose regimens and level of glycaemic control (see section 4.8, Description of selected adverse reactions).

At the beginning of the insulin treatment, refraction anomalies, oedema and injection site reactions (pain, redness, hives, inflammation, bruising, swelling and itching at the injection site) may occur. These reactions are usually of transitory nature. Fast improvement in blood glucose control may be associated with acute painful neuropathy, which is usually reversible. Intensification of insulin therapy with abrupt improvement in glycaemic control may be associated with temporary worsening of

diabetic retinopathy, while long-term improved glycaemic control decreases the risk of progression of diabetic retinopathy.

### Tabulated list of adverse reactions

Adverse reactions listed below are based on clinical trial data and classified according to MedDRA frequency and System Organ Class. Frequency categories are defined according to the following convention: Very common ( $\geq 1/10$ ); common ( $\geq 1/100$  to  $< 1/10$ ); uncommon ( $\geq 1/1,000$  to  $< 1/100$ ); rare ( $\geq 1/10,000$  to  $< 1/1,000$ ); very rare ( $< 1/10,000$ ); not known (cannot be estimated from the available data).

Immune system disorders	Uncommon – Urticaria, rash, eruptions
	Very rare – Anaphylactic reactions*
Metabolism and nutrition disorders	Very common – Hypoglycaemia*
Nervous system disorders	Rare – Peripheral neuropathy (painful neuropathy)
Eye disorders	Uncommon – Refraction disorders
	Uncommon – Diabetic retinopathy
Skin and subcutaneous tissue disorders	Uncommon – Lipodystrophy*
General disorders and administration site conditions	Uncommon – Injection site reactions
	Uncommon – Oedema

\* see section 4.8, Description of selected adverse reactions.

### Description of selected adverse reactions

#### *Anaphylactic reactions:*

The occurrence of generalised hypersensitivity reactions (including generalised skin rash, itching, sweating, gastrointestinal upset, angioneurotic oedema, difficulties in breathing, palpitation and reduction in blood pressure) is very rare but can potentially be life threatening.

#### *Hypoglycaemia:*

The most frequently reported adverse reaction is hypoglycaemia. It may occur if the insulin dose is too high in relation to the insulin requirement. Severe hypoglycaemia may lead to unconsciousness and/or convulsions and may result in temporary or permanent impairment of brain function or even death. The symptoms of hypoglycaemia usually occur suddenly. They may include cold sweats, cool pale skin, fatigue, nervousness or tremor, anxiousness, unusual tiredness or weakness, confusion, difficulty in concentration, drowsiness, excessive hunger, vision changes, headache, nausea and palpitation.

In clinical trials, the frequency of hypoglycaemia varied with patient population, dose regimens and level of glycaemic control. During clinical trials the overall rates of hypoglycaemia did not differ between patients treated with insulin aspart compared to human insulin.

#### *Lipodystrophy:*

Lipodystrophy (including lipohypertrophy, lipoatrophy) may occur at the injection site. Continuous rotation of the injection site within the particular injection area reduces the risk of developing these reactions.

### Paediatric population

Based on post-marketing sources and clinical trials, the frequency, type and severity of adverse reactions observed in the paediatric population do not indicate any differences to the broader experience in the general population.

### **Other special populations**

Based on post-marketing sources and clinical trials, the frequency, type and severity of adverse reactions observed in the elderly patients and in patients with renal or hepatic impairment do not indicate any differences to the broader experience in the general population.

### **Reporting of suspected adverse reactions**

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via [the national reporting system listed in Appendix V](#).

## **4.9 Overdose**

A specific overdose for insulin cannot be defined, however, hypoglycaemia may develop over sequential stages if too high doses relative to the patient's requirement are administered:

- Mild hypoglycaemic episodes can be treated by oral administration of glucose or sugary products. It is therefore recommended that the diabetic patient always carries sugar-containing products.
- Severe hypoglycaemic episodes, where the patient has become unconscious, can be treated with glucagon (0.5 to 1 mg) given intramuscularly or subcutaneously by a trained person, or with glucose given intravenously by physicians or other healthcare staff. Glucose must be given intravenously, if the patient does not respond to glucagon within 10 to 15 minutes. Upon regaining consciousness, administration of oral carbohydrates is recommended for the patient in order to prevent a relapse.

## **5. PHARMACOLOGICAL PROPERTIES**

### **5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: Drugs used in diabetes. Insulins and analogues for injection, fast-acting.  
ATC code: A10AB05.

#### **Mechanism of action and pharmacodynamic effects**

The blood glucose lowering effect of insulin aspart is due to the facilitated uptake of glucose following binding of insulin to receptors on muscle and fat cells and to the simultaneous inhibition of glucose output from the liver.

NovoRapid produces a more rapid onset of action compared to soluble human insulin, together with a lower glucose concentration, as assessed within the first four hours after a meal. NovoRapid has a shorter duration of action compared to soluble human insulin after subcutaneous injection.






Fig. I. Blood glucose concentrations following a single pre-meal dose of NovoRapid injected immediately before a meal (solid curve) or soluble human insulin administered 30 minutes before a meal (hatched curve) in patients with type 1 diabetes mellitus.

When NovoRapid is injected subcutaneously, the onset of action will occur within 10 to 20 minutes of injection. The maximum effect is exerted between 1 and 3 hours after injection. The duration of action is 3 to 5 hours.

### Clinical efficacy

Clinical trials in patients with type 1 diabetes have demonstrated a lower postprandial blood glucose with NovoRapid compared to soluble human insulin (Fig. I). In two long-term open label trials in patients with type 1 diabetes comprising 1070 and 884 patients, respectively, NovoRapid reduced glycated haemoglobin by 0.12 [95% C.I. 0.03; 0.22] percentage points and by 0.15 [95% C.I. 0.05; 0.26] percentage points compared to human insulin; a difference of limited clinical significance.

Clinical trials in patients with type 1 diabetes have demonstrated a reduced risk of nocturnal hypoglycaemia with insulin aspart compared with soluble human insulin. The risk of daytime hypoglycaemia was not significantly increased.

Insulin aspart is equipotent to soluble human insulin on a molar basis.

### Special populations

#### *Elderly (≥ 65 years old)*

A randomised, double-blind cross-over PK/PD trial comparing insulin aspart with soluble human insulin was performed in elderly patients with type 2 diabetes (19 patients aged 65-83 years, mean age 70 years). The relative differences in the pharmacodynamic properties ( $GIR_{max}$ ,  $AUC_{GIR, 0-120 \text{ min}}$ ) between insulin aspart and human insulin in the elderly were similar to those seen in healthy subjects and in younger patients with diabetes.

#### *Paediatric population*

A clinical trial comparing preprandial soluble human insulin with postprandial insulin aspart was performed in small children (20 patients aged 2 to less than 6 years, studied for 12 weeks, among those four were younger than 4 years old) and a single dose PK/PD trial was performed in children (6-12 years) and adolescents (13-17 years). The pharmacodynamic profile of insulin aspart in children was similar to that seen in adults.

The efficacy and safety of NovoRapid given as bolus insulin in combination with either insulin detemir or insulin degludec as basal insulin has been studied for up to 12 months, in two randomised controlled clinical trials in adolescents and children aged 1 to less than 18 years (n=712). The trials included 167 children aged 1-5 years, 260 aged 6-11 and 285 aged 12-17. The observed improvements in HbA1c and the safety profiles were comparable between all age groups.

#### *Pregnancy*

A clinical trial comparing safety and efficacy of insulin aspart vs. human insulin in the treatment of pregnant women with type 1 diabetes (322 exposed pregnancies (insulin aspart: 157; human insulin: 165)) did not indicate any adverse effect of insulin aspart on pregnancy or on the health of the foetus/newborn.

In addition the data from a clinical trial including 27 women with gestational diabetes randomised to treatment with insulin aspart vs. human insulin (insulin aspart: 14; human insulin: 13) showed similar safety profiles between treatments.

## **5.2 Pharmacokinetic properties**

### **Absorption, distribution and elimination**

In NovoRapid substitution of amino acid proline with aspartic acid at position B28 reduces the tendency to form hexamers as observed with soluble human insulin. NovoRapid is therefore more rapidly absorbed from the subcutaneous layer compared to soluble human insulin.

The time to maximum concentration is, on average, half of that for soluble human insulin. A mean maximum plasma concentration of  $492 \pm 256$  pmol/l was reached 40 (interquartile range: 30–40) minutes after a subcutaneous dose of 0.15 unit/kg bodyweight in type 1 diabetic patients. The insulin concentrations returned to baseline about 4 to 6 hours after dose. The absorption rate was somewhat slower in type 2 diabetic patients, resulting in a lower  $C_{\max}$  ( $352 \pm 240$  pmol/l) and later  $t_{\max}$  (60 (interquartile range: 50–90) minutes). The intra-individual variability in time to maximum concentration is significantly less for NovoRapid than for soluble human insulin, whereas the intra-individual variability in  $C_{\max}$  for NovoRapid is larger.

### **Special populations**

#### *Elderly ( $\geq 65$ years old)*

The relative differences in pharmacokinetic properties between insulin aspart and soluble human insulin in elderly patients (65-83 years, mean age 70 years) with type 2 diabetes were similar to those observed in healthy subjects and in younger patients with diabetes. A decreased absorption rate was observed in elderly patients, resulting in a later  $t_{\max}$  (82 (interquartile range: 60-120) minutes), whereas  $C_{\max}$  was similar to that observed in younger patients with type 2 diabetes and slightly lower than in patients with type 1 diabetes.

#### *Hepatic impairment*

A single dose pharmacokinetic study of insulin aspart was performed in 24 subjects with hepatic function ranging from normal to severely impaired. In patients with hepatic impairment, absorption rate was decreased and more variable, resulting in delayed  $t_{\max}$  from about 50 min in subjects with normal hepatic function to about 85 min in patients with moderate and severe hepatic impairment. AUC,  $C_{\max}$  and CL/F were similar in patients with reduced hepatic function compared with subjects with normal hepatic function.

#### *Renal impairment*

A single dose pharmacokinetic study of insulin aspart in 18 subjects with renal function ranging from normal to severely impaired was performed. No apparent effect of creatinine clearance values on AUC,  $C_{\max}$ , CL/F and  $t_{\max}$  of insulin aspart was found. Data were limited in patients with moderate and severe renal impairment. Patients with renal failure necessitating dialysis treatment were not investigated.

### *Paediatric population*

The pharmacokinetic and pharmacodynamic properties of NovoRapid were investigated in children (6–12 years) and adolescents (13–17 years) with type 1 diabetes. Insulin aspart was rapidly absorbed in both age groups, with similar  $t_{\max}$  as in adults. However,  $C_{\max}$  differed between the age groups, stressing the importance of the individual titration of NovoRapid.

### **5.3 Preclinical safety data**

Non-clinical data reveal no special hazard for humans based on conventional studies of safety pharmacology, repeated dose toxicity, genotoxicity and toxicity to reproduction and development.

In *in vitro* tests, including binding to insulin and IGF-1 receptor sites and effects on cell growth, insulin aspart behaved in a manner that closely resembled human insulin. Studies also demonstrate that the dissociation of binding to the insulin receptor of insulin aspart is equivalent to human insulin.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Glycerol  
Phenol  
Metacresol  
Zinc chloride  
Disodium phosphate dihydrate  
Sodium chloride  
Hydrochloric acid (for pH adjustment)  
Sodium hydroxide (for pH adjustment)  
Water for injections

### **6.2 Incompatibilities**

Substances added to NovoRapid may cause degradation of insulin aspart. This medicinal product must not be diluted or mixed with other medicinal products, except infusion fluids as described in section 4.2.

### **6.3 Shelf life**

Before opening: 30 months.

NovoRapid vial/NovoRapid Penfill/NovoRapid FlexPen/NovoRapid InnoLet/NovoRapid FlexTouch  
During use or when carried as a spare: The product must be stored for a maximum of 4 weeks. Store below 30°C.

### NovoRapid PumpCart

During use or when carried as a spare: NovoRapid PumpCart carried as a spare can be kept for up to 2 weeks below 30°C. Thereafter it can be used for up to 7 days below 37°C in an insulin infusion pump system designed to be used with this cartridge, such as the Accu-Chek Insight and YpsoPump insulin pumps.

### **6.4 Special precautions for storage**

For storage conditions of the medicinal product, see section 6.3.

Before opening: Store in a refrigerator (2°C - 8°C). Do not freeze.

#### NovoRapid vial/NovoRapid Penfill

During use or when carried as a spare: Store below 30°C. Do not refrigerate. Do not freeze. Keep the vial/cartridge in the outer carton in order to protect from light.

#### NovoRapid FlexPen/NovoRapid FlexTouch

During use or when carried as a spare: Store below 30°C. Can be stored in a refrigerator (2°C - 8°C). Do not freeze. Keep the pen cap on the pen in order to protect from light.

#### NovoRapid InnoLet

During use or when carried as a spare: Store below 30°C. Do not refrigerate. Do not freeze. Keep the pen cap on the pen in order to protect from light.

#### NovoRapid PumpCart

During use or when carried as a spare: Store below 37°C (in use) or store below 30°C (carried as a spare). Do not refrigerate. Do not freeze. Keep the cartridge in the outer carton in order to protect from light.

### **6.5 Nature and contents of container**

#### NovoRapid vial

10 ml solution in vial (type 1 glass) closed with a disc (bromobutyl/polyisoprene rubber) and a protective tamper-proof plastic cap.

Pack sizes of 1 or 5 vials of 10 ml or a multipack containing 5 packs of 1 x 10 ml vials. Not all pack sizes may be marketed.

#### NovoRapid Penfill

3 ml solution in cartridge (type 1 glass) with a plunger (bromobutyl) and a rubber closure (bromobutyl/polyisoprene).

Pack sizes of 5 and 10 cartridges. Not all pack sizes may be marketed.

#### NovoRapid FlexPen

3 ml solution in cartridge (type 1 glass) with a plunger (bromobutyl) and a rubber closure (bromobutyl/polyisoprene) contained in a pre-filled multidose disposable pen made of polypropylene.

Pack sizes of 1 (with or without needles), 5 (without needles) and 10 (without needles) pre-filled pens. Not all pack sizes may be marketed.

#### NovoRapid InnoLet

3 ml solution in cartridge (type 1 glass) with a plunger (bromobutyl) and a rubber closure (bromobutyl/polyisoprene) contained in a pre-filled multidose disposable pen made of polypropylene.

Pack sizes of 1, 5 and 10 pre-filled pens. Not all pack sizes may be marketed.

#### NovoRapid FlexTouch

3 ml solution in cartridge (type 1 glass) with a plunger (bromobutyl) and a rubber closure (bromobutyl/polyisoprene) contained in a pre-filled multidose disposable pen made of polypropylene.

Pack sizes of 1 (with or without needles), 5 (without needles) or a multipack with 2 x 5 (without needles) pre-filled pens of 3 ml. Not all pack sizes may be marketed.

#### NovoRapid PumpCart

1.6 ml solution in cartridge (type 1 glass) with a plunger (bromobutyl) and a rubber closure (bromobutyl/polyisoprene).

Pack size of 5 cartridges and a multipack containing 25 (5 packs of 5) cartridges. Not all pack sizes may be marketed.

## **6.6 Special precautions for disposal and other handling**

Do not use this medicinal product if you notice that the solution is not clear, colourless and aqueous.

NovoRapid which has been frozen must not be used.

The patient should be advised to discard the needle after each injection.

NovoRapid may be used in an infusion pump system (CSII) as described in section 4.2. Tubings in which the inner surface materials are made of polyethylene or polyolefin have been evaluated and found compatible with pump use.

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

Needles, syringes, cartridges, pre-filled pens and infusion sets must not be shared.

The cartridge must not be refilled.

In case of emergency in current NovoRapid users (hospitalisation or insulin pen malfunction), NovoRapid can be withdrawn with an U100 insulin syringe from a NovoRapid cartridge or pre-filled pen.

### NovoRapid PumpCart

NovoRapid PumpCart is a pre-filled cartridge ready for use directly in the pump. Please refer to the package leaflet where detailed instructions for use are provided.

To ensure correct dosing, NovoRapid PumpCart must not be used in an insulin pen.

NovoRapid PumpCart is only for use with an insulin infusion pump system designed to be used with this cartridge, such as the Accu-Chek Insight and YpsoPump insulin pumps, as described in section 4.2. Tubings in which the inner surface materials are made of polyethylene or polyolefin have been evaluated and found compatible with pump use.

## **7. MARKETING AUTHORISATION HOLDER**

Novo Nordisk A/S, Novo Allé, DK-2880 Bagsværd, Denmark

## **8. MARKETING AUTHORISATION NUMBERS**

### NovoRapid vial

EU/1/99/119/001

EU/1/99/119/008

EU/1/99/119/015

### NovoRapid Penfill

EU/1/99/119/003

EU/1/99/119/006

### NovoRapid FlexPen

EU/1/99/119/009

EU/1/99/119/010

EU/1/99/119/011

EU/1/99/119/017  
EU/1/99/119/018

NovoRapid InnoLet  
EU/1/99/119/012  
EU/1/99/119/013  
EU/1/99/119/014

NovoRapid FlexTouch  
EU/1/99/119/019  
EU/1/99/119/020  
EU/1/99/119/021  
EU/1/99/119/022  
EU/1/99/119/023

NovoRapid PumpCart  
EU/1/99/119/024  
EU/1/99/119/025

## **9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

Date of first authorisation: 7 September 1999  
Date of last renewal: 30 April 2009

## **10. DATE OF REVISION OF THE TEXT**

Detailed information on this medicinal product is available on the website of the European Medicines Agency <http://www.ema.europa.eu>.