

## SUMMARY OF PRODUCT CHARACTERISTICS

### 1. NAME OF THE MEDICINAL PRODUCT

**POLYGYNAX VIRGO, vaginal capsule**

### 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Neomycin sulfate.....	35 000 I.U.
Polymyxin B sulfate .....	35 000 I.U.
Nystatin .....	100 000 I.U.

For one vaginal capsule.

Excipient with known effect: hydrogenated soybean oil.

For the full list of excipients, see section 6.1.

### 3. PHARMACEUTICAL FORM

Vaginal capsule.

### 4. CLINICAL PARTICULARS

#### 4.1. THERAPEUTIC INDICATIONS

Local treatment of vaginitis due to sensitive germs and treatment of non-specific vaginitis.

The official recommendations concerning appropriate use of antibacterial products must be taken into consideration.

#### 4.2. Posology and method of administration

##### Posology

PARTICULARLY INTENDED FOR FOR FEMALE CHILDREN AND ADOLESCENTS.

One vulvo-vaginal instillation in the evening at bedtime after cleansing, for 6 consecutive days.

##### Method of administration

Instil the medicinal suspension in the vestibule and the vagina by pressing slowly on the capsule.

##### Practical advice

- The treatment should be associated with hygiene advice (wearing cotton underwear, avoiding the use of an internal tampon during treatment...) and, as far as possible, elimination of contributing factors.
- Do not discontinue the treatment during menstrual periods.

#### 4.3. Contraindications

This medicinal product is contraindicated in the following situations:

- Hypersensitivity to the active substances or to any excipients listed in section 6.1 (or relevant group sensitivity),
- In case of use of diaphragms and latex condoms,
- In case of allergy to peanut or soya, due to the presence of soybean oil.

This medicinal product is generally not recommended in combination with spermicides.

#### 4.4. Special warnings and precautions for use

##### Warnings:

In the event of local intolerance or allergic reaction, the treatment must be interrupted.

The sensitisation to antibiotics by local route may compromise the later use of the same antibiotic or related antibiotics when administered by the systemic route.

**Precautions for use:**

The duration of treatment should be limited because of the risk of selecting resistant germs and the risk of superinfection by these germs.

In the absence of data on the importance of neomycin and polymyxin B fractions resorbed by the mucosa, the risk of systemic effects, especially increased in the case of renal insufficiency, cannot be excluded.

This medicinal product contains soybean oil and may cause hypersensitivity reactions (urticaria, anaphylactic shock).

**4.5. Interaction with other medicinal products and other forms of interaction****Contraindicated combination**

**+ Condoms and Diaphragms**

Risk of rupture.

**Combination not recommended:**

**+ Spermicide**

Any local vaginal treatment is likely to inactivate a local spermicidal contraception.

**4.6. Fertility, pregnancy and lactation****Pregnancy**

Due to the presence of an aminoglycoside, neomycin, which can cause an ototoxic risk and the possibility of its systemic absorption, the use of this medicinal product is not recommended during pregnancy.

**Lactation**

Due to the digestive immaturity of the newborn and pharmacokinetic properties of this medicinal product, its prescription is not recommended during lactation.

**4.7. Effects on ability to drive and use machines**

Not relevant.

**4.8. Undesirable effects**

Possibility of contact allergic eczema, this occurs more frequently with long-term use. These lesions may spread far from the treated areas.

The risk of occurrence of toxic systemic effects (renal, auditory) is limited due to the short duration of treatment recommended.

**Reporting of suspected adverse reactions**

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the national reporting system: Agence nationale de sécurité du médicament et des produits de santé (ANSM - French Health Products Safety Agency) and Regional Pharmacovigilance Centers - Website: [www.ansm.sante.fr](http://www.ansm.sante.fr).

**4.9. Overdose**

An excessive and prolonged administration may produce systemic effects (auditory and renal) in particular in patients with renal insufficiency. A prolonged use also entails an increased risk of allergic eczema.

**5. PHARMACOLOGICAL PROPERTIES****5.1. Pharmacodynamic properties**

**Pharmacotherapeutic group: ANTIINFECTIVES AND ANTISEPTICS IN GYNECOLOGICAL USE (G. genito urinary system and sex hormones), ATC code: G01AA51.**

Combination of neomycin, polymyxin B and nystatin.  
Neomycin is an aminoglycoside antibiotic.  
Polymyxin B is a polypeptide antibiotic.  
Nystatin is an antifungal with anticandida activity.

## **ANTIBACTERIAL ACTIVITY SPECTRUM OF POLYMYXIN B AND NEOMYCIN**

### **POLYMYXIN B**

Critical concentrations separate sensitive strains from strains with intermediate sensitivity and the latter from resistant strains:

$S \leq 2 \text{ mg/l}$  and  $R > 2 \text{ mg/l}$

The prevalence of acquired resistance may vary with the geographical location and time for certain species. It is therefore useful to have information on the prevalence of local resistance, in particular when treating severe infections. These data can only be used as an orientation for the probabilities of sensitivity of a bacterial strain to this antibiotic.

When the variability of the prevalence of resistance is known in France for a bacterial species, it is provided in the following table:

Categories	Frequency of acquired resistance in France (>10%) (limit values)
<b><u>SENSITIVE SPECIES</u></b> <b>Aerobic Gram-negative bacteria</b> <i>Acinetobacter</i> <i>Aeromonas</i> <i>Alcaligenes</i> <i>Citrobacter freundii</i> <i>Citrobacter koseri</i> <i>Enterobacter</i> <i>Escherichia coli</i> <i>Klebsiella</i> <i>Moraxella</i> <i>Pseudomonas aeruginosa</i> <i>Salmonella</i> <i>Shigella</i> <i>Stenotrophomonas maltophilia</i>	0 – 30%
<b><u>RESISTANT SPECIES</u></b> <b>Aerobic Gram-positive bacteria</b> Cocci and bacilli  <b>Aerobic Gram-negative bacteria</b> <i>Branhamella catarrhalis</i> <i>Brucella</i> <i>Burkholderia cepacia</i> <i>Burkholderia pseudomallei</i> <i>Campylobacter</i> <i>Chryseobacterium meningosepticum</i> <i>Legionella</i> <i>Morganella</i> <i>Neisseria</i> <i>Proteus</i> <i>Providencia</i> <i>Serratia</i> <i>Vibrio cholerae El Tor</i>  <b>Anaerobic</b> Cocci and bacilli  <b>Others</b> Mycobacteria	

### **NEOMYCIN**

The prevalence of acquired resistance may vary with geographical location and time for certain species. It is therefore useful to have information on the prevalence of local resistance, in particular when treating severe infections. These data can only be used as an orientation for the probabilities of sensitivity of a bacterial strain to this antibiotic.

When the variability of the prevalence of resistance is known in France for a bacterial species, it is provided in the following table:

Categories	Frequency of acquired resistance in France (> 10%) (limit values)
<b><u>SENSITIVE SPECIES</u></b>	
<b>Aerobic Gram-positive bacteria</b>	
Corynebacterium	
Listeria monocytogenes	
Staphylococcus meti-S	
<b>Aerobic Gram-negative bacteria</b>	
<i>Acinetobacter</i> (essentially <i>Acinetobacter baumannii</i> )	50 – 75%
<i>Branhamella catarrhalis</i>	
<i>Campylobacter</i>	
<i>Citrobacter freundii</i>	20 – 25%
<i>Citrobacter koseri</i>	
<i>Enterobacter aerogenes</i>	?
<i>Enterobacter cloacae</i>	10 – 20%
<i>Escherichia coli</i>	15 – 25%
<i>Haemophilus influenzae</i>	25 – 35%
<i>Klebsiella</i>	10 – 15%
<i>Morganella morganii</i>	10 – 20%
<i>Proteus mirabilis</i>	20 – 50%
<i>Proteus vulgaris</i>	?
<i>Providencia rettgeri</i>	?
<i>Salmonella</i>	?
<i>Serratia</i>	?
<i>Shigella</i>	?
<i>Yersinia</i>	?

Categories	Frequency of acquired resistance in France (>10%) (limit values)
<b><u>MODERATELY SENSITIVE SPECIES</u></b> (in vitro of intermediate sensitivity) <b>Aerobic Gram-negative bacteria</b> <i>Pasteurella</i>	
<b><u>RESISTANT SPECIES</u></b> <b>Aerobic Gram-positive bacteria</b> Enterococci <i>Nocardia asteroides</i> <i>Staphylococcus meti-R*</i> <i>Streptococcus</i>  <b>Aerobic Gram-negative bacteria</b> <i>Alcaligenes denitrificans</i> <i>Burkholderia</i> <i>Flavobacterium sp.</i> <i>Providencia stuartii</i> <i>Pseudomonas aeruginosa</i> <i>Stenotrophomonas maltophilia</i>  <b>Anaerobic</b> Strictly anaerobic bacteria  <b>Others</b> <i>Chlamydia</i> Mycoplasma Rickettsias	

\*The frequency of resistance to methicillin is approximately 30 to 50% of all staphylococci and is mainly encountered in hospitals.

Remark: these spectra correspond to those of the systemic forms of these antibiotics. With local pharmaceutical presentations, concentrations obtained *in situ* are much higher than plasma concentrations. There is some remaining uncertainty concerning kinetics of *in situ* concentrations, local physicochemical conditions which can alter the antibiotic activity and product stability *in situ*.

## 5.2. Pharmacokinetic properties

Not provided.

## 5.3. Preclinical safety data

Not provided.

## 6. PHARMACEUTICAL PARTICULARS

### 6.1 List of excipients

PEG-6 stearate and glycol stearate and PEG-32 stearate (Tefose 63), hydrogenated soybean oil, dimeticone 1000.

Composition of the shell soft capsule: gelatine, glycerol, dimeticone 1000, residual purified water.

### 6.2. Incompatibilities

Not applicable.

### **6.3. Shelf-life**

2 years.

### **6.4. Special precautions for storage**

Do not store above 25°C.

### **6.5. Nature and contents of container**

6 capsules under blister (PVC/PVDC/Aluminium).

### **6.6. Special precautions for disposal and other handling**

No special requirements.

## **7. MARKETING AUTHORISATION HOLDER**

**LABORATOIRE INNOTECH INTERNATIONAL**  
22, Avenue Aristide Briand  
94 110 ARCUEIL

## **8. MARKETING AUTHORISATION NUMBER(S)**

- 34009 216 105 5 0: 6 capsules under blister PVC/PVDC/Aluminium.

## **9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

Date of first authorisation: 27<sup>th</sup> January 1997

Date of latest renewal: 27<sup>th</sup> January 2012

## **10. DATE OF REVISION OF THE TEXT**

01 December 2017

## **11. DOSIMETRY**

Not applicable.

## **12. INSTRUCTIONS FOR PREPARATION OF RADIOPHARMACEUTICALS**

Not applicable.

---

## **GENERAL CLASSIFICATION FOR SUPPLY**

List I.

Medicinal product subject to medical prescription.