

REVISED FORMULA

Trimex

ADULT • PAEDIATRIC • DIABETIC
COUGH EXPECTORANT

Presentation

The Trimex range is available as 3 different formulations, each made to give fast effective relief from chesty coughs.
Trimex Adult is an orange coloured clear syrup with a fruity odour and a pleasant (sweet) taste.
Trimex Paediatric is a reddish orange coloured viscous syrup with a fruity smell and a sweet minty taste.
Trimex Diabetic is a pale orange coloured translucent syrup with a vanilla-strawberry odour and a pleasant (mentholic) taste.

Formula:

Ingredients	Adult	Paediatric	Diabetic
Chlorpheniramine Maleate BP	2 mg	1 mg	2 mg
Pseudoephedrine Hydrochloride BP	30 mg	15 mg	30 mg
Guaifenesin BP	100 mg	50 mg	100 mg
Colour, flavour, excipients and syrup to	5 ml	5 ml	5 ml

Pharmacological Actions:

Trimex is a combination of 3 active ingredients whose pharmacological actions varies as per content.

Guaifenesin increases the volume and reduces the viscosity of tenacious sputum and is used as an expectorant for productive cough.

Chlorpheniramine Maleate is a H₁ antagonist and possesses considerable anti-histaminic and anti-cholinergic activity. It blocks the action of histamine that results in increased capillary permeability and formation of oedema and wheal which treats nasal allergies and reduces sneezing.

Pseudoephedrine Hydrochloride is both a direct and indirect acting sympathomimetic agent on adrenergic-receptors, which relieves nasal congestion.

Indications:

Trimex cough expectorant is indicated for the effective relief from chesty coughs by assisting in the loosening and removal of stubborn bronchial mucus. Its additional ingredients also assist to clear nasal congestion associated with chesty coughs.
 Trimex diabetic cough expectorant is suitable for Diabetics because it is Sugar Free.

Dosage and Administration: Number of 5ml spoonfuls to be taken three times a day

	Adult	Paediatric	Diabetic
Adults and Children Over 12 yrs	One to Two		One to Two
Children 6 - 12 yrs	Half to one	One to Two	Half to one
		Half to one	Half
Below 2 yrs	ON DOCTOR'S ADVICE		

Contra-indications:

Trimex Expectorant is contra-indicated in patients who have previously exhibited intolerance to it or any of its constituents or in patients who are taking or have taken mono-amine oxidase inhibitors within the preceding two weeks. It is also contra-indicated in patients with severe hypertension and coronary heart disease.

Precautions :

Trimex Expectorant may cause drowsiness and patients should not drive or operate machinery, if affected avoid alcohol. Trimex Expectorant should be used with caution in patients suffering from cardio-vascular disorders, hypertension, prostatic enlargement and bladder dysfunction. It should only be used in pregnancy if the potential benefits outweigh the potential unknown risks to the foetus.

Adverse Effects: :

Dryness of the mouth, throat and nose, abdominal pain with vomiting and diarrhoea, central nervous system (CNS) depressant activity, Gastro-intestinal disturbances, Hypersensitivity reaction, Blood disorders, Ventricular arrhythmias, Paradoxical CNS stimulation, sleep disturbances and skin rashes.

Storage:

Store below 30°C, in a dry and dark place. Keep out of reach of children.

Package Quantities

Trimex Expectorant Adult: 60ml, 100ml, 5L.
 Trimex Expectorant Diabetic: 100ml.
 Trimex Expectorant Paediatric: 60ml, 100ml, 5L.

Ref: P07T01/1



REGAL PHARMACEUTICALS LTD.
 Plot No. 7879/18, Off Baba Dogo Road, Ruaraka
 P.O. Box 44421 - 00100 GPO, Nairobi