



AGOG Pharma Ltd.



(WHO - GMP CERTIFIED - GOVT RECOGNISED EXPORT HOUSE)

Regd. Office & Factory : Plot No. 33, Sector II, The Vasai Taluka Industrial Co-op. Estate Ltd. Gaurai-pada, Vasai (E), Dist. Thane - 401 208. INDIA.  
Tel. : 95250 - 2455801 / 2452714 / 2453525 • Fax : 95250 - 2452074 (0091 - 250 - 2452074) • Email : agog@vsnl.net & agogpharma@rediffmail.com

### 1.5.3 Patient information leaflet (PIL)

**ZEPIN-200**  
CARBAMAZEPINE TABLETS BP 200 mg

**APPEARANCE:** White circular uncoated tablet having break line on one side & other side is "AGOG".

**COMPOSITION :** Each uncoated tablet contains :  
Carbamazepine BP ... 200 mg.

**CLINICAL PHARMACOLOGY:** It has demonstrated anticonvulsant properties in rats and mice with electrically & chemically induced seizures. It appears to act by reducing polysynaptic response and blocking the post tetanic potentiation.  
Carbamazepine has been shown to be effective in treating of psychomotor & grandmal seizures, as well as trigeminal neuralgia. It depresses thalamic potential & bulbar and polysynaptic reflexes, including linguomandibular reflex in cats.

**INDICATION:** Partial seizures with complex symptomatology (psychomotor temporal lobe); generalized tonic-clonic seizures (grand mixed seizure patterns or other partial or generalized seizure pain associated with trigeminal neuralgia, glossopharyngeal neuralgia; for prophylaxis of manic depressive psychosis in patient unresponsive to lithium therapy.

**DOSAGE & ADMINISTRATION** (direction for use)  
**Adult:** Initially 200 mg once or twice daily. Slowly to be raised to 400 mg 2-3 times daily.  
**Children:** 10-20 mg/kg body weight daily.

**SPECIAL PRECAUTIONS:** Under medical supervision. Pregnancy, lactation, severe cardiovascular diseases, hepatic or renal impairment. Monitor blood counts and liver function test.

**CONTRAINDICATION:** Hypersensitivity, atrioventricular block.

**Adverse relation/side effect:** Loss of Appetite, dryness of mouth, itching diarrhoea or constipation, headache, dizziness, somnolence, ataxia, disorder of visual accommodation.

**DRUG INTERACTION:** Failure of oral contraceptives. Toxic effect enhanced by propoxyphene. It should not be administered within 2 weeks of MAOIS therapy. Serum level decreases with phenytoin, Phenobarbital, primidone. Shorten half life of doxycycline, Neurotoxic reactions when combined with lithium.

**Storage:** Store under normal storage conditions (15°C to 30°C).  
Protect from Light.  
Keep all medicines out of reach of children.

**PRESENTATION:** Blister pack of 10 x 10 Tablets  
Jar pack of 1000 Tablets

Manufactured in India by:  
**AGOG PHARMA LTD.**  
Plot No. 33, Sector II, The Vasai Taluka Indl. Co-op. Estate Ltd., Vasai (East), Dist. Thane. INDIA.

