ZALAIN® vaginal suppository

Action pharmacologique
Sertaconazole is a new topical antimycotic endowed with a potent fungicidal activity, with a broad activity spectrum where the pathogenic yeasts (Candida spp, Candida albicans, Candida tropicalis y Pityrosporum orbiculare), dermatophytes (Trichophyton, Microsporum and Epidermophyton) and other causing and accompanying agents in skin and mucous membranes infections like gram-positives germs (Staphylococcus, and Streptococcus) are included.

Composition
Per vaginal suppository:
Nitrate de Sertaconazole……………….0.3g

The other ingredients (excipients) are:
Solid hemi-synthetic glycerides (Witepsol H19), solid hemi-synthetic glycerides (Suppocire NAI50) and colloidal anhydrous silicon.

Indications
Local treatment of infections by Candida of the vaginal mucosa

Dosage
One vaginal suppository at night when going to bed, introduced deeply into the vagina, preferably while lying down. The vaginal suppository can be used during menstruation. In general, 1 vaginal suppository suffices to eradicate the vaginal mycosis. However, if it persists, a second vaginal suppository can be applied after 7 days.

Contraindications
Known hypersensitivity to an antimycotic of the imidazoles group or to one of the excipients.

Interactions
Have not been described.

Precautions
Candidiases: it is not recommended to use a soap with acid pH (pH favouring the multiplication of Candida).

Side-effects
The absence of systemic absorption of Sertaconazole after the application of a vaginal suppository by vaginal route, makes it hardly probable that side-effects appear.

Use during pregnancy and lactation
Non embryotoxic or teratogenic effect has been observed in animal studies.

Overdosage
Non applicable

Preservation
Keep at a temperature lower than 30ºC.

Presentation
Gynecologic suppository, unit box.

Keep out of reach of children

On medical prescription

Manufactured by:
Trommsdorff GmbH & Co. KG, Germany for:

Ferrer Internacional, S.A.
Gran Vía Carlos III, 98
08028 Barcelona (Spain)