

**BIODEAL LABORATORIES LIMITED  
NAIROBI, KENYA**

**ANNEX VI (a): SUMMARY OF PRODUCT CHARACTERISTICS**

**QUERY RESPONSES ON REGISTRATION OF  
CANAZOL TABLETS- RWANDA FDA**

## **SUMMARY OF PRODUCT CHARACTERISTICS**

1. Name of the medicinal product
2. Qualitative and quantitative composition
3. Pharmaceutical form
4. Clinical particulars
  - 4.1 Therapeutic indications
  - 4.2 Posology and method of administration
  - 4.3 Contraindications
  - 4.4 Special warnings and precautions for use
  - 4.5 Interaction with other medicinal products and other forms of interaction
  - 4.6 Fertility, pregnancy and lactation
  - 4.7 Effects on ability to drive and use machines
  - 4.8 Undesirable effects
  - 4.9 Overdose
5. Pharmacological properties
  - 5.1 Pharmacodynamic properties
  - 5.2 Pharmacokinetic properties
  - 5.3 Preclinical safety data
6. Pharmaceutical particulars
  - 6.1 List of excipients
  - 6.2 Incompatibilities
  - 6.3 Shelf life
  - 6.4 Special precautions for storage
  - 6.5 Nature and contents of container
  - 6.6 Special precautions for disposal and other handling
7. Marketing authorisation holder
8. Marketing authorisation number(s)
9. Date of first authorisation/renewal of the authorisation
10. Date of revision of the text

### **1. Name of the medicinal product**

Canazol 100mg Pessary.

### **2. Qualitative and quantitative composition**

Clotrimazole 100mg.

For the full list of excipients, see section 6.1.

### **3. Pharmaceutical form**

Pessary

### **4. Clinical particulars**

#### **4.1 Therapeutic indications**

Canazol 100mg Pessaries are recommended for the treatment of candidal vaginitis.

#### **4.2 Posology and method of administration**

The pessaries should be inserted into the vagina, as high as possible, using the applicator provided. This is best achieved when lying back with legs bent up.

##### Adults:

Two pessaries should be inserted daily (preferably at night) for three consecutive days. Alternatively, one pessary may be inserted daily for six days, preferably at night. A second treatment may be carried out if necessary.

There is no separate dosage schedule for the elderly.

Canazol pessaries need moisture in the vagina in order to dissolve completely, otherwise undissolved pieces of the pessary might crumble out of the vagina. Pieces of undissolved pessary may be noticed by women who experience vaginal dryness. To help prevent this it is important that the pessary is inserted as high as possible into the vagina at bedtime.

##### Children:

Not for use in children under 16.

#### **4.3 Contraindications**

Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.

#### **4.4 Special warnings and precautions for use**

Medical advice should be sought if this is the first time the patient has experienced symptoms of candidal vaginitis.

Before using Canazol 100mg Pessaries, medical advice must be sought if any of the following are applicable:

- More than two infections of candidal vaginitis in the last six months.

- Previous history of a sexually transmitted disease or exposure to partner with sexually transmitted disease.
- Pregnancy or suspected pregnancy.
- aged under 16 or over 60 years.
- known hypersensitivity to imidazoles or other vaginal antifungal products.

Canazol 100mg Pessaries should not be used if the patient has any of the following symptoms whereupon medical advice should be sought:

- Irregular vaginal bleeding.
- Abnormal vaginal bleeding or a blood-stained discharge.
- Vulval or vaginal ulcers, blisters or sores.
- Lower abdominal pain or dysuria.
- Any adverse events such as redness, irritation or swelling associated with the treatment.
- Fever or chills.
- Nausea or vomiting.
- Diarrhoea.
- foul smelling vaginal discharge.

Treatment during the menstrual period should not be performed due to the risk of the pessary being washed out by the menstrual flow. The treatment should be finished before the onset of menstruation.

Do not use tampons, intravaginal douches, spermicides or other vaginal products while using this product.

Vaginal intercourse should be avoided in case of vaginal infection and while using this product because the partner could become infected.

When used in pregnancy, the pessary should be inserted without using an applicator (see "Pregnancy").

Patients should be advised to consult their physician if the symptoms have not been relieved within one week of using Canazol 100mg Pessaries. The pessaries can be used again if the candidal infection returns after 7 days. However, if the candidal infection recurs more than twice within six months, patients should be advised to consult their physician.

#### **4.5 Interaction with other medicinal products and other forms of interaction**

Laboratory tests have suggested that, when used together, this product may cause damage to latex contraceptives. Consequently the effectiveness of such contraceptives may be reduced. Patients should be advised to use alternative precautions for at least five days after using this product.

Concomitant treatment with vaginal clotrimazole and oral tacrolimus (FK-506; immunosuppressant) might lead to increased tacrolimus plasma levels and similarly with sirolimus. Patients should thus be

closely monitored for signs and symptoms of tacrolimus or sirolimus overdosage, if necessary by determination of the respective plasma levels.

#### **4.6 Fertility, pregnancy and lactation**

##### **Pregnancy:**

There are limited amount of data from the use of clotrimazole in pregnant women. Animal studies with clotrimazole have shown reproductive toxicity at high oral doses (see section 5.3). At the low systemic exposures of clotrimazole following vaginal treatment, harmful effects with respect to reproductive toxicity are not predicted.

Clotrimazole can be used during pregnancy, but only under the supervision of a physician or midwife.

During pregnancy the pessary should be inserted without using an applicator.

##### **Lactation:**

There are no data on the excretion of clotrimazole into human milk. However, systemic absorption is minimal after administration and is unlikely to lead to systemic effects. Clotrimazole may be used during lactation.

##### **Fertility:**

No human studies of the effects of clotrimazole on fertility have been performed, however, animal studies have not demonstrated any effects of the drug on fertility.

#### **4.7 Effects on ability to drive and use machines**

The medication has no or negligible influence on the ability to drive or use machinery.

#### **4.8 Undesirable effects**

Frequency not known. As the listed undesirable effects are based on spontaneous reports, assigning accurate frequency of occurrence for each is not possible.

Immune system disorders: anaphylactic reaction, angioedema, hypersensitivity.

Vascular disorder: syncope, hypotension.

Respiratory, thoracic and mediastinal disorders: dyspnea.

Gastrointestinal disorders: abdominal pain, nausea.

Skin and Subcutaneous Tissue Disorders: rash, urticaria, pruritus.

Reproductive system and breast disorders: vaginal exfoliation, vaginal discharge, vaginal haemorrhage, vulvovaginal discomfort, vulvovaginal erythema, vulvovaginal burning sensation, vulvovaginal pruritus, vulvovaginal pain.

General disorders and administration site conditions: application site irritation, oedema, pain.

## Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorization of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the <https://www.rwandafda.gov.rw/food-and-drugs-inspection-and-safety-monitoring>

### **4.9 Overdose**

No risk of acute intoxication is seen as it is unlikely to occur following a single vaginal or dermal application of an overdose (application over a large area under conditions favorable to absorption) or inadvertent oral ingestion. There is no specific antidote.

However, in the event of accidental oral ingestion, routine measures such as gastric lavage should be performed only if clinical symptoms of overdose become apparent (e.g. dizziness, nausea or vomiting). Gastric lavage should be carried out only if the airway can be protected adequately.

## **5. Pharmacological properties**

### **5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: Gynaecological anti-infective and antiseptics – imidazole derivatives

ATC Code: G01A F02

Mechanism of Action

Azoles (e.g. clotrimazole) are usually recommended for the local treatment of vulvovaginal candidosis that is characterized by vulvovaginal symptoms such as itching, burning, discharge, redness, swelling and soreness.

Clotrimazole acts against fungi by inhibiting ergosterol synthesis. Inhibition of ergosterol synthesis leads to structural and functional impairment of the fungal cytoplasmic membrane.

Clotrimazole has a broad antimycotic spectrum of action in vitro and in vivo, which includes dermatophytes, yeasts, moulds, etc.

Under appropriate test conditions, the MIC values for these types of fungi are in the region of less than 0.062-8.0 microgram/ml substrate. The mode of action of clotrimazole is fungistatic or fungicidal depending on the concentration of clotrimazole at the site of infection. In-vitro activity is limited to proliferating fungal elements; fungal spores are only slightly sensitive.

Primarily resistant variants of sensitive fungal species are very rare; the development of secondary resistance by sensitive fungi has so far only been observed in very isolated cases under therapeutic conditions.

### **5.2 Pharmacokinetic properties**

Pharmacokinetic investigations after vaginal application have shown that only a small amount of clotrimazole (3 – 10% of the dose) is absorbed. Due to the rapid hepatic metabolism of absorbed clotrimazole into pharmacologically inactive metabolites the resulting peak plasma concentrations of

clotrimazole after vaginal application of a 500mg dose were less than 10 ng/ml, reflecting that clotrimazole applied intra-vaginally does not lead to measurable systemic effects or side effects.

### **5.3 Preclinical safety data**

Non-clinical data reveal no special hazard for humans based on studies of repeated dose toxicity, genotoxicity and carcinogenicity.

Clotrimazole was not teratogenic in reproductive toxicity studies in mice, rats and rabbits. In rats high oral doses were associated with maternal toxicity, embryotoxicity, reduced fetal weights and decreased pup survival.

In rats clotrimazole and/or its metabolites were secreted into milk at levels higher than in plasma by a factor of 10 to 20 at 4 hrs after administration, followed by a decline to a factor of 0.4 by 24 hrs.

## **6. Pharmaceutical particulars**

### **6.1 List of excipients**

PVP-K30 BP

Sodium Lauryl Sulphate

Sodium bicarbonate

Cross Carmellose sodium

Aerosil

Lactose Monohydrate

Magnesium Stearate

Microcrystalline Cellulose

### **6.2 Incompatibilities**

Not applicable.

### **6.3 Shelf life**

36 months.

### **6.4 Special precautions for storage**

Store below 30°C. Protect from direct sunlight and moisture

### **6.5 Nature and contents of container**

Canazol (6's) Unit Boxes, Canazol literatures, Shippers, Unprinted plain foil Printed Canazol100mg foil  
Canazol Applicators

The pessaries and applicator are enclosed in a cardboard carton.

#### **6.6 Special precautions for disposal and other handling**

Any unused medicinal product or waste material should be disposed off in accordance with local requirements

#### **7. Marketing authorization holder**

Biodeal Laboratories Ltd  
Plot No.123, Lunga Lunga P.O Box 32040-00600, Nairobi, Kenya  
Telephone: +254 720 333829  
E-mail: [regulatoryaffairs@biodealkenya.com](mailto:regulatoryaffairs@biodealkenya.com)

#### **8. Marketing authorization number(s)**

H2015/CTD16/271

#### **9. Date of first authorization/renewal of the authorization**

Date of first authorization: 14 October 2015.

#### **10. Date of revision of the text**

15/09/2022