



AGOG Pharma Ltd.



(WHO - GMP CERTIFIED - GOVT RECOGNISED EXPORT HOUSE)

Regd. Office & Factory : Plot No. 33, Sector II, The Vasai Taluka Industrial Co-op. Estate Ltd. Gaurai-pada, Vasai (E), Dist. Thane - 401 208, INDIA.
Tel. : 95250 - 2455801 / 2452714 / 2453525 • Fax : 95250 - 2452074 (0091 - 250 - 2452074) • Email : agog@vsnl.net & agogpharma@rediffmail.com

LEAFLET

ERO-250 TABLETS

ERYTHROMYCIN STEARATE TABLETS BP 250 MG.

Description

Pink, circular standard convex film coated tablets with ERO-250 Printed on one side.

Pharmaceutical Form

Film Coated Tablets

Route of administration

Oral

Composition

Each film coated tablet contains :
Erythromycin Stearate BP equivalent to
Erythromycin 250 mg.

Clinical Pharmacology

Bacteriostatic, highly active against Str. Pyogenas and Str. Pneumoniae, N. gonorrhoeae, Clostridia, C. diphtheriae, Corynebacteria, Legionella and Mycoplasma. Moderately sensitive are H. influenzae, H. ducreyi, B. pertusis, Chlamydia trachomatis, Str. Viridan, N meningitidis and Rickettsiae.

Indications (s)

Upper respiratory tract infections. Acute pneumonia, Vaginitis due to chlamydia trachomatis, Impetigo, Wound and burn infections, infected eczema, Acne vulgaris, Sycosis vulgaris.

Dosage and Administration (Directions for use)

250 mg 6 hourly; the dose may be doubled 6-7 days. Syphilis 1 g 6 hourly for 10-15 days (30 days in late Syphilis). Prophylaxis against



AGOG Pharma Ltd.



(WHO - GMP CERTIFIED - GOVT RECOGNISED EXPORT HOUSE)

Regd. Office & Factory : Plot No. 33, Sector II, The Vasai Taluka Industrial Co-op. Estate Ltd. Gaurapada, Vasai (E), Dist. Thane - 401 208, INDIA.
Tel. : 95250 - 2455801 / 2452714 / 2453525 • Fax : 95250 - 2452074 (0091 - 250 - 2452074) • Email : agog@vsnl.net & agogpharma@rediffmail.com

streptococci: 250 mg 12 hourly. Children : 7.5 - 25 mg/kg body wt. hourly or 15-50 mg/kg body wt. 12 hourly. The dose can be doubled required in severe cases, or As directed by the Physician

Precaution & Warnings

Erythromycin is principally excreted by the liver. Caution should be exercised when Erythromycin is administered to patients with impaired hepatic function.

Prolonged or repeated use of Erythromycin may result in an overgrowth of nonsusceptible bacteria or fungi. If superinfection occurs, Erythromycin should be discontinued and appropriate therapy instituted.

Contraindications

Hypersensitivity to erythromycin.

Adverse reactions / Side effects

Mild to severe epigastric pain, Diarrhoea and other allergic reactions.

Storage

Store under normal storage conditions (15°C - 30°C). Protect from light.
Keep Medicine out of reach of children.

Packing & Presentation

Blister pack of 10 x 10 Tablets
Blister pack of 100 x 10 Tablets
Jar Pack of 250's, 500's, 1000's Tablets.

Manufactured in India by :



AGOG PHARMA LTD.

Plot No. 33, The Vasai Taluka Industrial Co. Op. Estate Ltd.,
Vasai (E), Dist. Thane, INDIA.