

105mm

Stopkof

Expectorant Cough Syrup

NPI Pharma

Composition:

Each 5 ml contains
Diphenhydramine Hydrochloride 13.5 mg
Ammonium Chloride 131.5 mg

Excipients:

Ammonium Chloride, Sugar, Sodium Benzoate, Sodium Saccharin, Sodium Citrate, Glycerin, Liquid Glucose, Menthol, Propylene Glycol, Citric Acid Anhydrous (For pH Adjustment), Trisodium Citrate Dihydrate (For pH Adjustment), Soluble Essence Cherry

Introduction:

Stopkof is an expectorant, containing Diphenhydramine Hydrochloride & Ammonium Chloride having antihistaminic, antitussive and anticholinergic properties.

Indications:

Stopkof is indicated for relieving of cough symptoms due to cold or allergy and relief of upper respiratory tract congestion.

Dosage:

For Adults:

5-10 ml of Stopkof syrup every 4 hours or as directed by the physician.

For Children:

6-12 years : 5 ml of Stopkof syrup 4 times a day Or as directed by the physician
Not recommended for children under 6 years.

Precautions:

Stopkof syrup should be used with caution in patients with prostatic enlargement, narrow angle glaucoma, stenosing peptic ulcer, pyloroduodenal obstruction, symptomatic prostatic hypertrophy or bladder neck obstruction and in patients with history of bronchial asthma, hyperthyroidism, cardiovascular disease or hypertension.

Concomitant use of Anti-depressants or Atropine will increase the chances of side effects such as drowsiness and or dry mouth.

Patients with Stopkof therapy should be monitored for signs of Ammonia toxicity.

Contraindications:

Stopkof syrup should not be given to patients having hypersensitivity to any of the active ingredients.
Stopkof syrup should not be given to patients receiving MAO inhibitors or CNS depressants; also it should not be

taken with alcoholic beverages and in patients with reduced liver function or metabolic alkalosis.

Use during pregnancy & lactation:

It is advisable to avoid use of stopkof syrup during pregnancy, in neonates and premature infants.

Side Effects:

Adverse effects are usually due to Diphenhydramine that enhances sedative effects, which can cause drowsiness or difficulty in concentration. Occasionally upset of stomach, constipation, nausea, dry mouth, nose or throat, difficulty-in passing urine, blurred vision may also occur.

Interaction with other drugs:

Stopkof syrup should not be given to patients receiving MAO inhibitors or CNS depressants.

Patient Instructions:

Stopkof syrup may cause sedation and drowsiness. Hence caution to be exercised while driving, operating machinery or performing tasks requiring mental alertness.

Storage: Store below 30° C , protected from light.

Presentation: 60 & 100 ml syrup in bottle.

THIS IS A MEDICAMENT

- Medicament is a product which affects your health and its consumption contrary to instruction is dangerous for you.
- Follow strictly the doctor's prescription, the method of use and the instruction of the pharmacist who sold the medicament.
- The doctor and the pharmacist are experts in medicine, its benefits and risks.
- Do not by yourself interrupt the period of treatment prescribed for you.
- Do not repeat the same prescription without consulting your doctor.

Keep medicament out of the reach of children.

Council of Arab Health Ministers
Union of Arab Pharmacists

Manufactured by :
National Pharmaceutical Industries Co. (SAOG)
Street No 15, Rusayl Industrial Area,
Muscat, Sultanate of Oman.
Rev. No: 00
Rev. Date 01/1/16
NCLA 003

150mm

Bar Code 2502