



Agog Pharma Ltd.



(WHO - GMP CERTIFIED - GOVT RECOGNISED EXPORT HOUSE)

Regd. Office & Factory : Plot No. 33, Sector II, The Vasai Taluka Industrial Co-op. Estate Ltd. Gauralpada, Vasai (E), Dist. Thane - 401 208. INDIA.
Tel. : 952550 - 2455801 / 2452714 / 2453525 • Fax : 95250 - 2452074 (0091 - 250 - 2452074) • Email : agog@vsnl.net & agogpharma@rediffmail.com

LEAFLET

AGOCARB-500

(Activated Charcoal Tablets 500 mg)

Composition:

Each uncoated tablet contains:
Activated Charcoal BP 500 mg
Excipients q.s.

The active ingredients is activated charcoal in the form of very fine powder. Agocarb-500 is used as an emergency treatment for acute oral poisoning or drug overdose. It works by adsorbing the drugs or toxic substances so reducing their absorption into the blood or helping to eliminate them from the body. Agocarb-500 itself is not absorbed into the blood and passes through the gut unchanged.

Precautions:

Tell the doctor or hospital staff if you have diabetes or any form of sugar intolerance, e.g. hereditary fructose intolerance, glucose-galactose malabsorption syndrome or sucrose-isomaltase deficiency. If Agocarb-500 is being given to your child then of course this information will apply to him/her. If you know what poison has been taken, please tell the hospital staff at once. Agocarb-500 is effective in most cases.

Drug Interaction:

Please tell your doctor or hospital staff if you are taking or have recently taken any other medicines, including medicines obtained without a prescription. Agocarb-500 will absorb most medicines and other chemical substances, and may therefore interfere with any medicines taken by mouth.

Pregnancy and breast feeding:

Please tell the doctor or hospital staff if you are pregnant, planning to become pregnant or are breast feeding. Agocarb-500 is generally regarded as safe, although the doctor may prefer a different treatment to be given under these circumstances.

Dosage:

Adults & Children over 12 years: 1-2 tablets after meals three times a day
Children under 12 years: Not recommended. Swallow with liquid.
Or as directed by the physician.

Side effects:

The most common side-effect associated with Agocarb-500 is black faces (stools). Multiple dose use of the medicine has been reported to cause obstruction of the bowel with abdominal pain. Accidental inhalation of the medicine into the lungs can cause lung damage. Other possible side effects associated with Agocarb-500 are stomach/intestinal problems including sickness, constipation and diarrhea. If any of the side effects become serious, or you notice any side effects not listed in this leaflet, please tell the doctor or hospital staff.

STORAGE:

Store under normal storage conditions (15°C to 30°C).
Protect from light. Keep all medicines out of reach of children.

PRESENTATION:

Jar Pack of 100 Tablets
Jar Pack of 1000 Tablets.
Blister pack of 10x10 Tablets



Manufactured in India by:

Agog PHARMA LTD.

Plot No. 33, Sector II, The Vasai Taluka Indl. Co-op. Estate Ltd.,
Vasai (East), Dist. Thane, INDIA.

11920101

125 MM

150 MM