

What is in this leaflet?

- What ARBITEL is and what it is used for
- What you need to know before you take ARBITEL
- How to take ARBITEL
- Possible Side Effects
- How to store ARBITEL
- Contents of the pack and other information

1. WHAT ARBITEL IS AND WHAT IT IS USED FOR

Telmisartan belongs to a class of medicines known as angiotensin II receptor antagonists. Angiotensin II is a substance produced in your body which causes your blood vessels to narrow, thus increasing your blood pressure. Telmisartan blocks the effect of angiotensin II so that the blood vessels relax, and your blood pressure is lowered. Telmisartan is used to treat essential hypertension (high blood pressure) in adults. ‘Essential’ means that the high blood pressure is not caused by any other condition.

High blood pressure, if not treated, can damage blood vessels in several organs, which could lead sometimes to heart attack, heart or kidney failure, stroke, or blindness. There are usually no symptoms of high blood pressure before damage occurs. Thus it is important to regularly measure blood pressure to verify if it is within the normal range.

Telmisartan is also used to reduce cardiovascular events (i.e. heart attack or stroke) in adults who are at risk because they have a reduced or blocked blood supply to the heart or legs, or have had a stroke or have high risk diabetes. Your doctor can tell you if you are at high risk for such events.

2. WHAT YOU NEED TO KNOW BEFORE YOU TAKE ARBITEL

Do not take Telmisartan

- if you are allergic to telmisartan or any other ingredients of this medicine (listed in section 6).
- if you are more than 3 months pregnant. (It is also better to avoid Telmisartan in early pregnancy – see pregnancy section.)
- if you have severe liver problems such as cholestasis or biliary obstruction (problems with drainage of the bile from the liver and gall bladder) or any other severe liver disease.
- if you have diabetes or impaired kidney function and you are treated with a blood pressure lowering medicine containing aliskiren.

If any of the above applies to you, tell your doctor or pharmacist before taking Telmisartan.

Other medicines and Telmisartan

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines. Your doctor may need to change the dose of these other medicines or take other precautions. In some cases you may have to stop taking one of the medicines. This applies especially to the medicines listed below taken at the same time with Telmisartan:

- Lithium containing medicines to treat some types of depression.
- Medicines that may increase blood potassium levels such as salt substitutes containing potassium, potassium-sparing diuretics (certain ‘water tablets’), ACE inhibitors, angiotensin II receptor antagonists, NSAIDs (non steroidal anti-inflammatory medicines, e.g. aspirin or ibuprofen), heparin, immunosuppressive (e.g. cyclosporin or tacrolimus), and the antibiotic trimethoprim.

- Diuretics ('water tablets'), especially if taken in high doses together with Telmisartan, may lead to excessive loss of body water and low blood pressure (hypotension).
- If you are taking an ACE-inhibitor or aliskiren (see also information under the headings "Do not take Telmisartan" and "Warnings and precautions").
- Digoxin.

The effect of Telmisartan may be reduced when you take NSAIDs (non steroidal anti-inflammatory medicines, e.g. aspirin or ibuprofen) or corticosteroids.

Telmisartan may increase the blood pressure lowering effect of other medicines used to treat high blood pressure or of medicines with blood pressure lowering potential (e.g. baclofen, amifostine).

Furthermore, low blood pressure may be aggravated by alcohol, barbiturates, narcotics or antidepressants. You may notice this as dizziness when standing up. You should consult with your doctor if you need to adjust the dose of your other medicine while taking Telmisartan.

Pregnancy and breast-feeding

Pregnancy: You must tell your doctor if you think you are (or might become) pregnant. Your doctor will normally advise you to stop taking

Telmisartan before you become pregnant or as soon as you know you are pregnant and will advise you to take another medicine instead of Telmisartan. Telmisartan is not recommended in early pregnancy, and must not be taken when more than 3 months pregnant, as it may cause serious harm to your baby if used after the third month of pregnancy.

Breast-feeding: Tell your doctor if you are breast-feeding or about to start breast-feeding. Telmisartan is not recommended for mothers who are breast-feeding, and your doctor may choose another treatment for you if you wish to breast-feed, especially if your baby is new-born, or was born prematurely.

Driving and using machines: Some people feel dizzy or tired when taking Telmisartan. If you feel dizzy or tired, do not drive or operate machinery.

3. HOW TO TAKE ARBITEL

Always take Telmisartan exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

The recommended dose of Telmisartan is one tablet a day. Try to take the tablet at the same time each day.

You can take Telmisartan with or without food. The tablets should be swallowed with some water or other non-alcoholic drink. It is important that you take Telmisartan every day until your doctor tells you otherwise. If you have the impression that the effect of

Telmisartan is too strong or too weak, talk to your doctor or pharmacist. For treatment of high blood pressure, the usual dose of Telmisartan for most patients is one 40 mg tablet once a day to control blood pressure over the 24 hour period. However, sometimes your doctor may recommend a lower dose of 20 mg or a higher dose of 80 mg. Alternatively, Telmisartan may be used in combination with diuretics ('water tablets') such as hydrochlorothiazide which has been shown to have an additive blood pressure lowering effect with Telmisartan.

For reduction of cardiovascular events, the usual dose of Telmisartan is one 80 mg tablet once a day. At the beginning of the preventive therapy with Telmisartan 80 mg, blood pressure should be frequently monitored.

If your liver is not working properly, the usual dose should not exceed 40 mg once daily.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

Warnings and precautions

Talk to your doctor if you are suffering or have ever suffered from any of the following conditions or illnesses:

- Kidney disease or kidney transplant.
- Renal artery stenosis (narrowing of the blood vessels to one or both kidneys).
- Liver disease.
- Heart trouble.
- Raised aldosterone levels (water and salt retention in the body along with imbalance of various blood minerals).

- Low blood pressure (hypotension), likely to occur if you are dehydrated (excessive loss of body water) or have salt deficiency due to diuretic therapy (‘water tablets’), low-salt diet, diarrhoea, or vomiting.
- Elevated potassium levels in your blood.
- Diabetes.

Talk to your doctor before taking Telmisartan:

- if you are taking any of the following medicines used to treat high blood pressure:
- an ACE-inhibitor (for example enalapril, lisinopril, ramipril), in particular if you have diabetes-related kidney problems.
- aliskiren.

Your doctor may check your kidney function, blood pressure, and the amount of electrolytes (e.g. potassium) in your blood at regular intervals. See also information under the heading

“Do not take Telmisartan”.

- if you are taking digoxin.

You must tell your doctor if you think you are (or might become) pregnant. Telmisartan is not recommended in early pregnancy, and must not be taken if you are more than 3 months pregnant, as it may cause serious harm to your baby if used at that stage (see pregnancy section).

In case of surgery or anaesthesia, you should tell your doctor that you are taking Telmisartan.

Telmisartan may be less effective in lowering the blood pressure in black patients.

Children and adolescents

The use of Telmisartan in children and adolescents up to the age of 18 years is not recommended.

If you take more Telmisartan than you should

If you accidentally take too many tablets, contact your doctor, pharmacist, or your nearest hospital emergency department immediately.

If you forget to take Telmisartan

If you forget to take a dose, do not worry. Take it as soon as you remember then carry on as before. If you do not take your tablet on one day, take your normal dose on the next day. ***Do not*** take a double dose to make up for forgotten individual doses.

4. POSSIBLE SIDE EFFECTS

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Some side effects can be serious and need immediate medical attention:

You should see your doctor immediately if you experience any of the following symptoms: Sepsis* (often called “blood poisoning”, is a severe infection with whole-body inflammatory response), rapid swelling of the skin and mucosa (angioedema); these side effects are rare (may affect up to 1 in 1,000 people) but are extremely serious and patients should stop taking the medicine and see their doctor immediately.

If these effects are not treated they could be fatal.

Possible side effects of Telmisartan:

Common side effects (may affect up to 1 in 10 people): Low blood pressure (hypotension) in users treated for reduction of cardiovascular events. Uncommon side effects (may affect up to 1 in 100 people):

Urinary tract infections, upper respiratory tract infections (e.g. sore throat, inflamed sinuses, common cold), deficiency in red blood cells (anaemia), high potassium levels, difficulty falling asleep, feeling sad (depression), fainting (syncope), feeling of spinning (vertigo), slow heart rate (bradycardia), low blood pressure (hypotension) in users treated for high blood pressure, dizziness on standing up (orthostatic hypotension), shortness of breath, cough, abdominal pain, diarrhoea, discomfort in the abdomen, bloating, vomiting, itching, increased sweating, drug rash, back pain, muscle cramps, muscle pain (myalgia), kidney impairment including acute kidney failure, pain in the chest, feeling of weakness, and increased level of creatinine in the blood.

Rare side effects (may affect up to 1 in 1,000 people): Sepsis* (often called “blood poisoning”, is a severe infection with whole-body inflammatory response which can lead to death), increase in certain white blood cells (eosinophilia), low platelet count (thrombocytopenia), severe allergic reaction (anaphylactic reaction), allergic

reaction (e.g. rash, itching, difficulty breathing, wheezing, swelling of the face or low blood pressure), low blood sugar levels (in diabetic patients), feeling anxious, somnolence, impaired vision, fast heart beat (tachycardia), dry mouth, upset stomach, taste disturbance (dysgeusia), abnormal liver function (Japanese patients are more likely to experience these side effect), rapid swelling of the skin and mucosa which can also lead to death (angioedema also with fatal outcome), eczema (a skin disorder), redness of skin, hives (urticaria), severe drug rash, joint pain (arthralgia), pain in extremity, tendon pain, flu-like-illness, decreased haemoglobin (a blood protein), increased levels of uric acid, increased hepatic enzymes or creatine phosphokinase in the blood.

Very rare side effects (may affect up to 1 in 10,000 people):

Progressive scarring of lung tissue (interstitial lung disease)**

* The event may have happened by chance or could be related to a mechanism currently not known.

** Cases of progressive scarring of lung tissue have been reported during intake of telmisartan. However, it is not known whether telmisartan was the cause.

5. HOW TO STORE ARBITEL

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton after “EXP”. The expiry date refers to the last day of that month.

This medicine does not require any special temperature storage conditions. You should store your medicine in the original package in order to protect the tablets from moisture. Remove your Telmisartan tablet from the blister only directly prior to intake.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. CONTENTS OF THE PACK AND OTHER INFORMATION

What Arbitel contains,

Each uncoated tablet contains

Telmisartan BP 80 mg

What Arbitel looks like and contents of the pack

White, circular ,flat, beveledged, uncoated tablets with ‘MICRO’ engraved on one side and breakline on other sides.

Manufacturer

MICRO LABS LIMITED

92, SIPCOT,

HOSUR-635 126, INDIA

Marketing Authorisation Holder

MICRO LABS LIMITED

27 Race course road

Bangalore-560001

INDIA

Date of Revision

April 2019