

MODULE -1
ADMINISTRATIVE INFORMATION FOR
BECOACTIN SYRUP

1.5.3 PATIENT INFORMATION LEAFLET

Read this entire leaflet carefully before you starts taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet?

- 1 What is Becoactin Syrup and what is it used for**
- 2 What you need to know before you take Becoactin Syrup**
- 3 How to take Becoactin Syrup**
- 4 Possible side effects**
- 5 How to store Becoactin Syrup**
- 6 Contents of the pack and other information**

1 WHAT IS BECOACTIN SYRUP AND WHAT IS IT USED FOR

Becoactin Syrup is an Appetite stimulant.

2. WHAT YOU NEED TO KNOW BEFORE YOU TAKE BECOACTIN SYRUP

Becoactin syrup is contraindicated in:

- Newborn or Premature infants
- Pregnancy & lactation
- In those patients who have shown hypersensitivity to any of the ingredients used in syrup

MODULE -1

ADMINISTRATIVE INFORMATION FOR BECOACTIN SYRUP

3. HOW TO TAKE BECOACTIN SYRUP

CHILDREN (2 to 6 years): ½ teaspoon (2.5 ml) to 1 teaspoonful (5 ml) 2 or 3 times a day

CHILDREN (7 to 14 years): 2 teaspoonfuls (10 ml) 2 or 3 times a day

ADOLESCENTS & ADULTS: 2 teaspoonfuls (10 ml) 3 times a day

OR as directed by the Physician

If you take more Becoactin Syrup than you should

It is important not to take doses. If you have taken more than you should, contact your doctor, pharmacist or nearest hospital casualty department immediately.

If you forget to take Becoactin Syrup

If you forget to take your syrup, take it as soon as you remember, unless it is nearly time for your next dose; do not take the dose you have missed. Do not take a double the dose to make up for a forgotten dose.

4. POSSIBLE SIDE EFFECTS

Confusion, hallucinations, convulsions, restlessness, blurred vision, fainting, unusual or irregular pulse, wheezing, dry mouth, dry nose, drowsiness (often transient).

5. HOW TO STORE BECOACTIN SYRUP

Protect from direct exposure to light.

Store below 30°C, at a dry place

Keep medicines out of reach of children.

6. CONTENTS OF THE PACK AND OTHER INFORMATION

What Becoactin Syrup contains

The active substance is Cyproheptadine Hydrochloride, Thiamine Hydrochloride, Riboflavin Pyridoxine Hydrochloride, Niacinamide and Cyanocobalamin.

What Becoactin Syrup looks like and contents of the pack

Yellow coloured, sweet liquid, pleasantly flavoured with pineapple packed in a 200 ml PET bottle.