

Read the patient information leaflet before use

BURNEM CREAM

(Acriflavine BPC 0.12 gm/100 gm)

Read all of this leaflet carefully before you start using this medicine.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, please ask your doctor or pharmacist.
- This medicine has been prescribed for you. Do not pass it on to others. It may harm them, even if their symptoms are the same as yours.
- If any of the side effects get serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

The name of your medicine is Acriflavine Cream, but it will be referred to as Burnem Cream throughout the remainder of this leaflet.

Index

- 1. What Burnem Cream is and what it is used for**
- 2. Before you use Burnem Cream**
- 3. How to use Burnem Cream**
- 4. Possible side effects**
- 5. How to store Burnem Cream**
- 6. Further information**

1. What Burnem Cream is and what it is used for

Burnem cream is an antiseptic (applied to living tissue/skin to reduce the possibility of infection) and bacteriostat (prevent or inhibit bacterial growth). Acriflavine is slow acting with bacteriostatic action against many gram-positive organisms and increased activity in alkaline solutions.

Indicated for minor cuts, burns, wounds, pus, bruises and topical skin infections.

2 Before you use.

Before using this drug, inform your doctor about your current list of medications, over the counter products (e.g., vitamins, herbal supplements, etc.), allergies, pre-existing diseases, and current health conditions (e.g., pregnancy, upcoming surgery, etc.). Some health conditions may make you more susceptible to the side-effects of the drug. Take as directed by your doctor or follow the direction

printed on the product insert. Dosage is based on your condition. Tell your doctor if your condition persists or worsens. Important counseling points are listed below.

- Avoid contact with mucous membrane & eyes
- Avoid use in body cavities.
- Breastfeeding
- Consult with doctor for precautions.
- Do not use it for a longer period and repeatedly.
- For topical use only
- Pregnant
- Pregnant, planning to get pregnant or breastfeeding.

Important information about some of the ingredients of Burnem Cream

Burnem Cream contains cetostearyl alcohol which can cause local skin reactions (e.g. contact dermatitis), propylene glycol may cause skin irritation and chlorocresol which may cause allergic reactions.

3. How to use.

Always use Burnem cream exactly as your doctor has told you. If you are not sure, check with your doctor or pharmacist.

For application to the skin.

Doses

- Gently apply a thin layer of cream to the affected area(s) two or three times a day.

If you use more Burnem Cream than you should

- Do not use the cream more often or for longer than your doctor tells you to.
- Do not use larger quantities than as directed, as this will increase the possibility of side effects.
- If you accidentally use too much on a few occasions do not worry.
- If you accidentally swallow some cream, small amounts are not harmful.

If you forget to use Burnem Cream

In case you forget, take it as soon as you notice. If it is close to the time of your next dose, skip the missed dose and resume your dosing schedule. Do not take extra dose to make up for a missed dose. If you are regularly missing doses, consider setting an alarm or asking a family member to remind you. Please consult your doctor to discuss changes in your dosing schedule or a new schedule to make up for missed doses if you have missed too many doses recently.

4. Possible side effects

The most reported side-effects of Burnem Cream are skin rash, skin irritation, red or itchy skin, irritation to the gastric mucosa, rashes, and cardiac arrhythmias.

The following is a list of possible side effects that may occur from the use of Burnem Cream. These side-effects are possible, but do not always occur. Some of the side-effects may be rare but serious. Consult your doctor if you observe any of the following side-effects, especially if they do not go away.

- Skin Rash
- Skin Irritation
- Red or Itchy Skin
- Irritation to The Gastric Mucosa
- Rashes
- Cardiac Arrhythmias
- Central Nervous System Depression
- Acidosis
- Cyanosis
- Pulmonary Edema
- Respiratory Failure
- Myocardial Damage
- Circulatory Failure
- Diarrhea
- Sweating
- Nausea
- Vomiting
- Metabolic Acidosis
- Local Edema
- Dizziness
- Headache

Reporting of side effects

If you get any side effects, talk to your doctor.

By reporting side affects you can help provide more information on the safety of this medicine.

5. How to store.

Keep all medicines out of the sight and reach of children.

Do not store above 30°C. Protect from light.

Once opened, do not use this medicine for more than one month.

Do not use Burnem Cream after the expiry date which is shown on the tube or carton. The expiry date refers to the last day of that month.

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines that are no longer required. This will help to protect the environment.

6. Further information

What Burnem cream contains

- The active substance (the ingredient that makes the cream work) is Acriflavine.

The other ingredients are Thymol, Cetostearyl Alcohol, Chlorocresol, Hard Paraffin, Borax, Stearic acid, Light Liquid Paraffin, Microcrystalline Wax, Sodium Acid Phosphate Dihydrate, Sodium Phosphate Dibasic, Propylene Glycol, Isopropyl alcohol, Purified water.

What Burnem Cream looks like and contents of the pack

BURNEM CREAM is available as orange yellow, smooth, homogenous, semi-solid preparation filled in 20 gm aluminium collapsible tubes.

Manufactured by:



NEM LABORATORIES PVT. LTD.,

133- Krishna Industrial Estate, Vasai (E),

Palghar – 401210, Maharashtra - INDIA

E-mail: nemlabs@gmail.com

This leaflet was last approved on “Mar-2022”