



LEAFLET

AGOBEN

Mebendazole Tablets / Oral Suspension USP

COMPOSITION :

For Mebendazole Tablets

Each Tablet contains : Mebendazole USP 100 mg

For Mebendazole Oral Suspension

Each 5 ml of Suspension contains : Mebendazole USP 100 mg

INDICATIONS :

Mebendazole is the drug of choice in : Ascariasis (roundworm infection), Trichuriasis (whipworm infection), enterobiasis (Pinworm or threadworm infection), ancylostomiasis & trichuriasis (hookworm infection). Also used in treatment of double/triple (mixed) infection (other infections where Mebendazole can be administered include those due to *Angiostrongylus cantonensis*, *gnathostoma spinigerum*, *Mansonella peritans*, *strongyloides stercoralis*, *Toxocara Spp.*)

DOSAGE :

Type of Helminth infection and Dosage schedule Enterobiasis (pinworm): 100 mg (1-tablet) once.

Ascariasis : 100 mg (1-tablet) twice daily for 3 consecutive days.

Trichuriasis (whipworm) : 100 mg (1-tablet) twice daily for 3 consecutive days.

Hookworm : 100 mg (1-tablet) twice daily for 3 consecutive days.

NOTE : IN ALL THE CASES, IF PATIENT IS NOT CURED 3 WEEKS AFTER TREATMENT, A SECOND COURSE SHOULD BE GIVEN.

Capillariasis : 400 mg per day (in 2 divided doses of 200 mg each) for atleast 20-days.

Hyalid Disease : 400-600 mg of Mebendazole 3 times a day for 21-30 days or longer.

or as directed by the physician.

ADVERSE EFFECTS :

Transient abdominal pain and diarrhoea : usually seen in-patients being treated for heavy infections. Adverse effects reported with high doses



AGOG Pharma Ltd.



(WHO - GMP CERTIFIED - GOVT RECOGNISED EXPORT HOUSE)

Regd. Office & Factory : Plot No. 33, Sector II, The Vasai Taluka Industrial Co-op. Estate Ltd. Gauripada, Vasai (E), Dist. Thane - 401 205, INDIA.
Tel. : 95250 - 2455801 / 2452714 / 2453525 • Fax : 95250 - 2452074 (0091 - 250 - 2452074) • Email : agog@vsnl.net & agogpharma@rediffmail.com

tried for hydatid disease include allergic conditions, raised liver enzyme values, alopecia and bone marrow depression.

WARNING :

Precautions : Mebendazole is teratogenic & embryotoxic in pregnant rats at doses as low as 100 mg/kg. Hence use of the same is not recommended in pregnant women. (during pregnancy, especially during the first trimester, Mebendazole should be used only if the potential benefit justifies the potential risk to the foetus). Caution should be exercised when Mebendazole is administered to nursing mothers. Not recommended for children under 2 years of age. Patients receiving high doses of Mebendazole such as those with hydatid disease, should be supervised closely with blood counts & liver function being monitored.

STORAGE :

Store below 30°C, Protect from light.

KEEP MEDICINE OUT OF THE REACH OF CHILDREN.

FOR SUSPENSION : SHAKE WELL BEFORE USE.

PRESENTATION :

A bulk pack of 100/250/500/1000 Tablets.

B blister of 6/10 Tablets.

Suspension in 30/60/100 ml in amber colour bottles.

STRICTLY TO BE USED UNDER THE ADVICE OF MEDICAL PRACTITIONER ONLY.

Manufactured in India by



AGOG PHARMA LTD.

Plot No. 33, Sector II, The Vasai Taluka Industrial Co-op. Estate Ltd.,
Vasai (E), Dist. Thane, INDIA