



AGOG Pharma Ltd.



(WHO - GMP CERTIFIED - GOVT RECOGNISED EXPORT HOUSE)

Regd. Office & Factory : Plot No. 33, Sector II, The Vasai Taluka Industrial Co-op. Estate Ltd. Gauraipada, Vasai (E), Dist. Thane - 401 208, INDIA.
Tel. : 95250 - 2455801 / 2452714 / 2453525 • Fax : 95250 - 2452074 (0091 - 250 - 2452074) • Email : agog@vsnl.net & agogpharma@rediffmail.com

LEAFLET

AGOMYCIN

SUSPENSION

Erythromycin Estolate Oral Suspension USP 125 mg / 5 ml

Appearance : Light Orange coloured suspension.

Composition:

Each 5 ml contains:

Erythromycin Estolate USP

equivalent to Erythromycin 125 mg.

Clinical Pharmacology : Bacteriostatic, highly active against *Str. Pyogenes* and *C. lostridia*, *C. diphtheriae*, *Listeria*, *Campylobacter*, *Legionella* and *Mycoplasma*. Moderately sensitive are *H. influenzae*, *H. duroyi*, *B. pertusis*, *Chlamydia trachomatis* and *Rickettsiae*. Erythromycin is used as an alternative to penicillins in patients allergic to the latter group and an alternative to tetracyclines in chlamydial infections and some other conditions.

Indication (s) : Upper respiratory tract infections. Acute pneumonia. Vaginitis due to *Chlamydia trachomatis*, Impetigo. Wound and burn infections. Infected eczema. Acne vulgaris. Sycosis vulgaris.

Dosage and Administration: 250 mg. 6 hrly by oral, the dose may be doubled for severe infections. Gonorrhoea: 500 mg. 6 hrly for 6-7 days. Syphilis: 1 g 6 hrly. for 10-15 days (30 days in late Syphilis). Prophylaxis against streptococci: 250 mg. 12 hrly. Children: 7.5 - 25 mg/kg. body wt. hrly or 15 to 50 mg/kg. body wt. 12 hrly. The dose can be doubled if required in severe cases or as directed by the physician.

Precautions & Warnings : Hepatitis with Cholestatic jaundice, resembling viral hepatitis or extra hepatic biliary obstruction and reversible abnormalities in LFT's may be associated with prolonged or repeated therapy. Interacts with Theophylline, digoxin, Carbamazepine, oral anticoagulants.

Paediatrics: Reduced dose necessary. Pregnancy: Safe. Lactation: Contraindicated. Elderly: No special problem.

Contraindications : Hypersensitivity to Erythromycin.

Adverse reactions/Side effects

Mild to severe epigastric pain, Diarrhoea and other allergic reactions.

Drug Interactions : Many potentiate action of carbamazepine, cyclosporin, Theophylline warfarin. Increases serum digoxin levels, reduces therapeutic effect of penicillins terfenadine and astemizole increase the risk the adverse effect on the heart.

Storage : Store under normal storage conditions (15°C to 30°C).

Protect from light.

Keep all medicines out of reach of children.

Presentation: 100 ml amber pet bottle.



Manufactured in India by:

AGOG PHARMA LTD.

Plot No. 33, Sector - II, The Vasai Taluka Indl.
Co-op. Estate Ltd., Vasai (East), Dist. Thane. INDIA.