

ATRACURIUM MEDIS

1. Name of the medicinal product

ATRACURIUM MEDIS 50 mg/5 ml, injectable solution
ATRACURIUM MEDIS 25 mg/2,5 ml, injectable solution

2. Qualitative and quantitative composition

Atracurium besilate HSE 10 mg/ml

For a full list of excipients, see section 6.1.

3. Pharmaceutical form

Solution for injection

4. Clinical particulars

4.1 Therapeutic indications

Atracurium Medis is a highly selective, competitive or non-depolarising neuromuscular blocking agent. It is used as an adjunct to general anesthesia or sedation in the intensive care unit (ICU), to relax skeletal muscles, and to facilitate tracheal intubation and mechanical ventilation.

4.2 Posology and method of administration

Route of administration: Intravenous injection or continuous infusion.

Used by injection in adults: Atracurium Medis is administered by intravenous injection.

The dosage range recommended for adults is 0.3 to 0.6 mg/kg (depending on the duration of full block required) and will provide adequate relaxation for about 15 to 35 minutes. Endotracheal intubation can usually be accomplished within 90 seconds from the intravenous injection of 0.5 to 0.6 mg/kg. Full block can be prolonged with supplementary doses of 0.1 to 0.2 mg/kg as required. Successive supplementary dosing does not give rise to accumulation of neuromuscular blocking effect. Spontaneous recovery from the end of full block occurs in about 35 minutes as measured by the restoration of the tetanic response to 95% of normal neuromuscular function.

The neuromuscular block produced by Atracurium Medis can be rapidly reversed by standard doses of anticholinesterase agents, such as neostigmine, accompanied or preceded by atropine, with no evidence of recurarisation.

Use as an infusion in adults: After an initial bolus dose of 0.3 to 0.6 mg/kg, Atracurium Medis can be used to maintain neuromuscular block during long surgical procedures by administration as a continuous infusion at rates of 0.3 to 0.6mg/kg/hour

Atracurium Medis can be administered by infusion during cardiopulmonary bypass surgery at the recommended infusion rates. Induced hypothermia to a body temperature of 25° to 26°C reduces the rate of inactivation of atracurium, therefore full neuromuscular block may be maintained by approximately half the original infusion rate at these low temperatures.

Atracurium Medis is compatible with the following infusion solutions for the times stated below:

<i>Infusion solution</i>	<i>Period of stability</i>
Sodium Chloride Intravenous Infusion British Pharmacopoeia (BP) (0.9% w/v)	24 hours
Glucose Intravenous Infusion BP (5% w/v)	8 hours
Ringer's Injection United States Pharmacopoeia (USP)	8 hours
Sodium Chloride (0.18%w/v) and Glucose (4% w/v) Intravenous Infusion BP	8 hours
Compound Sodium Lactate Intravenous Infusion BP (Hartmann's Solution for Injection)	4 hours

When diluted in these solutions to give atracurium besilate concentrations of 0.5 mg/ml and above, the resultant solutions will be stable in daylight for the stated periods at temperatures of up to 30°C.

Use in Children: The dosage in children over the age of one month is similar to that in adults on a bodyweight basis.

Use in Neonates: The use of Atracurium Medis is not recommended in neonates since there are insufficient data available (see section 5.1).

Use in the elderly: Atracurium Medis may be used at standard dosage in elderly patients. It is recommended, however, that the initial dose be at the lower end of the range and that it be administered slowly.

Use in patients with reduced renal and/or hepatic function: Atracurium Medis may be used at standard dosage at all levels of renal or hepatic function, including end stage failure.

Use in patients with cardiovascular disease: In patients with clinically significant cardiovascular disease, the initial dose of Atracurium Medis should be administered over a period of 60 seconds.

Use in intensive care unit (ICU) patients: After an optional initial bolus dose of Atracurium Medis of 0.3 to 0.6 mg/kg, Atracurium Medis can be used to maintain neuromuscular block by administering a continuous infusion at rates of between 11 and 13 micrograms/kg/min (0.65 to 0.78 mg/kg/hr). There may be wide inter-patient variability in dosage requirements and these may increase or decrease with time. Infusion rates as low as 4.5 microgram/kg/min (0.27 mg/kg/hr) or as high as 29.5 microgram/kg/min (1.77 mg/kg/hr) are required in some patients.

The rate of spontaneous recovery from neuromuscular block after infusion of Atracurium Medis in ICU patients is independent of the duration of administration.

Spontaneous recovery to a train-of-four ratio >0.75 (the ratio of the height of the fourth to the first twitch in a train-of-four) can be expected to occur in approximately 60 minutes. A range of 32 to 108 minutes has been observed in clinical trials.

Monitoring: In common with all neuromuscular blocking agents, monitoring of neuromuscular function is recommended during the use of Atracurium Medis in order to individualise dosage requirements.

4.3 Contraindications

Atracurium Medis is contraindicated in patients known to be hypersensitive to atracurium, cisatracurium or benzenesulfonic acid.

4.4 Special warnings and precautions for use

Precautions: In common with all the other neuromuscular blocking agents, Atracurium Medis paralyses the respiratory muscles as well as other skeletal muscles but has no effect on consciousness. Atracurium Medis should be administered only with adequate general anaesthesia and only by or under the close supervision of an experienced anaesthetist with adequate facilities for endotracheal intubation and artificial ventilation.

The potential for histamine release exists in susceptible patients during Atracurium Medis administration. Caution should be exercised in administering Atracurium Medis to patients with a history suggestive of an increased sensitivity to the effects of histamine. In particular, bronchospasm may occur in patients with a history of allergy and asthma.

High rates of cross-sensitivity between neuromuscular blocking agents have been reported. Therefore, where possible, before administering atracurium, hypersensitivity to other neuromuscular blocking agents should be excluded. Atracurium should only be used when absolutely essential in susceptible patients. Patients who experience a hypersensitivity reaction under general anaesthesia should be tested subsequently for hypersensitivity to other neuromuscular blockers.

Atracurium Medis does not have significant vagal or ganglionic blocking properties in the recommended dosage range. Consequently, Atracurium Medis has no clinically significant effects on heart rate in the recommended dosage range and it will not counteract the bradycardia produced by many anaesthetic agents or by vagal stimulation during surgery.

In common with other non-depolarising neuromuscular blocking agents, increased sensitivity to atracurium may be expected in patients with myasthenia gravis and other forms of neuromuscular disease.

As with other neuromuscular blocking agents severe acid-base and/or serum electrolyte abnormalities may increase or decrease the sensitivity of patients to Atracurium Medis.

Atracurium Medis should be administered over a period of 60 seconds to patients who may be unusually sensitive to falls in arterial blood pressure, for example those who are hypovolaemic.

Atracurium Medis is inactivated by high pH and so must not be mixed in the same syringe with thiopental or any alkaline agent.

When a small vein is selected as the injection site, Atracurium Medis should be flushed through the vein with physiological saline after injection. When other anaesthetic drugs are administered through the same in-dwelling needle or cannula as Atracurium Medis it is important that each drug is flushed through with an adequate volume of physiological saline. Atracurium Medis is hypotonic and must not be administered into the infusion line of a blood transfusion.

Studies in malignant hyperthermia in susceptible animals (swine), and clinical studies in patients susceptible to malignant hypothermia indicate that atracurium besilate does not trigger this syndrome

In common with other non-depolarising neuromuscular blocking agents, resistance may develop in patients suffering from burns. Such patients may require increased doses, dependent on the time elapsed since the burn injury and the extent of the burn.

Intensive Care Unit (ICU) patients: When administered to laboratory animals in high doses, Laudanosine, a metabolite atracurium besilate has been associated with transient hypotension and, in some species, cerebral excitatory effects. Although seizures have been seen in ICU patients receiving atracurium, a causal relationship to laudanosine has not been established (see Undesirable Effects).

4.5 Interaction with other medicinal products and other forms of interaction

The neuromuscular block produced by Atracurium Medis may be increased by the concomitant use of inhalational anaesthetics such as halothane, isoflurane, enflurane, desflurane, sévoflurane.

In common with all non-depolarising neuromuscular blocking agents the magnitude and/or duration of a non-depolarising neuromuscular block may be increased as a result of interaction with:

- antibiotics, including the aminoglycosides, polymyxins, spectinomycin, tetracyclines, lincomycin and clindamycin
- anti-arrhythmic drugs: propranolol, calcium channel blockers, lidocaine, procainamide and quinidine
- diuretics: furosemide and possibly mannitol, thiazide diuretics and acetazolamide
- magnesium sulfate
- ketamine
- lithium salts
- ganglion blocking agents, trimetaphan, hexamethonium.

Rarely certain drugs may aggravate or unmask latent myasthenia gravis or actually induce a myasthenic syndrome; increased sensitivity to Atracurium Medis would be consequent on such a development. Such drugs include various antibiotics, β - blockers (propranolol, oxprenolol), antiarrhythmic drugs (procainamide, quinidine), antirheumatic drugs (chloroquine, D- penicillamine), chlorpromazine, steroids, phenytoin and lithium.

The onset of non-depolarising neuromuscular block is likely to be lengthened and the duration of block shortened in patients receiving chronic anticonvulsant therapy.

The administration of combinations of non-depolarising neuromuscular blocking agents in conjunction with Atracurium Medis may produce a degree of neuromuscular blockage in excess of that which might be expected were an equipotent total dose of Atracurium Medis administered. Any synergistic effect may vary between different drug combinations.

A depolarising muscle relaxant such as suxamethonium chloride should not be administered to prolong the neuromuscular blocking effects of non-depolarising blocking agents such as atracurium, as this may result in a prolonged and complex block which can be difficult to reverse with anticholinesterase drugs.

Treatment with anticholinesterases, commonly used in the treatment of Alzheimer's disease e.g. donepezil, may shorten the duration and diminish the magnitude of neuromuscular blockade with atracurium

4.6 Fertility, pregnancy and lactation

Fertility

Fertility studies have not been performed

Pregnancy

Animal studies have indicated that atracurium besilate has no significant effects on foetal development.

In common with all neuromuscular blocking agents, Atracurium Medis should be used during pregnancy only if the potential benefit to the mother outweighs any potential risk to the fetus.

Atracurium Medis is suitable for maintenance of muscle relaxation during Caesarean section as it does not cross the placenta in clinically significant amounts following recommended doses.

Breast-feeding

Atracurium has a relatively high molecular weight and is highly ionized at physiological pH; these two factors considerably reduce the passage into breast milk. Also, although the acidity of milk is slightly higher than that of plasma, atracurium passed into milk is expected to be rapidly metabolized. However, it is recommended that breastfeeding should be discontinued for 24 hours after administration of atracurium.

4.7 Effects on ability to drive and use machines

This precaution is not relevant to the use of atracurium. Atracurium Medis will always be used in combination with a general anaesthetic and therefore the usual precautions relating to performance of tasks following general anaesthesia apply.

4.8 Undesirable effects

The most commonly reported adverse reactions during treatment are hypotension (mild, transient) and skin flushing, these events are attributed to histamine release. Very rarely, severe anaphylactoid or anaphylactic reactions have been reported in patients receiving atracurium in conjunction with one or more anaesthetic agents.

Adverse reactions are listed below by system organ class and frequency. Frequencies are defined as: very common > 1/10, common > 1/100 and < 1/10, uncommon > 1/1000 and < 1/100, rare > 1/10,000 and < 1/1000, very rare < 1/10,000.

Very common, common and uncommon frequencies were determined from clinical trial data. Rare and very rare frequencies were generally derived from spontaneous data. The frequency classification "Not known" has been applied to those reactions where a frequency could not be estimated from the available data.

Clinical Trial Data

<u>Vascular Disorders</u>	
Common	Hypotension (mild, transient)#, Skin flushing#
<u>Respiratory, thoracic and mediastinal disorders</u>	
Uncommon	Bronchospasm#

Post-Marketing Data

<u>Immune system disorders</u>	
Very rare	Anaphylactic reaction, anaphylactoid reaction including shock, circulatory failure and cardiac arrest
Very rarely, severe anaphylactoid or anaphylactic reactions have been reported in patients receiving atracurium in conjunction with one or more anaesthetic agents.	
<u>Nervous system disorder</u>	
Not known	Seizures
There have been reports of seizures in ICU patients who have been receiving atracurium concurrently with several other agents. These patients usually had one or more medical conditions predisposing to seizures (e.g. cranial trauma, cerebral oedema, viral encephalitis, hypoxic encephalopathy, uraemia). A causal relationship to laudanosine has not been established. In clinical trials, there appears to be no correlation between plasma laudanosine concentration and the occurrence of seizures.	
<u>Skin and subcutaneous tissue disorders</u>	
Rare	Urticaria
<u>Musculoskeletal and connective tissue disorders</u>	
Not known	Myopathy, muscle weakness

There have been some reports of muscle weakness and/or myopathy following prolonged use of muscle relaxants in severely ill patients in the ICU. Most patients were receiving concomitant corticosteroids. These events have been seen infrequently in association with atracurium and a causal relationship has not been established.

Events which have been attributed to histamine release are indicated by a hash (#)

4.9 Overdose

Symptoms: Prolonged muscle paralysis and its consequences are the main signs of overdosage.

Management: It is essential to maintain a patient airway together with assisted positive pressure ventilation until spontaneous respiration is adequate. Full sedation will be required since consciousness is not impaired. Recovery may be hastened by the administration of anticholinesterase agents accompanied by atropine, once evidence of spontaneous recovery is present.

5. Pharmacological properties

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Peripherally acting muscle relaxants: Other quaternary ammonium compounds.

ATC code: M03AC04.

Atracurium is a highly selective competitive (non-depolarising) neuromuscular blocking agent with an intermediate duration of action. Non-depolarising agents antagonise the neurotransmitter action of acetylcholine by binding with receptor sites on the motor-end-plate. Atracurium can be used in a wide range of surgical procedures and to facilitate controlled ventilation.

The onset of action of atracurium is 2 minutes on average. Its duration of action varies from 15 to 35 minutes, depending on the doses used

Atracurium provides good conditions for intubation. Under the usual conditions of anaesthesia, atracurium, by virtue of its particular and rapid elimination, does not accumulate, and several maintenance doses can therefore be administered successively. It can therefore be used during long-term surgical procedures. This elimination, independent of renal and hepatic functions, authorizes the use of atracurium in patients with renal and / or hepatic insufficiency.

The passage of atracurium through the placental barrier is insignificant, which allows its use during caesarean sections

Paediatric population:

The limited data in neonates from literature reports suggest variability in the time to onset and duration of action of atracurium in this population as compared to children (see section 4.2).

5.2 Pharmacokinetic properties

The pharmacokinetics of atracurium in man are essentially linear with the 0.3-0.6 mg/kg dose range. The elimination half-life is approximately 20 minutes, and the volume of distribution is 0.16 L/kg. Atracurium is 82% bound to plasma proteins.

Atracurium is degraded spontaneously mainly by a non-enzymatic decomposition process (Hofmann elimination) which occurs at plasma pH and at body temperature and produces breakdown products which are inactive. Degradation also occurs by ester hydrolysis catalysed by non-specific esterases. Elimination of atracurium is not dependent on kidney or liver function.

The main breakdown products are laudanosine and a monoquaternary alcohol which have no neuromuscular blocking activity. The monoquaternary alcohol is degraded spontaneously by Hofmann elimination and excreted by the kidney. Laudanosine is excreted by the kidney and metabolised by the liver. The half-life of laudanosine ranges from 3-6h in patients with normal kidney and liver function. It is about 15h in renal failure and is about 40h in renal and hepatic failure. Peak plasma levels of laudanosine are highest in patients without kidney or liver function and average 4 µg/ml with wide variation.

Concentration of metabolites are higher in ICU patients with abnormal renal and/or hepatic function (see Special Warnings and Special Precautions for Use). These metabolites do not contribute to neuromuscular block.

The clearance of atracurium is approximately 5 ml / min / kg.

Atracurium has a protein binding comparable to that of other curares which do not present a problem of drug interaction at this level. Placental passage of atracurium is variable (fetal-maternal ratios vary between 0.02 and 0.33).

5.3 Preclinical safety data

Atracurium besilate has been evaluated in three in vitro and in vivo genotoxicity tests. It has been shown to be weakly mutagenic in a single in vitro assay, in mouse lymphoma cell culture (MLA Test), at very high, cytotoxic doses, when metabolizing enzymes are added

6. Pharmaceutical particulars

6.1 List of excipients

20 percent benzene sulfonic acid solution, water for injections.

6.2 Incompatibilities

This medicine must not be mixed with other medicines except those mentioned in section 4.2.

6.3 Shelf life

24 months

6.4 Special precautions for storage

Store at temperatures between 2 and 8°C. Do not freeze. Keep container in the outer carton in order to protect from light. Any unused Atracurium Medis from opened ampoules or vials should be discarded.

6.5 Nature and contents of container

6.6 Special precautions for disposal and other handling

None

7. Marketing authorisation holder

Les Laboratoires MédiS.
Route de Tunis - KM 7 - BP 206 8000 Nabeul - Tunisie
Tel : +216 72 235 006 / Fax : +216 72 235 016
E-mail : contact@labomedis.com

8. Marketing authorisation number(s)

ATRACURIUM MEDIS 50 mg/5 ml, injectable solution: 923 340 2H
ATRACURIUM MEDIS 25 mg/2,5 ml, injectable solution: 923 340 1H

9. Date of first authorisation/renewal of the authorization

Date of first autorisation : 31/12/2005

Date of first renewal : 31/12/2010

Date of second renewal : 31/12/2015

Date of third renewal : 31/12/2020

10. Date of revision of the text

june 2021