







Regd. Office & Factory : Plot No. 33, Sector II, The Vasai Taluka Industrial Co-op. Estate Ltd. Gauraipada, Vasai (E), Dist. Thane - 401 208. INDIA l. : 95250 - 2455801 / 2452714 / 2453525 ● Fax : 95250 - 2452074 (0091 - 250 - 2452074) ● Email : agog@vsnl.net & agogpharma@rediffmail.cc

LEAFLET

CANDIZOL

(Clotrimazole Cream BP 1% w/w)

Composition:

BP 1 % w/w Clotrimazole

Cream base q.s.

It is a synthetic imidazole derivative. It is primarily used locally in the treatment of skin infections and vaginal infections due to yeasts and dermatophytes. In addition it has activity against certain gram positive bacteria and trichomonas spp.

Therapeutic class: Antifungal

Pharmacological action:

It studies of the mechanism of action, the minimum fungicidal concentration of Clotrimazole caused leakage of intracellular phosphorus compounds into the ambient medium with concomitant breakdown of cellular nucleic acids and accelerated potassium efflux. Both these events began rapidly and extensively after addition of the drug.

Indications:

Clotrimazole cream is a topically applied broad spectrum anti- fungal cream. Clotrimazole is also effective against Trichomonas, staphylococci, streptococci and bacterioides. Clotrimazole is indicated for the treatment of skin infections due to dermatophytes, yeasts, moulds and other fungi. These include ringworm (tinea) infections, interdigital mycoses(e.g.athletes foot), paronychias associated with nail mycoses, mycoses in skin folds, pityriasis versicolorerythrasma, scalp infection, Napkin rash Dhobi's itch intertrigo.

Dosage & Administration:

A thin layer of clotrimazole cream should be rubbed evenly on the affected area 2-3 timesdaily. For successful treatment Clotrimazole cream should be applied for a sufficient duration, depending on the extent and localization of the disease. In fungal infection of feet, treatment should be continued 2 weeks after the signs of the disease disappear, to prevent recurrence. The feet should be washed and dried thoroughly (especially between the toes) before applying the cream.

Contraindication:

Clotrimazole is contraindicated in individuals who have shown hypersensitivity to any of their components.



AGOG Pharma Ltd.





WHO - GMP CERTIFIED - GOVT RECOGNISED EXPORT HOUSE

Regd. Office & Factory: Plot No. 33, Sector II, The Vasai Taluka Industrial Co-op. Estate Ltd. Gauraipada, Vasai (E), Dist. Thane - 401 208. INDIA. Tal: 98750, -2455801 / 2452714 / 2453752 - Fay: 95250, -2459074 (2001 - 250.) - Email: -acor@wsnl. pet & acoronbarma@rediffmail.com

Precaution:

For external use only.

Clotrimazole cream should not be used by patients with a history of sensitivity to Clotrimazole.

Pregnant women should take the advice of a doctor before using Clotrimazole cream . Medical advice should be sought if symptoms persist. It is not for ophthalmic use.

Special precautions:

As with other creams, Canesten Cream may reduce theeffectiveness of rubber contraceptives, such as condoms ordiaphragms. Consequently, if you are using this cream onthe vulva or penis, you should use alternative precautionsfor at least five days after using this product.

Side effects:

Clotrimazole is well tolerated when applied topically. Rarely patients may experience local mild burning or irritation, immediately after applying cream. No systemic effects are to be expected on external application.

Symptoms and treatment of overdosage:

Clinically important toxicity would not be anticipated and no specific treatment is required.

Storage:

Store under normal storage conditions (15°C to 30°C).

Protect from light.

Keep out of reach of children.

Shelf life: 36 months.

Presentation:

20 gm tube in a carton



Manufactured in India by:

AGOG PHARMA LTD.

Plot No. 33, Sector II, The Vasai Taluka Indl. Co-op. Estate Ltd., Vasai (E), Dist. Thane. INDIA.