





LEAFLET

FOLON TABLETS

(Folic Acid Tablets BP 5 mg)

Yellow coloured circular flat bevelled edge uncoated tablets having breakline on one side and plain on the other side.

Label Claim

Each uncoated tablet cointains:

Folic acid BP 5 mg.

Excipients

q.s.

Clinical Pharmacology

After conversion into coenzyme forms it is concerned in single carbon unit transfer in the synthesis of purines, pyrimidines, and methionine.

Indication(s)

Prophylaxis to prevent folate deficiency in prematurity. Maintenance of adequate diatary folate requirement in pregnacy. Prophylaxis to prevent folate deficiency in chronic myelofibrosis, Maintenance of adequate dietary folate intake in chronic hemolytic states. Prophylaxis to prevent folate loss caused by renal dialysis, Prophylaxis to prevent neural tube defects in pregnancy, Netritional megaloblastic anemia, planning Megaloblastic anemia in pregnancy, Megaloblastic anemia caused by celiac disease. Megaloblastic anemia caused by hemolytic anemia, Treatment of folate deficiency in chronic myelofibrosis, Megaloblastic anemia caused by exfoliative dermatitis, Megaloblastic anemia as a result of congenital folate malabsorption, Megaloblastic anemia in infants reared on goat's milk, Megaloblastic anemia caused by tropical spruge, Megaloblastic anemia caused by anticonvulsant drug therapy. Treatment of sideroblastic anemia in the elderly. Treatment of folate deficiency in scury, Megaloblastic anemia related to antimalarial drug theraphy, Megaloblastic anemia resulting from a long-term course of methptrexate, Treatment of megaloblastic anemia caused by long-term sulfamethoxazole and trimethoprim combination prophylaxis.



Pharma Ltd.





(WHO - GMP CERTIFIED - GOVT RECOGNISED EXPORT HOUSE)

Dosage and Administration (directions for use) As directed by the physician.

Untreated cobalamin deficiency. Addisonian pernicious anemia or other vitamin B₁₂ deficiency, Malignant disease.

Adverse relations/Side effects

Such effect are rare. A few subjects have become sensitized to folic acid, and administration has been followed within an hour by an itchy skin eruption, malaise, and bronchospasm. In such cases, intradermal tests have been positive with folic acid but negative with leucovorin. Precipitation of status epilepticus may occur. When given long term to patients with cobalamin deficiency, the drug can precipitate a neuropathy.

Storage: Store under normal storage conditions (15°C - 30°C). Protect from light. Keep medicine out of reach of children.

Presentation:

A Bulk pack of 1000's Tablets.

Blister pack of 5 X 10 Tablets.
Blister pack of 10 X 10 Tablets.

Blister pack of 100 X 10 Tablets.

Manufactured in India by:



Plot No. 33, Sector – II, The Vasai Taluka Indl. Co-op. Estate Ltd., Vasai (East), Dist. Thane. INDIA.