

GLYKA MR 60 MG TABLETS
(Gliclazide Modified Release 60 mg Tablets)

1.6.1 SUMMARY OF PRODUCT CHARACTERISTICS

1. Name of the Medicinal Product:

Glyka MR Tablets 60 mg

2. Qualitative & Quantitative Composition:

Each modified release tablet contains Gliclazide 60 mg.

3. Pharmaceutical Form:

Modified Release Tablet. White oblong uncoated tablets, scored on both side, with 'GL 60' engraved on one side plain on other side

4. Clinical Particulars:

4.1 Therapeutic indications

Non-insulin-dependent diabetes (type 2) in adults when dietary measures, physical exercise and weight loss alone is not sufficient to control blood glucose.

4.2 Posology and method of administration

Posology

The daily dose of Gliclazide 60 mg MR Tablets may vary from one half to 2 tablets per day, i.e. from 30 to 120 mg taken orally in a single intake at breakfast time.

It is recommended to swallow the dose without crushing or chewing.

If a dose is forgotten, there must be no increase in the dose taken the next day.

As with any hypoglycaemic agent, the dose should be adjusted according to the individual patient's metabolic response (blood glucose, HbA1c).

Initial dose

The recommended starting dose is 30 mg daily (half a tablet of Glyka MR Tablets 60 mg). If blood glucose is effectively controlled, this dose may be used for maintenance treatment.

GLYKA MR 60 MG TABLETS

(Gliclazide Modified Release 60 mg Tablets)

If blood glucose is not adequately controlled, the dose may be increased to 60, 90 or 120 mg daily, in successive steps. The interval between each dose increment should be at least 1 month except in patients whose blood glucose has not reduced after two weeks of treatment. In such cases, the dose may be increased at the end of the second week of treatment.

The maximum recommended daily dose is 120 mg.

One Gliclazide 60 mg MR Tablets modified release tablet is equivalent to two Gliclazide 30 mg MR Tablets modified release tablets. The breakability of the Gliclazide 60 mg MR Tablets modified release tablet enables flexibility of dosing to be achieved.

Switching from another oral antidiabetic agent to Gliclazide 30 mg MR Tablets:

Gliclazide 60 mg MR Tablets can be used to replace other oral antidiabetic agents.

The dosage and the half-life of the previous antidiabetic agent should be taken into account when switching to Gliclazide 60 mg MR Tablets.

A transitional period is not generally necessary. A starting dose of 30 mg should be used and this should be adjusted to suit the patient's blood glucose response, as described above.

When switching from a hypoglycaemic sulfonylurea with a prolonged half-life, a treatment free period of a few days may be necessary to avoid an additive effect of the two products, which might cause hypoglycaemia. The procedure described for initiating treatment should also be used when switching to treatment with Gliclazide 60 mg MR Tablets, i.e. a starting dose of 30 mg/day, followed by a stepwise increase in dose, depending on the metabolic response.

Combination treatment with other antidiabetic agents:

Gliclazide 60 mg MR Tablets can be given in combination with biguanides, alpha glucosidase inhibitors or insulin. In patients not adequately controlled with Gliclazide 60 mg MR, concomitant insulin therapy can be initiated under close medical supervision.

Special Populations

Elderly

Gliclazide 60 mg MR should be prescribed using the same dosing regimen recommended for patients under 65 years of age.

GLYKA MR 60 MG TABLETS **(Gliclazide Modified Release 60 mg Tablets)**

Renal impairment

In patients with mild to moderate renal insufficiency, the same dosing regimen can be used as in patients with normal renal function with careful patient monitoring. These data have been confirmed in clinical trials.

Patients at risk of hypoglycaemia

- Undernourished or malnourished,
- Severe or poorly compensated endocrine disorders (hypopituitarism, hypothyroidism, adrenocorticotrophic insufficiency),
- Withdrawal of prolonged and/or high dose corticosteroid therapy,
- Severe vascular disease (severe coronary heart disease, severe carotid impairment, diffuse vascular disease); It is recommended that the minimum daily starting dose of 30 mg is used.

Paediatric population

The safety and efficacy of Gliclazide 60 mg MR in children and adolescents have not been established. No data are available

4.3 Contraindications

This medicine is contra-indicated in case of:

- Hypersensitivity to gliclazide or to any of the excipients listed in section 6.1, other sulfonylureas, sulfonamides,
- Type 1 diabetes,
- Diabetic pre-coma and coma, diabetic keto-acidosis,
- Severe renal or hepatic insufficiency: in these cases the use of insulin is recommended,
- Treatment with miconazole (see Section 4.5)
- Lactation (see Section 4.6).

4.4 Special warning & precautions for use

Hypoglycaemia:

This treatment should be prescribed only if the patient is likely to have a regular food intake (including breakfast). It is important to have a regular carbohydrate intake due to the increased risk

GLYKA MR 60 MG TABLETS **(Gliclazide Modified Release 60 mg Tablets)**

of hypoglycaemia if a meal is taken late, if an inadequate amount of food is consumed or if the food is low in carbohydrate. Hypoglycaemia is more likely to occur during low-calorie diets, following prolonged or strenuous exercise, alcohol intake or if a combination of hypoglycaemic agents is being used.

Hypoglycaemia may occur following administration of sulfonylureas (see Section 4.8). Some cases may be severe and prolonged. Hospitalisation may be necessary and glucose administration may need to be continued for several days. Careful selection of patients, of the dose used, and clear patient directions are necessary to reduce the risk of hypoglycaemic episodes

Factors which increase the risk of hypoglycaemia:

- patient refuses or (particularly in elderly subjects) is unable to co-operate,
- malnutrition, irregular mealtimes, skipping meals, periods of fasting or dietary changes,
- imbalance between physical exercise and carbohydrate intake,
- renal insufficiency,
- severe hepatic insufficiency,
- overdose of Gliclazide,
- certain endocrine disorders: thyroid disorders, hypopituitarism and adrenal insufficiency,
- concomitant administration of certain other medicines (see Section 4.5).

Renal and hepatic insufficiency: the pharmacokinetics and/or pharmacodynamics of gliclazide may be altered in patients with hepatic insufficiency or severe renal failure. A hypoglycaemic episode occurring in these patients may be prolonged, so appropriate management should be initiated.

Patient information: The risks of hypoglycaemia (see section 4.8), together with its symptoms treatment, and conditions that predispose to its development, should be explained to the patient and to family members. The patient should be informed of the importance of following dietary advice, of taking regular exercise, and of regular monitoring of blood glucose levels.

Blood glucose control in a patient receiving antidiabetic treatment may be affected by any of the following: St. John's Wort (*Hypericum perforatum*) preparations (see section 4.5), fever, trauma, infection or surgical intervention. In some cases, it may be necessary to administer insulin.

GLYKA MR 60 MG TABLETS **(Gliclazide Modified Release 60 mg Tablets)**

The hypoglycaemic efficacy of any oral antidiabetic agent, including gliclazide, is attenuated over time in many patients: this may be due to progression in the severity of the diabetes, or to a reduced response to treatment. This phenomenon is known as secondary failure which is distinct from primary failure, when an active substance is ineffective as first-line treatment. Adequate dose adjustment and dietary compliance should be considered before classifying the patient as secondary failure.

Dysglycaemia:

Disturbances in blood glucose, including hypoglycaemia and hyperglycaemia have been reported, in diabetic patients receiving concomitant treatment with fluoroquinolones, especially in elderly patients. Indeed, careful monitoring of blood glucose is recommended in all patients receiving at the same time Gliclazide Modified Release Tablets 60 mg and a fluoroquinolone.

Laboratory tests: Measurement of glycated haemoglobin levels (or fasting venous plasma glucose) is recommended in assessing blood glucose control. Blood glucose self-monitoring may also be useful.

Treatment of patients with G6PD-deficiency with sulphonylurea agents can lead to haemolytic anaemia. Since gliclazide belongs to the chemical class of sulphonylurea drugs, caution should be used in patients with G6PD-deficiency and a non-sulphonylurea alternative should be considered.

Excipients: Gliclazide 60 mg MR Tablet should not be administered to patients with rare hereditary problems of galactose intolerance, the Lapp lactase deficiency or glucose-galactose malabsorption.

4.5 Interaction with other medicinal products & other forms of interaction

The following products are likely to increase the risk of hypoglycaemia

Contra-indicated combination

Miconazole (systemic route, oromucosal gel): increases the hypoglycaemic effect with possible onset of hypoglycaemic symptoms, or even coma.

Combinations which are not recommended

Phenylbutazone (systemic route): increases the hypoglycaemic effect of sulphonylurea (displaces their binding to plasma proteins and/or reduces their elimination).

It is preferable to use a different anti-inflammatory agent, or else to warn the patient and emphasise the importance of self-monitoring. Where necessary, adjust the dose during and after treatment with the anti-inflammatory agent.

GLYKA MR 60 MG TABLETS **(Gliclazide Modified Release 60 mg Tablets)**

Alcohol: increases the hypoglycaemic reaction (by inhibiting compensatory reactions) that can lead to the onset of hypoglycaemic coma.

Alcohol or medicinal products containing alcohol should be avoided.

Combinations requiring precautions for use

Potential of the blood glucose lowering effect and thus, in some instances, hypoglycaemia may occur when one of the following drugs is taken: other antidiabetic agents (insulins, acarbose, metformin, thiazolidinediones, dipeptidyl peptidase-4 inhibitors, GLP-1 receptor agonists), beta-blockers, fluconazole, angiotensin converting enzyme inhibitors (captopril, enalapril), H₂-receptor antagonists, MAOIs, sulfonamides, clarithromycin and nonsteroidal anti-inflammatory agents.

The following products may cause an increase in blood glucose levels

Combination which is not recommended

Danazol: diabetogenic effect of danazol.

If the use of this active substance cannot be avoided, warn the patient and emphasise the importance of urine and blood glucose monitoring. It may be necessary to adjust the dose of the antidiabetic agent during and after treatment with danazol.

Combinations requiring precautions during use

Chlorpromazine (neuroleptic agent): high doses (>100 mg per day of chlorpromazine) increase blood glucose levels (reduced insulin release).

Warn the patient and emphasise the importance of blood glucose monitoring. It may be necessary to adjust the dose of the antidiabetic active substance during and after treatment with the neuroleptic agent.

Glucocorticoids (systemic and local route: intra-articular, cutaneous and rectal preparations) and tetracosactrin: increase in blood glucose levels with possible ketosis (reduced tolerance to carbohydrates due to glucocorticoids).

Warn the patient and emphasise the importance of blood glucose monitoring, particularly at the start of treatment. It may be necessary to adjust the dose of the antidiabetic active substance during and after treatment with glucocorticoids.

GLYKA MR 60 MG TABLETS **(Gliclazide Modified Release 60 mg Tablets)**

Ritodrine, salbutamol, terbutaline: (I.V.)

Increased blood glucose levels due to beta-2 agonist effects.

Emphasise the importance of monitoring blood glucose levels. If necessary, switch to insulin.

Saint John's Wort (*Hypericum perforatum*) preparations:

Gliclazide exposure is decreased by Saint John's Wort- *Hypericum perforatum*. Emphasize the importance of blood glucose levels monitoring.

The following products may cause dysglycaemia

Combinations requiring precautions during use

Fluoroquinolones: in case of a concomitant use of Gliclazide 60 mg MR Tablets and a fluoroquinolone, the patient should be warned of the risk of dysglycaemia, and the importance of blood glucose monitoring should be emphasized.

Combination which must be taken into account

Anticoagulant therapy (e.g. Warfarin...):

Sulfonylureas may lead to potentiation of anticoagulation during concurrent treatment.

Adjustment of the anticoagulant may be necessary.

4.6 Pregnancy and Lactation

Pregnancy

There is no experience with the use of gliclazide during pregnancy in humans, even though there are few data with other sulphonylurea.

In animal studies, gliclazide is not teratogenic. Control of diabetes should be obtained before the time of conception to reduce the risk of congenital abnormalities linked to uncontrolled diabetes.

Oral hypoglycaemic agents are not suitable, insulin is the drug of first choice for treatment of diabetes during pregnancy. It is recommended that oral hypoglycaemic therapy is changed to insulin before a pregnancy is attempted, or as soon as pregnancy is discovered.

Breastfeeding

GLYKA MR 60 MG TABLETS **(Gliclazide Modified Release 60 mg Tablets)**

It is not known whether gliclazide or its metabolites are excreted in breast milk. Given the risk of neonatal hypoglycaemia, the product is contraindicated in breast-feeding mother.

4.7 Effects on ability to drive & use machines

Gliclazide MR Tablets 60 mg has no known influence on the ability to drive and use machines. However, patients should be made aware of the symptoms of hypoglycaemia and should be careful if driving or operating machinery, especially at the beginning of treatment.

4.8 Undesirable Effects

Based on the experience with gliclazide, the following undesirable effects have been reported.

The most frequent adverse reaction with gliclazide is hypoglycaemia

As for other sulfonylureas, treatment with Gliclazide Modified Release can cause hypoglycaemia, if mealtimes are irregular and, in particular, if meals are skipped. Possible symptoms of hypoglycaemia are: headache, intense hunger, nausea, vomiting, lassitude, sleep disorders, agitation, aggression, poor concentration, reduced awareness and slowed reactions, depression, confusion, visual and speech disorders, aphasia, tremor, paresis, sensory disorders, dizziness, feeling of powerlessness, loss of self-control, delirium, convulsions, shallow respiration, bradycardia, drowsiness and loss of consciousness, possibly resulting in coma and lethal outcome.

In addition, signs of adrenergic counter-regulation may be observed: sweating, clammy skin, anxiety, tachycardia, hypertension, palpitations, angina pectoris and cardiac arrhythmia.

Usually, symptoms disappear after intake of carbohydrates (sugar). However, artificial sweeteners have no effect. Experience with other sulfonylurea shows that hypoglycaemia can recur even when measures prove effective initially.

If a hypoglycaemic episode is severe or prolonged, and even if it is temporarily controlled by intake of sugar, immediate medical treatment or even hospitalisation is required.

Other undesirable effects:

Gastrointestinal disturbances, including abdominal pain, nausea, vomiting dyspepsia, diarrhoea, and constipation have been reported: if these should occur they can be avoided or minimised if gliclazide is taken with breakfast.

The following undesirable effects have been more rarely reported:

- Skin and subcutaneous tissue disorders: rash, pruritus, urticaria, angioedema, erythema, maculopapular rashes, bullous reactions (such as Stevens-Johnson syndrome and toxic

GLYKA MR 60 MG TABLETS

(Gliclazide Modified Release 60 mg Tablets)

epidermal necrolysis), and exceptionally, drug rash with eosinophilia and systemic symptoms (DRESS).

- Blood and lymphatic system disorders: Changes in haematology are rare. They may include anaemia, leucopenia, thrombocytopenia, granulocytopenia. These are in general reversible upon discontinuation of medication.
- Hepato-biliary disorders: raised hepatic enzyme levels (AST, ALT, alkaline phosphatase), hepatitis (isolated reports). Discontinue treatment if cholestatic jaundice appears.

These symptoms usually disappear after discontinuation of treatment.

- Eye disorders: transient visual disturbances may occur especially on initiation of treatment, due to changes in blood glucose levels.
- Class attribution effects: As for other sulfonylureas, the following adverse events have been observed: cases of erythrocytopenia, agranulocytosis, haemolytic anaemia, pancytopenia, allergic vasculitis, hyponatraemia, elevated liver enzyme levels and even impairment of liver function (*e.g.* with cholestasis and jaundice) and hepatitis which regressed after withdrawal of the sulfonylurea or led to life-threatening liver failure in isolated cases.

4.9. Overdose

An overdose of sulphonylurea may cause hypoglycaemia.

Moderate symptoms of hypoglycaemia, without any loss of consciousness or neurological signs, must be corrected by carbohydrate intake, dose adjustment and/or change of diet. Strict monitoring should be continued until the doctor is sure that the patient is out of danger.

Severe hypoglycaemic reactions, with coma, convulsions or other neurological disorders are possible and must be treated as a medical emergency, requiring immediate hospitalisation. If hypoglycaemic coma is diagnosed or suspected, the patient should be given a rapid I.V. injection of 50 mL of concentrated glucose solution (20 to 30 %). This should be followed by continuous infusion of a more dilute glucose solution (10 %) at a rate that will maintain blood glucose levels above 1 g/L. Patients should be monitored closely and, depending on the patient's condition after this time, the doctor will decide if further monitoring is necessary. Dialysis is of no benefit to patients due to the strong binding of gliclazide to proteins.

GLYKA MR 60 MG TABLETS
(Gliclazide Modified Release 60 mg Tablets)

5. Pharmacological Properties:

5.1. Pharmacodynamics:

Pharmacotherapeutic group: Sulphonamides, Urea derivatives

ATC code: **A10BB09**.

Mode of action

Gliclazide is a hypoglycaemic sulphonylurea oral antidiabetic active substance differing from other related compounds by an N-containing heterocyclic ring with an endocyclic bond. Gliclazide reduces blood glucose levels by stimulating insulin secretion from the β -cells of the islets of Langerhans. Increase in postprandial insulin and C peptide secretion persists after two years of treatment. In addition to these metabolic properties, gliclazide has haemovascular properties.

Haemovascular properties:

Gliclazide decreases microthrombosis by two mechanisms which may be involved in complications of diabetes:

a partial inhibition of platelet aggregation and adhesion, with a decrease in the markers of platelet activation (beta thromboglobulin, thromboxane B2).

an action on the vascular endothelium fibrinolytic activity with an increase in tPA activity.

5.2. Pharmacokinetics:

Absorption

Plasma levels increase progressively during the first 6 hours, reaching a plateau which is maintained from the sixth to the twelfth hour after administration. Intra-individual variability is low. Gliclazide is completely absorbed. Food intake does not affect the rate or degree of absorption.

GLYKA MR 60 MG TABLETS **(Gliclazide Modified Release 60 mg Tablets)**

Distribution

Plasma protein binding is approximately 95%. The volume of distribution is around 30 liters. A single daily intake of Gliclazide 60 mg MR maintains effective gliclazide plasma concentrations over 24 hours.

Biotransformation

Gliclazide is mainly metabolised in the liver and excreted in the urine: less than 1% of the unchanged form is found in the urine. No active metabolites have been detected in plasma.

Elimination

The elimination half-life of gliclazide varies between 12 and 20 hours.

Linearity/non-linearity

The relationship between the dose administered ranging up to 120 mg and the area under the concentration time curve is linear.

Special populations

Elderly

No clinically significant changes in pharmacokinetic parameters have been observed in elderly patients.

53. Preclinical Safety Data

Preclinical data reveal no special hazards for humans based on conventional studies of repeated dose toxicity and genotoxicity. Long term carcinogenicity studies have not been done. No teratogenic changes have been shown in animal studies, but lower foetal body weight was observed in animals receiving doses 25 fold higher than the maximum recommended dose in humans. Fertility and reproductive performance were unaffected after gliclazide administration in animal studies.

GLYKA MR 60 MG TABLETS
(Gliclazide Modified Release 60 mg Tablets)

6. Pharmaceutical Particulars:

6.1. List of excipients:

| S. No. | Ingredient | Function |
|--------|----------------------------|-------------------|
| 1. | Lactose Monohydrate | Filler/Diluent |
| 2. | Hypromellose | Release Retardant |
| 3. | Maltodextrin | Filler/Diluent |
| 4. | Colloidal Anhydrous Silica | Glidant |
| 5. | Magnesium Stearate | Lubricant |

6.2. Incompatibilities:

Not Applicable

6.3. Shelf life:

36 months

6.4. Special precautions for storage:

Store below 30°C.

6.5. Nature & contents of container:

Alu/Alu blister placed in a printed carton along with a pack insert. 30's pack (3×10's Blister)

6.6 Manufacturer:

Neopharma, Abu Dhabi, UAE

Plot No. A1 89-95,

Industrial City of Abu Dhabi (ICAD),

Mussafah, Abu Dhabi, UAE

GLYKA MR 60 MG TABLETS
(Gliclazide Modified Release 60 mg Tablets)

6.7 Marketing Authorization Holder:

Neopharma, Abu Dhabi, UAE

Plot No. A1 89-95,

Industrial City of Abu Dhabi (ICAD),

Mussafah, Abu Dhabi, UAE

Date of Revision of Text:

July 2019