

Description <u>Description</u>

Ibumex tablets are available as round, sugar-coated, pink coloured tablets containing 200mg of ibuprofen BP.

Ibumex suspension is an orange coloured, pleasandy flavoured liquid containing 100mg of ibuprofen BP in 5ml.

Indications.

Ibuprofen is a non-steroidal anti-inflammatory drug with buprofen is a not-inflammatory and anti-pyretic actions. Ibuprofen tablets are used for the relief of pain associated with low backache, tendinitis and frozen shoulder. It can with low backache, tendinitis and frozen shoulder is can take the transfer of the profession of the state of the part of the profession of the part of the profession of the profes with low backache, tendinitis and frozen shoulder. It can also be used in soft tissue injuries like sprains and strains. The analgesic effect of ibuprofen makes it of use to relieve pain associated with mild to moderate toothache and headache. Ibuprofen suspension is indicated for the reduction of fever and for the relief of mild to moderate pain in children older than one year. pain in children older than one year.

Dosage and Administration

Ibuprofen should be taken three to four times a day, preferably after food, as indicated below.

Adults and Children over 12 years

Adults and California to the Landon One to two lournes to blets
One to two lournes to blets
One to two lournes to blets
I had en and or 11 jea to
lourner suspension is recomment
8-12 yrs Two 5 mt poorfuls

COP 3 yrs

One 5 mil spoonful Half a.5 mil spoonful

Underlyr Not recommended Or as directed by a doctor

DO NOT EXCEED THE STATED DOSE

If symptoms persists seek medical advice

Contra-indications, Warnings etc Ibuprofen should NOT be taken by patients with severe or active peptic ulceration, or with a history of it.

- Precautions

 Medical advice should be sought if suffering from astima or aspirin sensitivity, before taking ibuprofen.

 Pregnant or lactating women should take the advice of a doctor or pharmacist before taking ibuprofen.

Side effects

The side effects of [buprofen include dyspepsia, gastro-lintestinal intolerance, bleeding, skin rashes and oedema.

Storage | Storag

place. Keep out of reach of children.

Package Quantities
20's and 100's (Bilster Packad)
100's (Bilster Packad)
1000's

60ml and 100ml ibumex Suspension:

Bef: P07/02/1

REGAL PHARMACEUTICALS LTD.
Plot No. 7879/18, Off Baba Dogo Road, Ruaraka
RO. Box 44421 - 00100 GPO, Nairobi