



**AGOG Pharma Ltd.**

(WHO - GMP CERTIFIED - GOVT RECOGNISED EXPORT HOUSE)



CD# 11-059 13

**Regd. Office & Factory :** Plot No. 33, Sector II, The Vasai Taluka Industrial Co-op Estate Ltd. Gaurapada, Vasai (E), Dist. Thane - 401 208, INDIA.  
Tel. : 95250 - 2455801 / 2452714 / 2453525 • Fax : 95250 - 2452074 (0091 - 250 - 2452074) • Email : agog@vsnl.net & agogpharma@rediffmail.com

## LEAFLET

### TRIMAGO

(Co-trimoxazole Suspension & Tablets BP)

#### Contents :

##### Tablets :

Trimago tablets are white biconvex uncoated tablets each containing 80 mg Trimethoprim BP and 400 mg, Sulfamethoxazole BP.  
Suspension : Trimago Suspension is available as Orange coloured Suspension containing 200 mg Sulfamethoxazole BP and 40 mg Trimethoprim BP per 5ml.

#### Pharmacological actions :

The combination of Trimethoprim and Sulfamethoxazole, acts synergistically to produce a bactericidal or bacteriostatic action against a wide range of gram-positive and gram - negative organisms and some protozoa. Both antibiotics acts at different points of the folate metabolic pathway.

#### Indications :

Trimago is an antibacterial agent, effective against wide range of gram-positive and gram - negative organisms. Trimago is indicated for the treatment of respiratory tract, genito-urinary tract, gastro intestinal tract, skin and other bacterial infections.

#### Dosage and Administration :

Trimago should be taken preferably after food, minimize the possibilities of gastro-intestinal disturbances, as indicated below.

**Adults and Children over 12 years :**  
2 tablets every 12 hours.

#### Children :

6-12 years: 10 ml every 12 hours, or 1 tablet twice a day.

6 months - 5 years : 5ml every 12 hours.

8 weeks - 5 months: 2.5ml every 12 hours.

For severe infections in all age groups the above dosage may be increased by 50%.

In acute infections, Trimago should be taken for at least five days or until the patient has been symptom free for two days or as directed by the physician.

#### Contra-indications :

Trimago should not be taken by patients with a history of sensitivity to Sulphonamide, Trimethoprim or Co-trimoxazole. Trimago should be avoided in patients with marked liver damage. Trimago should not be given to premature babies or to full term infants during their first six weeks of life. Regular monthly blood counts are advisable when Trimago is given for long periods.

#### Precautions :

Pregnant or lactating women should take the advice of a doctor, before taking Trimago.

#### Side Effects :

1. The side effect of Trimago include nausea, vomiting, diarrhoea and skin rashes. Trimago should be stopped if rash develops.
2. Severe but rare side effect include depression, dizziness and blood dyscrasias.

#### Storage :

Store under normal storage conditions (15°C to 30°C)

Protect from light. Keep all Medicines out of reach of children.  
For Suspension : Shake well Before Use.

#### Legal Category :

Prescription Only Medicine (POM).

#### PRESENTATION :

Trimago Tablets : Blister pack of 10 X 10 tablets  
Blister pack of 100 X 10 tablets  
Bulk pack of 1000 tablets / 250 tablets

Trimago Suspension : 60 ml / 100 ml in amber coloured pet bottles.

Manufactured in India by



**AGOG PHARMA LTD.**

Plot No. 33, Sector II, The Vasai Taluka Industrial Co-op. Estate Ltd.,  
Vasai (E), Dist. Thane, INDIA



**AGOG Pharma Ltd.**

(WHO - GMP CERTIFIED - GOVT RECOGNISED EXPORT HOUSE)



CGN: 11-099 13

**Regd. Office & Factory :** Plot No. 33, Sector II, The Vasai Taluka Industrial Co-op. Estate Ltd. Gaurai-pada, Vasai (E), Dist. Thane - 401 208. INDIA.  
Tel. : 95250 - 2455801 / 2452714 / 2453525 • Fax : 95250 - 2452074 (0091 - 250 - 2452074) • Email : agog@vsnl.net & agogpharma@rediffmail.com

### 1.5.3 Patient Information Leaflet (PIL)

#### TRIMAGO

(Co-trimoxazole Suspension & Tablets BP)

##### Contents :

##### Tablets :

Trimago tablets are white biconvex uncoated tablets each containing 80 mg Trimethoprim BP and 400 mg. Sulfamethoxazole BP.  
Suspension : Trimago Suspension is available as Orange coloured Suspension containing 200 mg Sulfamethoxazole BP and 40 mg Trimethoprim BP per 5ml.

##### Pharmacological actions :

The combination of Trimethoprim and Sulfamethoxazole, acts synergistically to produce a bactericidal or bacteriostatic action against a wide range of gram-positive and gram - negative organisms and some protozoa. Both antibiotics acts at different points of the folate metabolic pathway.

##### Indications :

Trimago is an antibacterial agent, effective against wide range of gram-positive and gram - negative organisms. Trimago is indicated for the treatment of respiratory tract, genito-urinary tract, gastro intestinal tract, skin and other bacterial infections.

##### Dosage and Administration :

Trimago should be taken preferably after food, minimize the possibilities of gastro-intestinal disturbances, as indicated below.

##### Adults and Children over 12 years :

2 tablets every 12 hours.

##### Children :

6-12 years : 10 ml every 12 hours, or 1 tablet twice a day.

6 months - 5 years : 5ml every 12 hours.

8 weeks - 5 months : 2.5ml every 12 hours.

For severe infections in all age groups the above dosage may be increased by 50%.

In acute infections, Trimago should be taken for at least five days or until the patient has been symptom free for two days or as directed by the physician.

##### Contra-indications :

Trimago should not be taken by patients with a history of sensitivity to Sulphonamide, Trimethoprim or Co-trimoxazole. Trimago should be avoided in patients with marked liver damage. Trimago should not be given to premature babies or to full term infants during their first six weeks of life. Regular monthly blood counts are advisable when Trimago is given for long periods.

##### Precautions :

Pregnant or lactating women should take the advice of a doctor, before taking Trimago.

##### Side Effects :

1. The side effect of Trimago include nausea, vomiting, diarrhoea and skin rashes. Trimago should be stopped if rash develops.
2. Severe but rare side effect include depression, dizziness and blood dyscrasias.

##### Storage :

Store under normal storage conditions (15°C to 30°C)  
Protect from light. Keep all Medicines out of reach of children.  
For Suspension : Shake well Before Use.

##### Legal Category :

Prescription Only Medicine (POM).

##### PRESENTATION :

Trimago Tablets : Blister pack of 10 X 10 tablets  
Blister pack of 100 X 10 tablets  
Bulk pack of 1000 tablets / 250 tablets

Trimago Suspension : 60 ml / 100 ml in amber coloured pet bottles.

Manufactured in India by



**AGOG PHARMA LTD.**

Plot No. 33, Sector II, The Vasai Taluka Industrial Co-op. Estate Ltd.,  
Vasai (E), Dist. Thane, INDIA