



AGQG Pharma Ltd.

(WHO - GMP CERTIFIED - GOVT RECOGNISED EXPORT HOUSE)



CGN: 11-05913

Regd. Office & Factory : Plot No. 33, Sector II, The Vasai Taluka Industrial Co-op. Estate Ltd. Gauripada, Vasai (E), Dist. Thane - 401 208. INDIA.
Tel. : 95250 - 2455801 / 2452714 / 2453525 • Fax : 95250 - 2452074 (0091 - 250 - 2452074) • Email : agog@vsnl.net & agogpharma@rediffmail.com

Brand Name : TRIMAGO ORAL SUSPENSION	
Generic Name : Paediatric Co-Trimoxazole Oral Suspension BP	2021
Module 1 Administrative Information and Product Information	
1.5 Product Information	Confidential

1.5 PRODUCT INFORMATION

1.5.1 Prescribing Information (Summary of Products Characteristics)

1. NAME OF DRUG PRODUCT

1. Name of drug product

Paediatric Co-Trimoxazole Oral Suspension BP

1.1 (Trade) name of product

TRIMAGO ORAL SUSPENSION

1.2 Strength

Each 5 ml contains:
Sulfamethoxazole 200 mg
Trimethoprim 40 mg

1.3 Pharmaceutical Dosage Form

Oral Liquid Suspension

**AQOQ Pharma Ltd.****(WHO - GMP CERTIFIED - GOVT RECOGNISED EXPORT HOUSE)**

CIN: 11-09913

Regd. Office & Factory : Plot No. 33, Sector II, The Vasai Taluka Industrial Co-op. Estate Ltd. Gaurapada, Vasai (E), Dist. Thane - 401 208. INDIA.
 Tel. : 95250 - 2455801 / 2452714 / 2453525 • Fax : 95250 - 2452074 (0091 - 250 - 2452074) • Email : aqoq@vsnl.net & aqoqpharma@rediffmail.com

2. QUALITATIVE AND QUANTITATIVE COMPOSITIONS

2.1 Qualitative Declaration

Each 5 ml contains:

Sulfamethoxazole 200 mg

Trimethoprim 40 mg

2.2 Quantitative Declaration

Ingredients	Specification	Label Claim	Overages	Qty. / liq.
<u>ACTIVE</u>				
Sulfamethoxazole	BP	200 mg	-	204.0 mg
Trimethoprim	BP	40 mg	-	48.80 mg
<u>NON ACTIVE</u>				
Methyl Paraben Sodium	BP	-	-	7.50 mg
Propyl Paraben Sodium	BP	-	-	2.50 mg
Carboxy methyl Cellulose Sodium	BP	-	-	40.0 mg
Sodium Saccharin	BP	-	-	2.50 mg
Citric acid monohydrate	BP	-	-	5.00 mg
Sodium Benzoate	BP	-	-	7.50 mg
Colour sunset yellow supra	Inhouse	-	-	0.16 mg
Essence Orange Sweet No.1	Inhouse	-	-	0.01 ml
Sucrose	BP	-	-	1500 mg
Purified Water	Inhouse	-	-	Q.S to 2000 ml

BP 2019 = British Pharmacopoeia 2019.



AGOG Pharma Ltd.

(WHO - GMP CERTIFIED - GOVT RECOGNISED EXPORT HOUSE)



CPN: 11-09918

Regd. Office & Factory : Plot No. 33, Sector II, The Vasai Taluka Industrial Co-op. Estate Ltd. Gauripada, Vasai (E), Dist. Thane - 401 208. INDIA.
Tel. : 95250 - 2455801 / 2452714 / 2453525 • Fax : 95250 - 2452074 (0091 - 250 - 2452074) • Email : agog@vsnl.net & agogpharma@rediffmail.com

3. PHARMACEUTICAL DOSAGE FORM

Liquid Oral Suspension

Light orange coloured, thick, uniform suspension on shaking.



4. CLINICAL PARTICULARS

4.1 Therapeutic Indications

Co-trimoxazole is indicated in adults and children (>12 to <18 years old) and adults (>18 years old).

Co-trimoxazole indicated for the treatment of the following infections when owing to sensitive organisms.

- Treatment and prevention of *Pneumocystis jiroveci* pneumonitis or 'PJP'.
- Treatment and prophylaxis of toxoplasmosis
- Treatment of nocardiosis.

The following infections may be treated with co-trimoxazole where there is bacterial evidence of sensitivity to co-trimoxazole and good reason to prefer the combination of antibiotics in co-trimoxazole to a single antibiotic:

- Acute uncomplicated urinary tract infection
- Acute otitis media
- Acute exacerbation of chronic bronchitis

Consideration should be given to official guidance on the appropriate use of antibacterial agents.

4.2 Posology and Method of Administration

General Dosage Recommendations

Where dosage is expressed as "Liquid" this refers to the Paediatric suspension, i.e. 40 mg Trimethoprim BP and 200 mg Sulfamethoxazole BP. If other formulations are to be used appropriate adjustment should be made.

Standard dosage recommendations for acute infections

Children over 12 years old (>12 to <18 years old):

The standard dosage for children is equivalent to approximately 6 mg trimethoprim and 30 mg sulfamethoxazole per kg body weight per day, given in two equally divided doses. The schedules for children are according to the child's age and provided in the table below:

Age	Tablets
>12 to <18 years old	10 ml every 12 hours

Treatment should be continued until the patient has been symptom free for two days; the majority will require treatment for at least 5 days. If clinical improvement is not evident after 7 days therapy, the patient should be reassessed.



As an alternative to Standard Dosage for acute uncomplicated lower urinary tract infections, short-term therapy of 1 to 3 days duration has been shown to be effective.

Impaired hepatic function:

No data are available relating to dosage in patients with impaired hepatic function.

Pneumocystis jirovecii pneumonitis

Treatment - Children (>12 to <18 years old)

A higher dosage is recommended, using 20 mg trimethoprim and 100 mg sulfamethoxazole per kg of body weight per day in two or more divided doses for two weeks. The aim is to obtain peak plasma or serum levels of trimethoprim of greater than or equal to 5 microgram/ml (verified in patients receiving 1-hour infusions of intravenous Co-Trimoxazole).

Prevention - Children (>12 to <18 years old):

The standard dosage for children is equivalent to approximately 6 mg trimethoprim and 30 mg sulfamethoxazole per kg body weight per day, given in two equally divided doses. The following dose schedules may be used for the duration of the period at risk:

Age	Tablets
6-12 years old	10 ml every 12 hours
6 months - 5 years old	5 ml every 12 hours
8 weeks – 5 months old	2.5 ml every 12 hours

Method of administration:

Oral.

It may be preferable to take Co-Trimoxazole with some food or drink to minimise the possibility of gastrointestinal disturbances.

4.3 Contraindications

- Hypersensitivity to the active substance, sulfonamides, trimethoprim, Co-trimoxazole or to any of the Excipients.
- Contra-indicated in patients showing marked severe liver parenchymal damage.
- Contra-indicated in severe renal insufficiency where repeated measurements of the plasma concentration cannot be performed.
- Co-Trimoxazole should not be given to infants during the first 6 weeks of life.
- Co-Trimoxazole should not be given to patients with a history of drug-induced immune thrombocytopenia with use of trimethoprim and/or sulfonamides.
- Co-Trimoxazole should not be given to patients with acute porphyria.



AQOQ Pharma Ltd.

(WHO - GMP CERTIFIED - GOVT RECOGNISED EXPORT HOUSE)



CRN: 11-05918

Regd. Office & Factory : Plot No. 33, Sector II, The Vasai Taluka Industrial Co-op. Estate Ltd. Gaurapada, Vasai (E), Dist. Thane - 401 208. INDIA.
Tel. : 95250 - 2455801 / 2452714 / 2453525 • Fax : 95250 - 2452074 (0091 - 250 - 2452074) • Email : aqoq@vsnl.net & aqoqpharma@rediffmail.com

4.4 Special Warnings and Precautions for Use

Fatalities, although very rare, have occurred due to severe reactions including Stevens-Johnson syndrome, toxic epidermal necrolysis, fulminant hepatic necrosis, agranulocytosis, aplastic anaemia, other blood dyscrasias and hypersensitivity of the respiratory tract.

- Life-threatening cutaneous reactions Stevens-Johnson syndrome (SJS) and toxic epidermal necrolysis (TEN) have been reported with the use of co-trimoxazole.

- Patients should be advised of the signs and symptoms and monitored closely for skin reactions. The highest risk for occurrence of SJS or TEN is within the first weeks of treatment.

- If symptoms or signs of SJS or TEN (e.g. progressive skin rash often with blisters or mucosal lesions) are present, co-trimoxazole treatment should be discontinued.

- The best results in managing SJS and TEN come from early diagnosis and immediate discontinuation of any suspect drug. Early withdrawal is associated with a better prognosis.

- If the patient has developed SJS or TEN with the use of co-trimoxazole, co-trimoxazole must not be re-started in this patient at any time.

- At the start of treatment, the occurrence of a generalised febrile erythema associated with pustules, should raise the suspicion of acute generalised exanthematouspustulosis (AGEP) it requires cessation of treatment and contraindicates any new administration of Co-Trimoxazole alone or in combination with other drugs.

Particular care is *always* advisable when treating elderly patients because, as a group, they are more susceptible to adverse reactions and more likely to suffer serious effects as a result particularly when complicating conditions exist, e.g. impaired kidney and/or liver function and/or concomitant use of other drugs.

For patients with known renal impairment special measures should be adopted.

An adequate urinary output should be maintained at all times. Evidence of crystalluria *in vivo* is rare, although sulfonamide crystals have been noted in cooled urine from treated patients. In patients suffering from malnutrition the risk may be increased.

Regular monthly blood counts are advisable when co-trimoxazole is given for long periods, or to folate deficient patients or to the elderly; since there exists a possibility of asymptomatic changes in haematological laboratory indices due to lack of available folate. Supplementation with folic acid may be considered during treatment but this should be initiated with caution due to possible interference with antimicrobial efficacy.

In glucose-6-phosphate dehydrogenase (G-6-PD) deficient patients haemolysis may occur.

Co-trimoxazole should be given with caution to patients with severe allergy or bronchial asthma.

Co-trimoxazole should not be used in the treatment of streptococcal pharyngitis due to Group A beta-haemolytic *streptococci*; eradication of these organisms from the oropharynx is less effective than with penicillin.

Trimethoprim has been noted to impair phenylalanine metabolism but this is of no significance in phenylketonuric patients on appropriate dietary restriction.



AGOQ Pharma Ltd.

(WHO - GMP CERTIFIED - GOVT RECOGNISED EXPORT HOUSE)



Regd. Office & Factory : Plot No. 33, Sector II, The Vasai Taluka Industrial Co-op. Estate Ltd. Gauripada, Vasai (E), Dist. Thane - 401 208, INDIA.
Tel. : 95250 - 2455801 / 2452714 / 2453525 • Fax : 95250 - 2452074 (0091 - 250 - 2452074) • Email : agoq@vsnl.net & agoqpharma@rediffmail.com

The administration of co-trimoxazole to patients known or suspected to be at risk of acute porphyria should be avoided. Both trimethoprim and sulfonamides (although not specifically sulfamethoxazole) have been associated with clinical exacerbation of porphyria.

Close monitoring of serum potassium is warranted in patients at risk of hyperkalaemia and hyponatraemia.

Co-Trimoxazole has been associated with metabolic acidosis when other possible underlying causes have been excluded. Close monitoring is always advisable when metabolic acidosis is suspected.

Except under careful supervision co-trimoxazole should not be given to patients with serious haematological disorders. Co-trimoxazole has been given to patients receiving cytotoxic therapy with little or no additional effect on the bone marrow or peripheral blood.

The combination of antibiotics in co-trimoxazole should only be used where, in the judgement of the physician, the benefits of treatment outweigh any possible risks; consideration should be given to the use of a single effective antibacterial agent.

This medicine contains less than 1 m/mol sodium (23 mg) per tablet, that is to say essentially 'sodium-free'.

4.5 Interaction with Other Drugs, Other Forms of Interactions

Interaction with laboratory tests: trimethoprim may interfere with the estimation of serum/plasma creatinine when the alkaline picrate reaction is used. This may result in overestimation of serum/plasma creatinine of the order of 10%. The creatinine clearance is reduced: the renal tubular secretion of creatinine is decreased from 23% to 9% whilst the glomerular filtration remains unchanged.

Zidovudine: in some situations, concomitant treatment with zidovudine may increase the risk of haematological adverse reactions to co-trimoxazole. If concomitant treatment is necessary, consideration should be given to monitoring of haematological parameters.

Cyclosporin: reversible deterioration in renal function has been observed in patients treated with co-trimoxazole and cyclosporin following renal transplantation.

Rifampicin: concurrent use of rifampicin and Co-Trimoxazole results in a shortening of the plasma half-life of trimethoprim after a period of about one week. This is not thought to be of clinical significance.

When trimethoprim is administered simultaneously with drugs that form cations at physiological pH, and are also partly excreted by active renal secretion (e.g. procainamide, amantadine), there is the possibility of competitive inhibition of this process which may lead to an increase in plasma concentration of one or both of the drugs.

Diuretics (thiazides): in elderly patients concurrently receiving diuretics, mainly thiazides, there appears to be an increased risk of thrombocytopenia with or without purpura.

Pyrimethamine: occasional reports suggest that patients receiving pyrimethamine at doses in excess of 25 mg weekly may develop megaloblastic anaemia should co-trimoxazole be prescribed concurrently.

Warfarin: co-trimoxazole has been shown to potentiate the anticoagulant activity of warfarin via stereo-selective inhibition of its metabolism. Sulfamethoxazole may displace warfarin



from plasma-albumin protein-binding sites *in vitro*. Careful control of the anticoagulant therapy during treatment with Co-Trimoxazole is advisable.

Phenytoin: co-trimoxazole prolongs the half-life of phenytoin and if co-administered could result in excessive phenytoin effect. Close monitoring of the patient's condition and serum phenytoin levels are advisable.

Digoxin: concomitant use of trimethoprim with digoxin has been shown to increase plasma digoxin levels in a proportion of elderly patients.

Methotrexate: co-trimoxazole may increase the free plasma levels of methotrexate. If Co-Trimoxazole is considered appropriate therapy in patients receiving other anti-folate drugs such as methotrexate, a folate supplement should be considered.

Trimethoprim interferes with assays for serum methotrexate when dihydrofolate reductase from *Lactobacillus casei* is used in the assay. No interference occurs if methotrexate is measured by radioimmuno assay.

Lamivudine: administration of trimethoprim /sulfamethoxazole 160 mg/800 mg (co-trimoxazole) causes a 40% increase in lamivudine exposure because of the trimethoprim component. Lamivudine has no effect on the pharmacokinetics of trimethoprim or sulfamethoxazole.

Interaction with sulfonylurea hypoglycaemic agents is uncommon but potentiation has been reported.

Hyperkalaemia: caution should be exercised in patients taking any other drugs that can cause hyperkalaemia, for example ACE inhibitors, angiotensin receptor blockers and potassium-sparing diuretics such as spironolactone. Concomitant use of trimethoprim-sulfamethoxazole (co-trimoxazole) may result in clinically relevant hyperkalaemia.

Repaglinide: trimethoprim may increase the exposure of repaglinide which may result in hypoglycaemia.

Folic acid: folic acid supplementation has been shown to interfere with the antimicrobial efficacy of trimethoprim-sulfamethoxazole. This has been observed in *Pneumocystis jirovecii* pneumonia prophylaxis and treatment.

Contraceptives: oral contraceptive failures have been reported with antibiotics. The mechanism of this effect has not been elucidated. Women on treatment with antibiotics should temporarily use a barrier method in addition to the oral contraceptive, or choose another method of contraception.

Azathioprine: There are conflicting clinical reports of interactions between azathioprine and trimethoprim-sulfamethoxazole, resulting in serious haematological abnormalities.

4.6 Use in Pregnancy and Lactation

Pregnancy

Trimethoprim and sulfamethoxazole cross the placenta and their safety in pregnant women has not been established. Case-control studies have shown that there may be an association between exposure to folate antagonists and birth defects in humans.

Trimethoprim is a folate antagonist and, in animal studies, both agents have been shown to cause foetal abnormalities (see section 5.3).

Co-trimoxazole should not be used in pregnancy, particularly in the first trimester, unless clearly necessary. Folate supplementation should be considered if co-trimoxazole is used in pregnancy.

Sulfamethoxazole competes with bilirubin for binding to plasma albumin. As significantly maternally derived drug levels persist for several days in the newborn, there may be a risk of precipitating or exacerbating neonatal hyperbilirubinaemia, with an associated theoretical risk of kernicterus, when co-trimoxazole is administered to the mother near the time of delivery. This theoretical risk is particularly relevant in infants at increased risk of hyperbilirubinaemia, such as those who are preterm and those with glucose-6-phosphate dehydrogenase deficiency.

Breast-feeding

The components of co-trimoxazole (trimethoprim and sulfamethoxazole) are excreted in breast milk. Administration of co-trimoxazole should be avoided in late pregnancy and in lactating mothers where the mother or infant has, or is at particular risk of developing, hyperbilirubinaemia. Additionally, administration of co-trimoxazole should be avoided in infants younger than eight weeks in view of the predisposition of young infants to hyperbilirubinaemia.

4.7 Effects on ability to drive and operate machine

There have been no studies to investigate the effect of co-trimoxazole on driving performance or the ability to operate machinery. Further a detrimental effect on such activities cannot be predicted from the pharmacology of the drug. Nevertheless the clinical status of the patient and the adverse events profile of co-trimoxazole should be borne in mind when considering the patients ability to operate machinery.

4.8 Undesirable effects

The frequency categories associated with the adverse events below are estimates. For most events, suitable data for estimating incidence were not available. In addition, adverse events may vary in their incidence depending on the indication.

Data from large published clinical trials were used to determine the frequency of very common to rare adverse events. Very rare adverse events were primarily determined from post-marketing experience data and therefore refer to reporting rate rather than a "true" frequency.

The following convention has been used for the classification of adverse events in terms of frequency: Very common $\geq 1/10$, common $\geq 1/100$ and $< 1/10$, uncommon $\geq 1/1000$ and $< 1/100$, rare $\geq 1/10,000$ and $< 1/1000$, very rare $< 1/10,000$, not known - cannot be estimated from the available data.

System Organ Class	Frequency	Side effects
Infections and infestations	Common	Overgrowth fungal.
	Very rare	Pseudomembranous colitis
Blood and lymphatic system	Very rare	Leukopenia, neutropenia,



disorders		thrombocytopenia, agranulocytosis, anaemia megaloblastic, aplastic anaemia, haemolytic anaemia, methaemoglobinaemia, eosinophilia, purpura, haemolysis in certain susceptible G-6-PD deficient patients.
Immune system disorders	Very rare	Serum sickness, anaphylactic reaction, allergic myocarditis, hypersensitivity vasculitis resembling Henoch-Schoenleinpurpura, periarteritis nodosa, systemic lupus erythematosus. Severe hypersensitivity reactions associated with PJP*, rash, pyrexia, neutropenia, thrombocytopenia, hepatic enzyme increased, hyperkalaemia, hyponatraemia, rhabdomyolysis.
Metabolism and nutrition disorders	Very common	Hyperkalaemia.
	Very rare	Hypoglycaemia, hyponatraemia, decreased appetite, metabolic acidosis
Psychiatric disorders	Very rare	Depression, hallucination.
	Not known	Psychotic disorder.
Nervous system disorders	Common	Headache.
	Very rare	Meningitis aseptic *, convulsions, neuropathy peripheral, ataxia, dizziness.
Ear and labyrinth disorders	Very rare	Vertigo, tinnitus
Eye disorders	Very rare	Uveitis
Respiratory, thoracic and mediastinal disorders	Very rare	Cough *, dyspnoea*, lung infiltration*.
Gastrointestinal disorders	Common	Nausea, diarrhoea.
	Uncommon	Vomiting.
	Very rare	Glossitis, stomatitis, pancreatitis.
Hepatobiliary disorders	Very rare	Transaminases increased, blood bilirubin increased, cholestatic jaundice, hepatic necrosis
Skin and subcutaneous tissue disorders*	Common	Rash
	Very rare	Photosensitivity reaction, angioedema, dermatitis exfoliative, fixed drug eruption, erythema multiforme, Stevens-Johnson syndrome (SJS) *, toxic epidermal necrolysis (TEN) *. Acute generalised exanthematouspustulosis (AGEP).
	Not known	Acute febrile neutrophilicdermatosis (Sweet's syndrome)
Musculoskeletal and connective tissue disorders	Very rare	Arthralgia, myalgia.



AGQO Pharma Ltd.

(WHO - GMP CERTIFIED - GOVT RECOGNISED EXPORT HOUSE)



CG# 11-09818

Regd. Office & Factory : Plot No. 33, Sector II, The Vasai Taluka Industrial Co-op. Estate Ltd. Gauripada, Vasai (E), Dist. Thane - 401 208. INDIA.
Tel. : 95250 - 2455801 / 2452714 / 2453525 • Fax : 95250 - 2452074 (0091 - 250 - 2452074) • Email : agqo@vsnl.net & agqopharma@rediffmail.com

Renal and urinary disorders	Very rare	Renal impairment (sometimes reported as renal failure), tubulointerstitial nephritis and uveitis syndrome, renal tubular acidosis
-----------------------------	-----------	---

4.9 Overdoses

Symptoms

Nausea, vomiting, dizziness and confusion are likely signs/symptoms of overdosage. Bone marrow depression has been reported in acute trimethoprim overdosage.

Treatment

If vomiting has not occurred, induction of vomiting may be desirable. Gastric lavage may be useful, though absorption from the gastrointestinal tract is normally very rapid and complete within approximately two hours. This may not be the case in gross overdosage. Dependant on the status of renal function administration of fluids is recommended if urine output is low.

Both trimethoprim and active sulfamethoxazole are moderately dialysable by haemodialysis. Peritoneal dialysis is not effective.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmaco-Kinetic Properties

Absorption

After oral administration trimethoprim and sulfamethoxazole are rapidly and nearly completely absorbed. The presence of food does not appear to delay absorption. Peak levels in the blood occur between one and four hours after ingestion and the level attained is dose related. Effective levels persist in the blood for up to 24 hours after a therapeutic dose. Steady state levels in adults are reached after dosing for 2-3 days. Neither component has an appreciable effect on the concentrations achieved in the blood by the other.

Distribution

Approximately 50% of trimethoprim in the plasma is protein bound.

Tissue levels of trimethoprim are generally higher than corresponding plasma levels, the lungs and kidneys showing especially high concentrations. Trimethoprim concentrations exceed those in plasma in the case of bile, prostatic fluid and tissue, saliva, sputum and vaginal secretions. Levels in the aqueous humor, breast milk, cerebrospinal fluid, middle ear fluid, synovial fluid and tissue (intestinal) fluid are adequate for antibacterial activity. Trimethoprim passes into amniotic fluid and foetal tissues reaching concentrations approximating those of maternal serum.

Approximately 66% of sulfamethoxazole in the plasma is protein bound. The concentration of active sulfamethoxazole in amniotic fluid, aqueous humour, bile, cerebrospinal fluid, middle ear fluid, sputum, synovial fluid and tissue (interstitial) fluids is of the order of 20 to 50% of the plasma concentration.

Biotransformation

Renal excretion of intact sulfamethoxazole accounts for 15-30% of the dose. This drug is more extensively metabolised than trimethoprim, via acetylation, oxidation or glucuronidation. Over a 72 hour period, approximately 85% of the dose can be accounted for in the urine as unchanged drug plus the major (N4-acetylated) metabolite.

Elimination

The half-life of trimethoprim in man is in the range 8.6 to 17 hours in the presence of normal renal function. It is increased by a factor of 1.5 to 3.0 when the creatinine clearance is less than 10 ml/minute. There appears to be no significant difference in elderly patients compared with young patients.

The principal route of excretion of trimethoprim is renal and approximately 50% of the dose is excreted in the urine within 24 hours as unchanged drug. Several metabolites have been identified in the urine. Urinary concentrations of trimethoprim vary widely.

The half-life of sulfamethoxazole in man is approximately 9 to 11 hours in the presence of normal renal function.



AQOQ Pharma Ltd.

(WHO - GMP CERTIFIED - GOVT RECOGNISED EXPORT HOUSE)



CGN: 11-09918

Regd. Office & Factory : Plot No. 33, Sector II, The Vasai Taluka Industrial Co-op. Estate Ltd. Gauripada, Vasai (E), Dist. Thane - 401 208, INDIA.
Tel. : 95250 - 2455801 / 2452714 / 2453525 • Fax : 95250 - 2452074 (0091 - 250 - 2452074) • Email : aqoq@vsnl.net & aqoqpharma@rediffmail.com

There is no change in the half-life of active sulfamethoxazole with a reduction in renal function but there is prolongation of the half-life of the major, acetylated metabolite when the creatinine clearance is below 25 ml /minute.

The principal route of excretion of sulfamethoxazole is renal; between 15% and 30% of the dose recovered in the urine is in the active form.

The pharmacokinetics in the paediatric population with normal renal function of both components of Co-Trimoxazole, TMP and SMZ are age dependent. Elimination of TMP-SMZ is reduced in neonates, during the first two months of life, thereafter both TMP and SMZ show a higher elimination with a higher body clearance and a shorter elimination half-life. The differences are most prominent in young infants (> 1.7 months up to 24 months) and decrease with increasing age, as compared to young children (1 year up to 3.6 years), children (7.5 years and < 10 years) and adults.

Trimethoprim is a weak base with a pKa of 7.4. It is lipophilic. Tissue levels of trimethoprim are generally higher than corresponding plasma levels, the lungs and kidneys showing especially high concentrations. Trimethoprim concentrations exceed those in plasma in the case of bile, prostatic fluid and tissue, saliva, sputum and vaginal secretions. Levels in the aqueous humor, breast milk, cerebrospinal fluid, middle ear fluid, synovial fluid and tissue (intestinal) fluid are adequate for antibacterial activity. Trimethoprim passes into amniotic fluid and foetal tissues reaching concentrations approximating those of maternal serum.

In elderly patients there is a reduced renal clearance of sulfamethoxazole.

Special patient population

Renal impairment

The elimination half-life of trimethoprim is increased by a factor of 1.5-3.0 when the creatinine clearance is less than 10 mL/minute. When the creatinine clearance falls below 30 mL/min the dosage of Co-Trimoxazole should be reduced.

Hepatic impairment

Caution should be exercised when treating patients with severe hepatic parenchymal damage as there may be changes in the absorption and biotransformation of trimethoprim and sulfamethoxazole.

Elderly patients

In elderly patients, a slight reduction in renal clearance of sulfamethoxazole but not trimethoprim has been observed.

5.2 Pharmacodynamic properties

Pharmacotherapeutic group: Combinations of sulfonamides and trimethoprim, incl. derivatives;

Mechanism of Action

Co-trimoxazole is an antibacterial drug composed of two active principles, sulfamethoxazole and trimethoprim. Sulfamethoxazole is a competitive inhibitor of dihydropteroatesynthetase



AGOQ Pharma Ltd.

(WHO - GMP CERTIFIED - GOVT RECOGNISED EXPORT HOUSE)



CIN: 11-05913

Regd. Office & Factory : Plot No. 33, Sector II, The Vasai Taluka Industrial Co-op. Estate Ltd. Gauripada, Vasai (E), Dist. Thane - 401 208, INDIA.
Tel. : 95250 - 2455801 / 2452714 / 2453525 • Fax : 95250 - 2452074 (0091 - 250 - 2452074) • Email : agoq@vsnl.net & agoqpharma@rediffmail.com

enzyme. Sulfamethoxazole competitively inhibits the utilisation of para-aminobenzoic acid (PABA) in the synthesis of dihydrofolate by the bacterial cell resulting in bacteriostasis. Trimethoprim reversibly inhibits bacterial dihydrofolate reductase (DHFR), an enzyme active in the folate metabolic pathway converting dihydrofolate to tetrahydrofolate. Depending on the conditions the effect may be bactericidal. Thus trimethoprim and sulfamethoxazole block two consecutive steps in the biosynthesis of purines and therefore nucleic acids essential to many bacteria. This action produces marked potentiation of activity *in vitro* between the two agents.

Trimethoprim binds to plasmodial DHFR but less tightly than to the bacterial enzyme. Its affinity for mammalian DHFR is some 50,000 times less than for the corresponding bacterial enzyme.

Mechanism of resistance

In vitro studies have shown that bacterial resistance can develop more slowly with both sulfamethoxazole and trimethoprim in combination than with either sulfamethoxazole or trimethoprim alone.

Resistance to sulfamethoxazole may occur by different mechanisms. Bacterial mutations cause an increase in the concentration of PABA and thereby out-compete with sulfamethoxazole resulting in a reduction of the inhibitory effect on dihydropteroatesynthetase enzyme. Another resistance mechanism is plasmid-mediated and results from production of an altered dihydropteroatesynthetase enzyme, with reduced affinity for sulfamethoxazole compared to the wild-type enzyme.

Resistance to trimethoprim occurs through a plasmid-mediated mutation which results in production of an altered dihydrofolate reductase enzyme having a reduced affinity for trimethoprim compared to the wild-type enzyme.

Trimethoprim binds to plasmodial DHFR but less tightly than to bacterial enzyme. Its affinity for mammalian DHFR is some 50,000 times less than for the corresponding bacterial enzyme.

Many common pathogenic bacteria are susceptible *in vitro* to trimethoprim and sulfamethoxazole at concentrations well below those reached in blood, tissue fluids and urine after the administration of recommended doses. In common with other antibiotics, however, *in vitro* activity does not necessarily imply that clinical efficacy has been demonstrated and it must be noted that satisfactory susceptibility testing is achieved only with recommended media free from inhibitory substances, especially thymidine and thymine.

5.3 Pre-clinical safety data

Reproductive toxicology: At doses in excess of recommended human therapeutic dose, trimethoprim and sulfamethoxazole have been reported to cause cleft palate and other foetal abnormalities in rats, findings typical of a folate antagonist. Effects with trimethoprim were preventable by administration of dietary folate. In rabbits, foetal loss was seen at doses of trimethoprim in excess of human therapeutic doses



AGOG Pharma Ltd.

(WHO - GMP CERTIFIED - GOVT RECOGNISED EXPORT HOUSE)



CPN: 11-05913

Regd. Office & Factory : Plot No. 33, Sector II, The Vasai Taluka Industrial Co-op. Estate Ltd. Gauripada, Vasai (E), Dist. Thane - 401 208. INDIA.
Tel. : 95250 - 2455801 / 2452714 / 2453525 • Fax : 95250 - 2452074 (0091 - 250 - 2452074) • Email : agog@vsnl.net & agogpharma@rediffmail.com

6. PHARMACEUTICAL PARTICULARS

6.1 List of Excipients

Methyl Paraben Sodium	BP	7.50 mg
Propyl Paraben Sodium	BP	2.50 mg
Carboxy methyl Cellulose Sodium	BP	40.0 mg
Sodium Saccharin	BP	2.50 mg
Citric acid monohydrate	BP	5.00 mg
Sodium Benzoate	BP	7.50 mg
Colour sunset yellow supra	Inhouse	0.16 mg
Essence Orange Sweet No.1	Inhouse	0.01 ml
Sucrose	BP	1500 mg
Purified Water	BP	Q.S. to 2000 ml

6.2 Incompatibilities

None reported

6.3 Shelf-Life

36 months from the date of manufacture.

6.4 Special Precautions for Storage

Store below 30°C.
Protect from light.

6.5 Nature and Contents of Container

Bottle pack of 60 ml & 100 ml



AGOG Pharma Ltd.

(WHO - GMP CERTIFIED - GOVT RECOGNISED EXPORT HOUSE)



CRN: 11-09918

Regd. Office & Factory : Plot No. 33, Sector II, The Vasai Taluka Industrial Co-op. Estate Ltd. Gauripada, Vasai (E), Dist. Thane - 401 208. INDIA.
Tel. : 95250 - 2455801 / 2452714 / 2453525 • Fax : 95250 - 2452074 (0091 - 250 - 2452074) • Email : agog@vsnl.net & agogpharma@rediffmail.com

**ANIL K. PANDEY
DIRECTOR**

**Date :
Director of the manufacturer
(Signature, Full name, Stamp)**



**ANIL K. PANDEY
DIRECTOR**

**Date :
Director of applicant company
(Signature, Full name, Stamp)**

