



MEPACO
MEDIFOOD



Factories:
Enshas El Raml, Sharkeiya
Tel : +2 055 2820406
+2 055 2822711
Fax : +2 055 2822690
Mob: +2 0122 7693336
+2 0122 7693337
Abo Sultan - Fayed
Tel : +2 064 - 3400360
Fax: +2 064 - 3400876

Scientific office:
Secondary Trade Building,
Entrance No. 5, Fifth Area
Sheraton Heliopolis, Cairo
Zip code 11361 Heliopolis
Tel : +2 02 22673175
+ 2 02 22673176
Mob : +2 0122 7693377
+2 0122 7693388
Fax : +2 02 22674934

Alexandria Office:
21, Mostafa Kamel Towers
Flat: 33
Tel : +2 03 54 57 140
Mob: +2 0122 7693399
Fax : +2 03 54 57 140

الصناع:
أنصاص الرمل - الشرقية
تلفون: +2 055 282 401
+2 055 282 2711
فاكس: +2 055 282 2719
 محمول: +2 0122 769 3336
+2 0122 769 3337
أبو سلطان - فايد
تلفون: +2 064 340 360
+2 064 340 871
فاكس: +2 064 340 871

الكتب العلمي:
مدخل ٥ - المبني التجاري الثاني
المنطقة الخامسة، مساكن
الشيراتون رقم بريدي ١١٣٦١
هليوبوليس - الجريدة
تلفون: +2 02 227 2175
+2 02 227 2171
محفظ: +2 0122 769 3377
+2 0122 769 3388
فاكس: +2 02 227 24734
مكتب الأسكندرية:

عمرات ضباط مصرفي كامل
عمارة ٢١ شقة ٣٣
تلفون: +2 03 571 40
+2 0122 769 3399
محفظ: +2 03 571 40
فاكس: +2 03 571 40

Arab Company for Pharmaceuticals. & Medicinal Plants S.A.E MEPACO - MEDIFOOD

الشركة العربية للأدوية والنباتات الطبية ش.م.م ميباكو - ميدي فود

Summary of Product Characteristics

1. Name Of The Medicinal Product:

Vitamin E Forte Soft Gelatin Capsules.

2. Qualitative And Quantitative Composition

Each Capsule Contains:

dl-Alpha Tocopherol Acetate Equivalent to 400 I.U Natural Vitamin E .	400 mg	BP 2018
---	--------	---------

Composition of soft gel. Capsule (Oval 6):

Gelatin bloom (150 -160)	144 mg	BP 2018
Glycerol	40.30 mg	BP 2018
Sorbitol solution 70%	27.84 mg	BP 2018
Propyl paraben	0.072 mg	BP 2018
Methyl paraben	0.144 mg	BP 2018

3. Pharmaceutical Form:

Capsules for oral administration

4. Clinical Particulars:

4.1 Therapeutic Indications:

Vitamin E capsules is indicated for

- Hyperlipoproteinemas, atherosclerosis and cardiomyopathy.
- Haemolytic anemia.
- Male infertility, habitual abortion and post IUD insertion.
- Epilepsy, Parkinson's disease and larvide dyskinesia.
- Cholestatic liver disease.

4.2 Posology And Method Of Administration:

One capsule daily.

4.3 Contraindications:

Vitamin E is contraindicated in the following conditions:

- Retinitis pigmentosa (RP): Investigators hypothesised that the 400 IU day-1 dosage of vitamin E may inhibit the absorption or transport of vitamin A, as the participants

Factories:

Enshas El Raml, Sharkeiya

Tel : +2 055 2820406

+2 055 2822711

Fax : +2 055 2822690

Mob: +2 0122 7693336

+2 0122 7693337

Abo Sultan - Fayed

Tel : +2 064 - 3400360

Fax: +2 064 - 3400876

Scientific office:

Secondary Trade Building,

Entrance No. 5, Fifth Area

Sheraton Heliopolis, Cairo

Zip code 11361 Heliopolis

Tel : +2 02 22673175

+2 02 22673176

Mob : +2 0122 7693377

+2 0122 7693388

Fax : +2 02 22674934

Alexandria Office:

21, Mostafa Kamel Towers

Flat: 33

Tel : +2 03 54 57 140

Mob: +2 0122 7693399

Fax : +2 03 54 57 140

الصناع:

أنصاص الرمل - الشرقية

٤٠٥٥٢٨٢٤٠١

+٢٠٥٥٦٨٢٧٦١١

فاكس: ٤٠٥٥٢٨٢٣٩٠

+٢٠١٢٢٧٦٩٣٣٦

موبايل: ٠١٢٢٧٦٩٣٣٧

+٢٠١٢٢٧٦٩٣٣٧

أبو سلطان - فايد

٤٠٦٤٣٤٠٣٦٠

تلفون: ٥٠١٢٤٣٤٠٨٧٦

فاكس: ٥٠١٢٤٣٤٠٨٧٦

المكتب العلمي:

مدخل ٥ - المبني التجاري الثاني

المنطقة الخامسة، مساكن

الشبريون رقم بريدي ١١٣٦١

هليوبوليس - الخيرية

تلفون: ٤٠٢٢٧٣١٧٥

+٢٠٢٢٧٣١٧٦

موبايل: ٠١٢٢٧٦٩٣٧٧

+٢٠١٢٢٧٦٩٣٨٨

فاكس: ٤٠٢٢٧٤٩٣٤

مكتب الأسكندرية:

مطارات ضباط مصطفى كامل

عمارة ٢١ شقة

٣٣ تليفون: ٤٠٣٥٤٥٧١٤٠

+٢٠١٢٢٧٦٩٣٩٩

موبايل: ٠١٢٢٧٦٩٣٤٠

+٢٠٣٥٤٥٧١٤٠

receiving this dosage of vitamin E had significantly decreased serum retinol levels compared with those not receiving vitamin E (**Grover and Samson , 2014;**

Pang et al., 2001)

- Vitamin K deficiency: Vitamin E may exacerbate the effects of Vitamin K deficiency and has been associated with an increased risk of haemorrhagic stroke, and a reduced risk of ischaemic stroke and ischaemic heart disease.

- Oral anticoagulants administration: The action of oral anticoagulants may be increased by concurrent administration of vitamin E. Concurrent vitamin E and warfarin use has been associated with abnormal bleeding, and vitamin E appears to add to aspirin's blood thinning effect. (**Bartlett, H., et al., 2005).**

- Angioplasty, a heart procedure: vitamin E or other antioxidant vitamins (beta-carotene, vitamin C) should be avoided immediately before and following angioplasty . This may be due to interference with proper healing.

• 4.4 Special Warnings And Precautions For Use:

• Precautions:

Vitamin E is a natural blood thinner , so it may increase the risk of bleeding, especially if you are taking blood-thinners such as warfarin ,aspirin, or clopidogrel .
 Vitamin E can cause a temporary elevation of blood pressure.

Vitamin E in 400 IU per day or higher; may increase the risk of death.

Vitamin E should not be taken in higher amounts during pregnancy or breastfeeding.
 Oral administration of large doses (200 mg per day) of ahyperosmolar vitamin E preparation to low-birth weight infants hasbeen associated with the development of necrotizing enterocolitis

• 4.5 Interaction With Other Medicinal Products And Other Forms Of Interaction:

Vitamin E may cause the following interactions between the following medications:

Antidepressant medications: Vitamin E interferes with the absorption of the antidepressant desipramine (**Scuntaro et al., 1996).**
 Antipsychotic medications: Vitamin E can interfere with the body's absorption of the antipsychotic medication such as chlorpromazine (**Soares and McGrath, 2001).**

Aspirin: Because vitamin E can increase the risk of bleeding (**Podszun and Frank, 2014).**

AZT: Vitamin E may protect against toxicity and side effects from AZT, a medication used to treat HIV and AIDS (**García-de-la-Asunción et al., 2007).**

Beta-blockers: Vitamin E interferes with the body's absorption of propranolol and Other beta-blockers include: Acebutolol, Atenolol,Bisoprolol, Carvedolol, Metoprolol, Nadolol, and Propranolol.

Cholesterol-lowering medications:

Factories:

Enshas El Raml, Sharkeiya
Tel : +2 055 2820406
+2 055 2822711
Fax : +2 055 2822690
Mob: +2 0122 7693336
+2 0122 7693337

Abo Sultan - Fayed

Tel : +2 064 - 3400360
Fax: +2 064 - 3400876

Scientific office:

Secondary Trade Building,
Entrance No. 5, Fifth Area
Sheraton Heliopolis, Cairo
Zip code 11361 Heliopolis
Tel : +2 02 22673175
+2 02 22673176
Mob : +2 0122 7693377
+2 0122 7693388
Fax : +2 02 22674934

Alexandria Office:

21, Mostafa Kamel Towers
Flat: 33
Tel : +2 03 54 57 140
Mob: +2 0122 7693399
Fax : +2 03 54 57 140

الصانع:
أنساخ الرمل - الشرقية
تلفون: +٢٠٠٥٤٢٨٠٤١
+٢٠٠٥٤٢٨٣٧١١
فاكس: +٢٠٠٥٤٨٢٢١٩٠
محمول: +٢٠١٢٢٧٩٣٣٦٩
+٢٠١٢٢٧٩٣٣٧

أبو سلطان - فايد
تلفون: +٢٠٠٤٣٤٠٣٦٠
+٢٠١٤٣٤٠٨٧١

المكتب العلمي:
مدخل ٥ - المبني التجاري الثاني
المنطقة الخامسة، مساكن
الشيراتون رقم بريدي ١١٣٦١
هليوبوليس - الخيرية
تلفون: +٢٠٢٢٧٣١٧٥
+٢٠٢٢٧٣٧٦١
محمول: +٢٠١٢٢٧٩٣٧٧
+٢٠١٢٢٧٩٣٨٨
فاكس : +٢٠٢٢٧٩٤٩٣٤

مكتب الأسكندرية:
عمارات ضباط مصرفي كامل
عمارة ٢١ شقة ٣٣
تلفون: +٢٠٣٥٤٥٧١٤٠
+٢٠١٢٢٧٩٣٣٩٩
محمول: +٢٠١٢٢٧٩٣٣٩٩
فاكس : +٢٠٣٥٤٥٧١٤٠

- **Bile acid sequestrants:** may reduce how much vitamin E your body absorbs (**Sutken et al., 2006**).
- **Statins:** Taking a combination of antioxidants, including vitamins E and C, selenium, and beta-carotene, along with niacin and simvastatin (Zocor) seems to keep niacin and Zocor from raising “good” HDL cholesterol (**Sutken et al., 2006**).
- **Gemfibrozil:** may also reduce vitamin E levels (**Sutken et al., 2006**).
- Cyclosporine:** Vitamin E may interact with cyclosporine, a medication used to treat cancer, making both vitamin E and cyclosporine less effective (**Chang et al., 1996**).
- Chemotherapy drugs:** vitamin E may reduce the effectiveness of chemotherapy drugs, including chlorambucil, busulfan, doxorubicine, carmustine, thiopeta, and cyclophosphamideradicals.
- Tamoxifen:** Vitamin E when taken along with tamoxifen, reduced its side effect. made the anti-cancer action of the tamoxifen stronger. (**Podszun and Frank, 2014**)
- Warfarin :** Taking vitamin E at the same time as other blood-thinners can increase the risk of bleeding (**Podszun and Frank, 2014**).
- Orlistat :** It may prevent the absorption of vitamin E (**McDuffie et al., 2002**).

4.6 Pregnancy And Lactation:

-Vitamin E is likely safe when taken by mouth in recommended daily amounts during breast-feeding.

-When used in the recommended daily amount, vitamin E is possibly safe for pregnant women. There has been some concern that taking vitamin E supplements might be harmful to the fetus when taken in early pregnancy (Boskovic et al, 2005).

4.7 Effects On Ability To Drive And Use Machines:

Not applicable

4.8 Undesirable Effects:

1. Adverse effect:

Greater dose more than 300 Units a day and long-term use may results in the following side effects:

1.1. Gastrointestinal

Gastrointestinal side effects associated with oral doses greater than 300 units daily have rarely included nausea, diarrhea, and intestinal cramps.

1.2. Hematologic

Hematologic side effects associated with oral doses greater than 300units daily have rarely included thrombophlebitis.

1.3. Ocular

Ocular side effects associated with oral doses greater than 300 units daily have rarely included blurred vision.

1.4. Dermatologic

Dermatologic side effects associated with oral doses greater than 300units daily have rarely included rash. Dermatologic side effects associated with topical application of vitamin E have included growth of white hair at site of alopecia and contact dermatitis.

Factories:

Enshas El Raml, Sharkeiya
Tel : +2 055 2820406
+2 055 2822711
Fax : +2 055 2822690
Mob: +2 0122 7693336
+2 0122 7693337

Abo Sultan - Fayed

Tel : +2 064 - 3400360
Fax: +2 064 - 3400876

Scientific office:

Secondary Trade Building,
Entrance No. 5, Fifth Area
Sheraton Heliopolis, Cairo
Zip code 11361 Heliopolis
Tel : +2 02 22673175
+2 02 22673176
Mob : +2 0122 7693377
+2 0122 7693388
Fax : +2 02 22674934

Alexandria Office:

21, Mostafa Kamel Towers
Flat : 33
Tel : +2 03 54 57 140
Mob: +2 0122 7693399
Fax : +2 03 54 57 140

الصانع:
أنصاف الرمل - الشرقية
تلفون: ٢٠٣٥٤٢٨٠٤١٠
٢٠٣٥٤٨٢٧٦١١
فاكس: ٢٠٣٥٤٨٢٣٩٠
موبايل: ٢٠١٢٢٧٦٩٣٣٣٦
٢٠١٢٢٧٦٩٣٣٧

أبو سلطان - فايد
تلفون: ٢٠٣٤٣٤٠٣٠٠
٢٠٣٤٣٤٠٨٧٦
فاكس: ٢٠٣٤٣٤٠٨٧٦

الكتب العلمي:
مدخل ٥ - المبنى التجاري الثانى
المنطقة الخامسة، مساكن
الشيرانى رقم بريدى ١١٣٦١
هليوبوليس - الخيرية
تلفون: ٢٠٢٢٧٣١٧٥
٢٠٢٢٧٣١٧٦
موبايل: ٢٠١٢٢٧٦٩٣٧٧
٢٠١٢٢٧٦٩٣٨٨
فاكس : ٢٠٢٢٧٤٩٣٤

مكتب الأسكندرية:
عقارات ضباط مصطفى كامل
عمارة ٢١ شقة ٣٣
تلفون: ٢٠٣٥٤٥٧١٤٠
٢٠١٢٢٧٦٩٣٩٩
موبايل: ٢٠٣٥٤٥٧١٤٠
فاكس : ٢٠٣٥٤٥٧١٤٠

1.5. Genitourinary

Genitourinary side effects associated with oral doses greater than 300 units daily have rarely included gonadal dysfunction and breast soreness.

Endocrine side effects associated with oral doses greater than 300 units daily have rarely included decrease in serum thyroxine and triiodothyronine.

1.7. Metabolic

Metabolic side effects associated with oral doses greater than 300 units daily have rarely included creatinuria, increased serum creatinine kinase, increased serum cholesterol and triglycerides and increased urinary estrogens and androgens.

1.8. Other

Other side effects may include fatigue, emotional disturbances, weakness, and headache. (Meyers et al., 1996)

(<http://www.drugs.com/sfx/nutr-e-sol-side-effects.html>).

• 4.9 Overdose:

Occurrence of vitamin E overdose is rare, However high doses of alphatocopherol supplements can cause hemorrhage and interrupt blood coagulation in animals, and in vitro data suggest that high doses inhibit platelet aggregation (Sesso et al., 2008).

5. Pharmacological Properties:

Vitamin E Forte Soft Gelatin is regarded as the most recommended concentrated preparation of natural alpha tocopherol, which is recognized the most active form of tocopherol.

Alpha tocopherol is considered the major lipid soluble antioxidant, protecting lipids in body tissues from peroxidative damage. In particular, it protects polyunsaturated fatty acids (PUFA) within biological membrane phospholipids or in plasma low density lipoproteins.

The medical trials and epidemiological studies have provided evidence that high blood levels of alpha tocopherol as anti-oxidant may thwart the incidence of atherosclerosis, congestive cardiomyopathy, Cholestatic liver disease, carcinogenesis and neurological problems associated with abetalipoproteinemia.

Also, the administration of high levels of high levels of alpha tocopherol has important role in the treatment of Haemolytic anemia.

Alpha tocopherol, also influence hormonal processes and fertility. In addition, the anti-inflammatory effect of alpha tocopherol is useful for habitual abortion and post IUD insertion



Arab Company for Pharmaceuticals. & Medicinal Plants S.A.E MEPACO - MEDIFOOD

الشركة العربية للأدوية والنباتات الطبية ش.م.م ميباكو - ميدي فود

6. Pharmaceutical Particulars:

- 6.1 List Of Excipients :**

No excipient added.

- 6.2 Incompatibilities:**

- All soft gelatin capsule Components comply with their BP or USP Pharmacopoeia monograph.
- None of the excipients contains materials of animal or human origin.
- No genetically modified organisms (GMO) have been used in the preparation of these products.
- No incompatibilities with the excipients or packaging materials have been shown during research of pre-formulation and formulation studies and nor during manufacturing processing and stability.

- 6.3 Shelf Life**

3 years

- 6.4 Special Precautions For Storage:**

No special precaution for storage

Store in a dry place at temperature not exceeding 30° C.

- 6.5 Nature And Contents Of Container:**

Carton box containing 2 (Aluminum- PVC) strips each of 10 capsules and inner leaflet.

- 6.6 Special Precautions For Disposal And Other Handling:**

Keep out of reach of children

- 7. Marketing Authorization Holder:**

Arab Company for Pharmaceuticals & Medicinal Plants (Mepaco – Medi food) Egypt.

- 8. Marketing Authorization Number(S)**

Reg. No: 1742/1997

Reg. No: 24576/2006

9. Date Of First Authorization/Renewal Of The Authorization:

Reg. No: 1742/1997

Authorization date: 05/01/1997

Reg. No: 24576/2006

Authorization date: 13/06/2006

R&D Manager

Dr. / Mohamed Eldegwy



www.mepaco-pharma.net

Operating Under Law 43 Issued 1974 - Commercial Register No. 866 - Capital 400 EGP Million

Page 5 of 5

خاضعة لـحكام قانون ٤٣ لسنة ١٩٧٤ وتعديلاته سجل رقم ٨٦٦ إستثمار رأس المال المرخص به ٤٠٠ مليون جنيه مصرى



Factories:

Enshas El Raml, Sharkeiya

Tel : +2 055 2820406

+2 055 2822711

Fax : +2 055 2822690

Mob: +2 0122 7693336

+2 0122 7693337

Abo Sultan - Fayed

Tel : +2 064 - 3400360

Fax: +2 064 - 3400876

Scientific office:

Secondary Trade Building,

Entrance No. 5, Fifth Area

Sheraton Heliopolis, Cairo

Zip code 11361 Heliopolis

Tel : +2 02 22673175

+2 02 22673176

Mob : +2 0122 7693377

+2 0122 7693388

Fax : +2 02 22674934

Alexandria Office:

21, Mostafa Kamel Towers

Flat: 33

Tel : +2 03 54 57 140

Mob: +2 0122 7693399

Fax : +2 03 54 57 140

المصانع:

أنصاص الرمل - الشرقية

تلفون: +2 0552820406

+2 0552822711

فاكس: +2 0552822711

موبايل: +2 01227693377

+2 01227693388

فاكس: +2 01227693377

أبو سلطان - فايد

تلفون: +2 0643400300

+2 0643400876

فاكس: +2 0643400876

الكتب العلمي:

مدخل ٥ - البني التجاري الثاني

المنطقة الخامسة، مساكن

الشبراتون رقم بريدي ١١٣٦١

هليوبوليس - الغربية

تلفون: +2 0122731750

+2 0122731761

موبايل: +2 01227693377

+2 01227693388

فاكس: +2 01227693377

مكتب الأسكندرية:

عمارات ضباط مصطفى كامل

عمارة ٢١ شقة ٣٣

تلفون: +2 035457140

+2 01227693399

فاكس: +2 035457140