

AGOG Pharma Ltd.



(WHO - GMP CERTIFIED - GOVT RECOGNISED EXPORT HOUSE)

Regd. Office & Factory: Plot No. 33, Sector II, The Vasai Taluka Industrial Co-op. Estate Ltd. Gauraipada, Vasai (E), Dist. Thane - 401 208. INDIA. Tel.: 95250 - 2452801 / 2452714 / 2453525 • Fax: 95250 - 2452074 (0091 - 250 - 2452074) • Email: agog@vsnl.net & agogpharma@rediffmail.com

1.5.3 Patient information leaflet (PIL)



Composition:

Bisacodyl is a popular non-prescription type of stimulant laxative. The drug reduces short term constipation and is also used for preparing the colon or rectum for an examination or surgical procedure.

Bisacodyl works by stimulating colon movement constipation is generally relieved within 15 minutes to 60 minutes after administration of a suppository form and in six to twelve hours after oral administration.

Dosage: The drug is taken at night if administered orally and during morning if taken through the rectum.

Adult: Constipation 1-2 tabs at night or 1×10 mg suppose in the morning. Before surgery or radiological procedure 2-4 tabs night before then 1×10 mg suppose the following morning.

Overdosage:

Signs of overdose may include diarrhoea and abdominal cramps. Diarrhoea may lead to excessive loss of water and salts. Seek advice from your doctor, pharmacist.

Side effects: Bisacodyl has certain side effects that can affect individuals in different ways. The following are some of the side effects, that are often associated with the drug. Nausea, Muscle Weakness, diarrhoea and cramps. Nursing and pregnant women should consult a doctor before taking the drug.

Storage: Store under normal storage condition (15°C to 30°C). Protect from light.

Keep all medicines out of reach of children.

Presentation: Jar Pack of 1000 Tablets. Blister pack of 10 × 10 Tablets.



Manufactured in India by:

AGOG PHARMA LTD.

Plot No. 33, Sector II, The Vasai Taluka Indl. Co-op. Estate Ltd., Vasai (East), Dist. Thane. INDIA.