

Patient information leaflet

Terbutaline Sulphate, Bromhexine Hydrochloride & Guaifenesin Syrup

TusQ® -X LIQUID

Terbutaline Sulphate, Bromhexine Hydrochloride & Guaifenesin Syrup

Read all of this leaflet carefully before you start taking this medicine.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, health care provider or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If any of these side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor, health care provider or pharmacist.

In this leaflet:

- a) What TusQ-X Liquid is and what it is used for
- b) Before you take TusQ-X Liquid
- c) How to take TusQ-X Liquid
- d) Possible side effects
- e) How to store TusQ-X Liquid
- f) Further information

a) WHAT TUSQ-X LIQUID IS AND WHAT IT IS USED FOR

TusQ-X Liquid contains Terbutaline - a medicine that relaxes the muscles in the air passage of the lungs, Bromhexine-which clears mucus from the airways, Guaiphenesin-that thins and loosens the mucus from the airways.

It is used for symptomatic treatment of productive cough associated with respiratory disorders.

b) BEFORE YOU TAKE TUSQ-X LIQUID

Do not take TusQ-X Liquid:

- If you are allergic to Bromhexine, Terbutaline, Guaiphenesin or any of the other ingredients of this medicine.
- If you have any heart disease

- If you have high levels of thyroid hormone in your body.

Take special care with TusQ-X Liquid:

- If you have severe heart disease and if you experience chest pain or other symptoms of worsening heart disease.
- If you have a history of gastric ulcer
- If you have kidney or liver problem.

Taking other medicines

Please tell your doctor, healthcare care provider or pharmacist if you are taking or have recently taken any other medicines, including medicines obtained without a prescription.

Taking TusQ-X Liquid with food and drink

- TusQ-X Liquid should be taken with or after food, or as prescribed.
- Avoid drinking alcohol

Pregnancy and breast-feeding

If you are pregnant or breast-feeding, ask your doctor, healthcare care provider or pharmacist for advice before taking this medicine.

Driving and using machines

Terbutaline or Bromhexine or Guaiphenesin has no or negligible influence on the ability to drive and use machines. However, caution should be exercised if patients experience dizziness. If affected by dizziness, patients should avoid tasks such as driving a vehicle or operating machinery.

c) HOW TO TAKE TUSQ-X LIQUID

Always take TusQ-X Liquid exactly as your doctor or health care provider has told you.

The usual dose is;

- In children above 12 years: 10 ml to 3 times daily
- In children
 - 6 to 12 years: 5 ml to 3 times daily
 - 2 to 6 years: 2.5 ml 3 times daily

Use in children below 2 years of age

TUSQ-X+ Liquid should not recommended be used in children below 2 years of age.

If you take more TusQ-X Liquid then you should

Symptoms of overdose include headache, anxiety, tremor, palpitations, increased heart rate, irregular heartbeat, abdominal (stomach) discomfort, nausea, vomiting, and drowsiness.

1. Tell your doctor, health care provider, pharmacist or nearest hospital casualty department immediately.
2. Take the bottle and any remaining suspension with you.

If you forget to take TusQ-X Liquid

If you forget to take a dose take it as soon as you remember, but if it is almost time for your next dose, skip the missed dose and continue as usual.

Do not take a double dose to make up for a forgotten dose.

If you stop taking Mefal-Spas tablets

Do not stop taking medicines on your own. Always consult your doctor, health care provider or pharmacist before stopping

d) POSSIBLE SIDE EFFECTS

Like all medicines, TusQ-X Liquid can cause side effects, although not everybody gets them.

If any of the side effects gets serious, please tell your doctor, health care provider or pharmacist;

Seek medical help if you have tremor, headache, increased heart rate, palpitations, muscle cramps, nervousness, dizziness, anxiety, reduced sleep, nausea, dry mouth and sweating.

e) HOW TO STORE TUSQ-X LIQUID

- Keep out of the reach and sight of children.
- Store at a temperature not exceeding 30⁰ C.
- Protect from light.
- Do not use TusQ-X Liquid after the expiry date which is stated on the blister carton/bottle/container after 'Expiry Date'. The expiry date refers to the last day of that month.
- Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

f) FURTHER INFORMATION

What TusQ-X Liquid contains

The active substances are Terbutaline Sulphate, Bromhexine Hydrochloride and Guaifenesin.

The other ingredient(s) are Sodium Benzoate, Xanthan Gum, Purified Water, Glycerol, Propylene Glycol, Sucralose, Sodium Citrate, Citric Acid Monohydrate, Mixed Fruit Flavour, Liquorice Flavour, Colour Tartrazine, Colour Brilliant Blue and Levomenthol.

What TusQ-X Liquid look like and contents of the bottle

TusQ-X Liquid is a green coloured clear liquid with pleasant flavour.

It is available in a 100 ml Pet bottle with measuring cup.

Name and full address of Marketing Authorization Holder and Manufacturing site:

Marketing Authorization Holder

Blue Cross Laboratories Pvt Ltd.

Peninsula Chambers, Ganpatrao Kadam Marg,

Lower Parel, Mumbai - 400013, Maharashtra, India.

Manufacturing site:

Blue Cross Laboratories Pvt Ltd.

A-12, MIDC, Ambad, Nashik – 422010, Maharashtra, India.

This leaflet was prepared on May 2024.