



COMMON TECHNICAL DOCUMENT

PRODUCT: DIGOXIN INJECTION BP 0.25 mg/ml

1.8	Product Information
1.8.1	Summary of Product Characteristics

SUMMARY OF PRODUCT CHARACTERISTICS**1. NAME OF THE PRODUCT:**

Digoxin Injection BP

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each ml contains

Digoxin BP 0.25 mg

Ethanol BP 0.125 ml

3. PHARMACEUTICAL FORM:

Solution for Injection

4. CLINICAL PARTICULARS**4.1. THERAPEUTIC INDICATIONS:****Cardiac failure**

Digoxin is indicated in the management of chronic cardiac failure where the dominant problem is systolic dysfunction. Its therapeutic benefit is greatest in those patients with ventricular dilatation.

Digoxin is specifically indicated where cardiac failure is accompanied by atrial fibrillation.

Supraventricular arrhythmias

Digoxin is indicated in the management of certain supraventricular arrhythmias, particularly chronic atrial flutter and fibrillation.

4.2 POSOLOGY AND METHOD OF ADMINISTRATION:**Posology:**

The dose of digoxin for each patient has to be tailored individually according to age, lean body weight and renal function.

Suggested doses are intended only as an initial guide.

In cases where cardiac glycosides have been taken in the preceding two weeks the recommendations for initial dosing of a patient should be reconsidered and a reduced dose is advised.

The difference in bioavailability between injectable digoxin and oral formulations must be



COMMON TECHNICAL DOCUMENT

PRODUCT: DIGOXIN INJECTION BP 0.25 mg/ml

considered when changing from one dosage form to another. For example if patients are switched from oral to the I.V. formulation the dosage should be reduced by approximately 33%.

Adults and pediatric populations over 10 years

Parenteral loading:

Parenteral loading should only be used in patients who have not been given cardiac glycosides within the preceding two weeks.

The total loading dose of parenteral digoxin is 500 to 1000 micrograms (0.5 to 1.0 mg) depending on age, lean body weight and renal function. The total loading dose should be administered in divided doses with approximately half of the total dose given as the first dose and further fractions of the total dose given at intervals of four to eight hours. An assessment of clinical response should be performed before giving each additional dose.

Each dose should be given by I.V. infusion (see Method of Administration, below) over 10 to 20 mins.

Maintenance dose:

The maintenance dosage should be based upon the percentage of the peak body stores lost each day through elimination. The following formula has had wide clinical use:

$$\text{Maintenance dose} = \text{Peak body stores} \times \frac{\text{daily loss in percent}}{100}$$

$$\text{Where: Peak body stores} = \text{personalised loading dose} \\ \text{daily loss (in percent)} = 14 + \text{creatinine clearance (C}_{cr}\text{)}/5$$

C_{cr} is creatinine clearance corrected to 70 kg bodyweight or 1.73 m² body surface area. If only serum creatinine (S_{cr}) concentrations are available, a C_{cr} (corrected to 70 kg bodyweight) may be estimated in men as

$$C_{cr} = \frac{(140 - \text{age})}{S_{cr} \text{ (in mg/100 ml)}}$$

NOTE: Where serum creatinine values are obtained in micromol/l, these may be converted to mg/100 ml (mg %) as follows:

$$S_{cr} \text{ (mg/100 ml)} = \frac{S_{cr} \text{ (micromol/l)} \times 113.12}{10,000} \\ = \frac{S_{cr} \text{ (micromol/l)}}{88.4}$$

Where 113.12 is the molecular weight of creatinine.

**COMMON TECHNICAL DOCUMENT****PRODUCT: DIGOXIN INJECTION BP 0.25 mg/ml**

For women, this result should be multiplied by 0.85.

N.B. These formulae cannot be used for creatinine clearance in children.

In practice, this will mean that most patients with heart failure will be maintained on 125 to 250 micrograms (0.125 to 0.25 mg) digoxin daily; however in those who show increased sensitivity to the adverse effects of digoxin, a dose of 62.5 micrograms (0.0625 mg) daily or less may suffice. Conversely, some patients may require a higher dose.

Neonates, infants and paediatric populations up to 10 years of age

If cardiac glycosides have been given in the two weeks preceding commencement of digoxin therapy, it should be anticipated that optimum loading doses of digoxin will be less than those recommended below.

In the newborn, particularly in the premature infant, renal clearance of digoxin is diminished and suitable dose reductions must be observed, over and above general dosage instructions.

Beyond the immediate newborn period, children generally require proportionally larger doses than adults on the basis of body weight or body surface area, as indicated in the schedule below. Children over ten years of age require adult dosages in proportion to their body weight.

Parenteral loading dose:

The I.V. loading dose in the above groups should be administered in accordance with the following schedule:

Preterm neonates less than 1.5 kg	-	20 micrograms/kg over 24 h.
Preterm neonates 1.5 kg to 2.5 kg	-	30 micrograms/kg over 24 h.
Term neonates to 2 years	-	35 micrograms/kg over 24 h.
2 to 5 years	-	35 micrograms/kg over 24 h.
5 to 10 years	-	25 micrograms/kg over 24 h.

The loading dose should be administered in divided doses with approximately half the total dose given as the first dose and further fractions of the total dose given at intervals of four to eight hours, assessing clinical response before giving each additional dose. Each dose should be given by I.V. infusion (see Method of Administration, below) over 10 to 20 mins.

Maintenance dose:

The maintenance dose should be administered in accordance with the following schedule:

Preterm neonates:

Daily dose = 20 % of 24 h loading dose.

Term neonates and children up to 10 years:

**COMMON TECHNICAL DOCUMENT****PRODUCT: DIGOXIN INJECTION BP 0.25 mg/ml**

Daily dose = 25 % of 24 h loading dose.

These dosage schedules are meant as guidelines and careful clinical observation and monitoring of serum digoxin levels (see Section 4.4) should be used as a basis for adjustment of dosage in these paediatric patient groups.

Elderly

The possibility of reduced renal function and lower lean body mass should be taken into account when dealing with elderly patients. If necessary, the dosage should be reduced and adjusted to the changed pharmacokinetics to prevent elevated serum digoxin levels and the risk of toxicity. The serum digoxin levels should be checked regularly and hypokalaemia should be avoided.

Renal impairment

The dosing recommendations should be reconsidered if patients are elderly or there are other reasons for the renal clearance of digoxin being reduced. A reduction in both initial and maintenance doses should be considered (see Section 4.4).

Method of administration

Dilution of digoxin injection:

Digoxin injection can be administered undiluted or diluted with a 4-fold or greater volume of 0.9% Sodium Chloride Injection, 0.18 % Sodium Chloride/4% Glucose Injection or 5% Glucose Injection. A 4-fold volume of diluent equates to adding one 2 ml ampoule of digoxin to 6 ml of injection solution. The use of less than a 4-fold volume of diluent could lead to precipitation of digoxin.

Digoxin injection, when diluted in the ratio of 1 to 250 is known to be compatible with the following infusion solutions and stable for up to 48 h at room temperature (20 to 25 °C):

- Sodium Chloride I.V. Infusion, B.P., 0.9 % w/v.
- Sodium Chloride (0.18 % w/v) and Glucose (4 % w/v) I.V. Infusion, B.P.
- Glucose I.V. Infusion, B.P., 5 % w/v.

A ratio of 1 to 250 can be obtained for example by diluting one 2 ml ampoule with 500 ml of infusion solution.

Dilution should be carried out either under full aseptic conditions or immediately before use. Any unused solution should be discarded (see Section 6.6).

Administration of digoxin injection:

Each dose should be given by I.V. infusion over 10 to 20 mins.

The total loading dose should be administered in divided doses with approximately half of the total

**COMMON TECHNICAL DOCUMENT****PRODUCT: DIGOXIN INJECTION BP 0.25 mg/ml**

dose given as the first dose and further fractions of the total dose given at intervals of four to eight hours. An assessment of clinical response should be performed before giving each additional dose.

The I.M. route is painful and is associated with muscle necrosis. This route cannot be recommended.

Rapid I.V. injection can cause vasoconstriction producing hypertension and/or reduced coronary flow. A slow injection rate is therefore important in hypertensive heart failure and acute myocardial infarction.

4.3 CONTRAINDICATIONS:

Digoxin is contraindicated in:

- intermittent complete heart block or second degree atrioventricular block, especially if there is a history of Stokes-Adams attacks.
- Arrhythmias caused by cardiac glycoside intoxication.
- Supraventricular arrhythmias associated with an accessory atrioventricular pathway, as in the Wolff-Parkinson-White syndrome, unless the electrophysiological characteristics of the accessory pathway and any possible deleterious effect of digoxin on these characteristics have been evaluated. If an accessory pathway is known or suspected to be present and there is no history of previous supraventricular arrhythmias, digoxin is similarly contraindicated.
- Ventricular tachycardia or ventricular fibrillation.
- hypertrophic obstructive cardiomyopathy, unless there is concomitant atrial fibrillation and heart failure but even then caution should be exercised if digoxin is to be used.
- Hypersensitivity to the active substance, other digitalis glycosides or to any of the excipients listed in section 6.1.

4.4 SPECIAL WARNINGS AND PRECAUTIONS FOR USE:

Patients receiving digoxin should have their serum electrolytes and renal function (serum creatinine concentration) assessed periodically; the frequency of assessments will depend on the clinical setting.

Serum concentrations of digoxin may be expressed in Conventional Units of nanograms/ml or SI Units of nanomol/l. To convert nanograms/ml to nanomol/l, multiply nanograms/ml by 1.28.

The serum concentration of digoxin can be determined by radioimmunoassay.

Blood should be taken six hours or more after the last dose of digoxin.

There are no rigid guidelines as to the range of serum concentrations that are most efficacious. Post hoc analyses of heart failure patients in the Digitalis Investigation Group trial suggest that the optimal trough digoxin serum level may be 0.5 nanogram/ml (0.64 nanomol/l) to 1.0 nanogram/ml (1.28 nanomol/l).

**COMMON TECHNICAL DOCUMENT****PRODUCT: DIGOXIN INJECTION BP 0.25 mg/ml**

Digoxin toxicity is more commonly associated with serum digoxin concentrations greater than 2 nanogram/ml. However, serum digoxin concentration should be interpreted in the clinical context. Toxicity may occur with lower digoxin serum concentrations. In deciding whether a patient's symptoms are due to digoxin, the clinical state together with the serum potassium level and thyroid function are important factors (see Section 4.9).

Determination of the serum digoxin concentration may be very helpful in making a decision to treat with further digoxin, but other glycosides and endogenous digoxin-like substances, including metabolites of digoxin, can interfere with the assays that are available and one should always be wary of values which do not seem commensurate with the clinical state of the patient. Observations while temporary withholding digoxin might be more appropriate.

Arrhythmias

Arrhythmias may be precipitated by digoxin toxicity, some of which can resemble arrhythmias for which the drug could be advised. For example, atrial tachycardia with varying atrioventricular block requires particular care as clinically the rhythm resembles atrial fibrillation.

Many beneficial effects of digoxin on arrhythmias result from a degree of atrioventricular conduction blockade. However, when incomplete atrioventricular block already exists the effects of a rapid progression in the block should be anticipated. In complete heart block the idioventricular escape rhythm may be suppressed.

Sinoatrial disorder

In some cases of sinoatrial disorder (i.e. Sick Sinus Syndrome) digoxin may cause or exacerbate sinus bradycardia or cause sinoatrial block.

Myocardial infarction

The administration of digoxin in the period immediately following myocardial infarction is not contraindicated. However, the use of inotropic drugs in some patients in this setting may result in undesirable increases in myocardial oxygen demand and ischaemia, and some retrospective follow-up studies have suggested digoxin to be associated with an increased risk of death. The possibility of arrhythmias arising in patients who may be hypokalaemic after myocardial infarction and are likely to be haemodynamically unstable must be borne in mind. The limitations imposed thereafter on direct current cardioversion must also be remembered.

Cardiac amyloidosis

Treatment with digoxin should generally be avoided in patients with heart failure associated with cardiac amyloidosis. However, if alternative treatments are not appropriate, digoxin can be used to control the ventricular rate in patients with cardiac amyloidosis and atrial fibrillation.

**COMMON TECHNICAL DOCUMENT****PRODUCT: DIGOXIN INJECTION BP 0.25 mg/ml****Myocarditis**

Digoxin can rarely precipitate vasoconstriction and therefore should be avoided in patients with myocarditis.

Beri-beri heart disease

Patients with beri-beri heart disease may fail to respond adequately to digoxin if the underlying thiamine deficiency is not treated concomitantly.

Constrictive pericarditis

Digoxin should not be used in constrictive pericarditis unless it is used to control the ventricular rate in atrial fibrillation or to improve systolic dysfunction.

Exercise tolerance

Digoxin improves exercise tolerance in patients with impaired left ventricular systolic dysfunction and normal sinus rhythm. This may or may not be associated with an improved haemodynamic profile. However, the benefit of digoxin in patients with supraventricular arrhythmias is most evident at rest, less evident with exercise.

Withdrawal

In patients receiving diuretics and an ACE inhibitor, or diuretics alone, the withdrawal of digoxin has been shown to result in clinical deterioration.

Electrocardiography

The use of therapeutic doses of digoxin may cause prolongation of the PR interval and depression of the ST segment on the electrocardiogram.

Digoxin may produce false positive ST-T changes on the electrocardiogram during exercise testing. These electrophysiologic effects reflect an expected effect of the drug and are not indicative of toxicity.

Severe respiratory disease

Patients with severe respiratory disease may have an increased myocardial sensitivity to digitalis glycosides.

Hypokalaemia

**COMMON TECHNICAL DOCUMENT****PRODUCT: DIGOXIN INJECTION BP 0.25 mg/ml**

Hypokalaemia sensitises the myocardium to the actions of cardiac glycosides.

Hypoxia, hypomagnesaemia and hypercalcaemia

Hypoxia, hypomagnesaemia and marked hypercalcaemia increase myocardial sensitivity to cardiac glycosides.

Thyroid disease

Administering digoxin to a patient with thyroid disease requires care. Initial and maintenance doses of digoxin should be reduced when thyroid function is subnormal. In hyperthyroidism there is relative digoxin resistance and the dose may have to be increased. During the course of treatment of thyrotoxicosis, dosage should be reduced as the thyrotoxicosis comes under control.

Malabsorption

Patients with malabsorption syndrome or gastro-intestinal reconstructions may require larger doses of digoxin.

Chronic congestive cardiac failure

Although many patients with chronic congestive cardiac failure benefit from acute administration of digoxin, there are some in whom it does not lead to constant, marked or lasting hemodynamic improvement. It is therefore important to evaluate the response of each patient individually when digoxin is continued long-term.

Direct current cardioversion:

The risk of provoking dangerous arrhythmias with direct current cardioversion is greatly increased in the presence of digitalis toxicity and is in proportion to the cardioversion energy used.

For elective direct current cardioversion of a patient who is taking digoxin, the drug should be withheld for 24 h before cardioversion is performed. In emergencies, such as cardiac arrest when attempting cardioversion the lowest effective energy should be applied.

Direct current cardioversion is inappropriate in the treatment of arrhythmias thought to be caused by cardiac glycosides.

Excipients warnings

Ethanol

- Digoxin Solution for injection contains 10% by volume of ethanol (alcohol) or 0.2 ml of ethanol in each 2 ml ampoule (0.5 mg digoxin), i.e. up to 176 mg of ethanol per ampoule, equivalent to

**COMMON TECHNICAL DOCUMENT****PRODUCT: DIGOXIN INJECTION BP 0.25 mg/ml**

approximately 5 ml (one teaspoon) of beer, or less than 2 ml of wine. Harmful to those suffering from alcoholism. To be taken into account in pregnant or breast-feeding woman, children and high-risk groups such as patients with liver disease, or epilepsy.

4.5 INTERACTION WITH OTHER MEDICINAL PRODUCTS AND OTHER FORMS OF INTERACTION:

These may arise from effects on the renal excretion, tissue binding, plasma protein binding, and distribution within the body, gut absorptive capacity, P-glycoprotein activity and sensitivity to digoxin. Consideration of the possibility of an interaction whenever concomitant therapy is contemplated is the best precaution and a check on serum digoxin concentration is recommended when any doubt exists.

Digoxin is a substrate of P-glycoprotein. Thus, inhibitors of P-glycoprotein may increase blood concentrations of digoxin by enhancing its absorption and/or by reducing its renal clearance (see Section 5.2). Induction of P-glycoprotein can result in decreases in plasma concentrations of digoxin.

Combinations that should be avoided**Combinations which can increase effects of digoxin when co-administered:**

Digoxin, in association with beta-adrenoceptor blocking drugs, may increase atrio-ventricular conduction time.

Agents causing hypokalemia or intracellular potassium deficiency may cause increased sensitivity to digoxin; they include lithium salts, corticosteroids, carbenoxolone and some diuretics. Co-administration with diuretics such as loop or hydrochlorothiazide should be under close monitoring of serum electrolytes and renal function.

Calcium, particularly if administered rapidly by the I.V. route, may produce serious arrhythmias in digitalised patients.

Sympathomimetic drugs have direct positive chronotropic effects that can promote cardiac arrhythmias and may also lead to hypokalaemia, which can lead to or worsen cardiac arrhythmias. Concomitant use of digoxin and sympathomimetics may increase the risk of cardiac arrhythmias.

Combinations requiring caution**Combinations which can increase the effects of digoxin when co-administered:**

amiodarone, canagliflozin, daclatasvir, flibanserin, flecainide, prazosin, propafenone, quinidine, spironolactone, macrolide antibiotics e.g. erythromycin and clarythromycin, tetracycline (and possibly other antibiotics), gentamicin, isavuconazole, itraconazole, ivacaftor, quinine, trimethoprim, alprazolam, indomethacin, propantheline, mirabegron, nefazodone, atorvastatin, ciclosporine, epoprostenol (transient), vasopressin receptor antagonists (tolvaptan and conivaptan), carvedilol, ritonavir/ritonavir containing regimens, taleprevir, dronedarone, ranolazine, simeprevir,

**COMMON TECHNICAL DOCUMENT****PRODUCT: DIGOXIN INJECTION BP 0.25 mg/ml**

telmisartan, lapatinib, ticagrelor, vandetanib, velpatasvir, venetoclax and vemurafenib. Care should be taken when any of the above medicinal products are used in combination with digoxin. Serum digoxin concentrations should be monitored and used for titration of digoxin.

The concomitant use of digoxin and sennosides may be associated with a moderate increase in the risk of digoxin toxicity in heart failure patients.

Patients receiving digoxin are more susceptible to the effects of suxamethonium-exacerbated hyperkalaemia.

Co-administration of lapatinib with orally administered digoxin resulted in an increase in the AUC of digoxin. Caution should be exercised when dosing digoxin concurrently with lapatinib.

Drugs that modify afferent and efferent arteriole vascular tone may alter glomerular filtration. Angiotensin converting enzyme inhibitors (ACEIs) and angiotensin receptor blockers (ARBs) decrease angiotensin II-mediated efferent arteriole vasoconstriction, while non-steroidal anti-inflammatory drugs (NSAIDs) and cyclooxygenase-2 enzyme (COX-2) inhibitors decrease prostaglandin-mediated afferent arteriole vasodilation. ARBs, ACEIs, NSAIDs, and COX-2 inhibitors did not significantly alter digoxin pharmacokinetics or did not alter PK parameters in a consistent manner. However, these drugs may modify renal function in some patients, resulting in a secondary increase in digoxin.

Calcium channel blocking agents may either increase or cause no change in serum digoxin levels. Verapamil, felodipine and tiapamil increase serum digoxin levels. Nifedipine and diltiazem may increase or have no effect on serum digoxin levels while isradipine causes no change. Calcium channel blockers are also known to have depressant effects on sinoatrial and atrioventricular nodal conduction, particularly diltiazem and verapamil.

Proton pump inhibitors (PPI) are able to increase plasma levels of digoxin by inhibiting its efflux. Metabolism of digoxin in the gastrointestinal tract is inhibited by omeprazole, resulting in increased plasma levels of digoxin. Similar effects have been reported with pantoprazole and rabeprazole to a lesser extent.

Combinations which can decrease the effects of digoxin when co-administered:

Antacids, some bulk laxatives, kaolin-pectin, acarbose, neomycin, penicillamine, rifampicin, some cytostatics, metoclopramide, sulfasalazine, adrenaline, salbutamol, cholestyramine, phenytoin, St John's wort (*Hypericum perforatum*), bupropion and supplemental enteral nutrition.

Bupropion and its major circulating metabolite, with and without digoxin, stimulated OATP4C1-mediated digoxin transport. Digoxin has been identified as a substrate for aOATP4C1 in the basolateral side of the proximal renal tubules. Binding of bupropion and its metabolites to OATP4C1 could possibly increase the transport of digoxin and therefore, increase the renal secretion of digoxin.

**COMMON TECHNICAL DOCUMENT****PRODUCT: DIGOXIN INJECTION BP 0.25 mg/ml****Other interactions**

Milrinone does not alter steady-state serum digoxin levels.

4.6 PREGNANCY & LACTATION:**Pregnancy:**

The use of digoxin in pregnancy is not contraindicated, although the dosage may be less predictable in pregnant than in non-pregnant women, with some requiring an increased dosage of digoxin during pregnancy. As with all drugs, use should be considered only when the expected clinical benefit of treatment to the mother outweighs any possible risk to the developing foetus.

Despite extensive antenatal exposure to digitalis preparations, no significant adverse effects have been observed in the foetus or neonate when maternal serum digoxin concentrations are maintained within the normal range. Although it has been speculated that a direct effect of digoxin on the myometrium may result in relative prematurity and low birthweight, a contributing role of the underlying cardiac disease cannot be excluded. Maternally-administered digoxin has been successfully used to treat foetal tachycardia and congestive heart failure.

Adverse foetal effects have been reported in mothers with digitalis toxicity.

Breast-feeding

Although digoxin is excreted in breast milk, the quantities are minute and breast feeding is not contraindicated.

Fertility

There is no information available on the effect of digoxin on human fertility.

No data are available on whether or not digoxin has teratogenic effects.

4.7 Effects on ability to drive and use machines

Since central nervous system and visual disturbances have been reported in patients receiving digoxin, patients should exercise caution before driving, using machinery or participating in dangerous activities.

4.8 UNDESIRABLE EFFECTS:**Summary of the safety profile**

In general, the adverse reactions of digoxin are dose-dependent and occur at doses higher than those needed to achieve a therapeutic effect.

Hence, adverse reactions are less common when digoxin is used within the recommended dose range or therapeutic serum concentration range and when there is careful attention to concurrent medications and conditions.

Tabulated list of adverse reactions



COMMON TECHNICAL DOCUMENT

PRODUCT: DIGOXIN INJECTION BP 0.25 mg/ml

Adverse reactions are listed below by system organ class and frequency. Frequencies are defined as: Very common $\geq 1/10$

Common $\geq 1/100$ and $< 1/10$

Uncommon $\geq 1/1000$ and $< 1/100$

Rare $\geq 1/10,000$ and $< 1/1000$

Very rare $< 1/10,000$, including isolated reports.

Very common, common and uncommon events were generally determined from clinical trial data. The incidence in placebo was taken into account. Adverse drug reactions identified through post-marketing surveillance were considered to be rare or very rare (including isolated reports).

System Organ Class	Frequency	Side effects
Blood and lymphatic system disorders	Very rare	Thrombocytopenia
Metabolism and nutrition disorders	Very rare	Decreased appetite
Psychiatric disorders	Uncommon	Depression
	Very rare	Psychotic disorder, apathy, confusional state
Nervous system disorders	Common	Nervous system disorder, dizziness
	Very rare	Headache
Eye disorders	Common	Visual impairment (blurred vision or xanthopsia)
Cardiac disorders	Common	Arrhythmia, conduction disorder, bigeminy, trigeminy, PR prolongation, sinus bradycardia
	Very rare	Supraventricular tachyarrhythmia, atrial tachycardia (with or without block), supraventricular tachycardia (nodal arrhythmia) tachycardia, ventricular arrhythmia, ventricular extrasystoles, electrocardiogram ST segment depression
Gastrointestinal disorders	Common	Nausea, vomiting, diarrhoea
	Very rare	Intestinal ischaemia, gastrointestinal necrosis
Skin and subcutaneous tissue Disorders	Common	Rash*

**COMMON TECHNICAL DOCUMENT****PRODUCT: DIGOXIN INJECTION BP 0.25 mg/ml**

Reproductive system and breast disorders	Very rare	Gynaecomastia*
General disorders and administration site conditions	Very rare	Fatigue, malaise, asthenia

* See “Description of selected adverse reactions”

Description of selected adverse reactions

Skin and subcutaneous tissue disorders

Skin rashes of urticarial or scarlatiniform character may be accompanied by pronounced eosinophilia.

Reproductive system and breast disorders

Gynaecomastia can occur with long term administration.

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the Yellow Card Scheme

Website: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store.

4.9 OVERDOSE:**Symptoms and signs**

The symptoms and signs of toxicity are generally similar to those described in Section 4.8, but may be more frequent and can be more severe.

Signs and symptoms of digoxin toxicity become more frequent with levels above 2.0 nanograms/ml (2.56 nanomol/l although there is considerable inter-individual variation. However, in deciding whether a patient's symptoms are due to digoxin, the clinical state, together with serum electrolyte levels and thyroid function are important factors (see Section 4.2). In patients undergoing haemodialysis, digoxin use is associated with increased mortality; patients with low pre-dialysis potassium concentrations are most at risk.

Adults

In adults without heart disease, clinical observation suggests that an overdose of digoxin of 10 to 15 mg was the dose resulting in death of half of the patients. If more than 25 mg of digoxin was ingested by an adult without heart disease, death or progressive toxicity responsive only to digoxin-binding Fab antibody fragments resulted.

**COMMON TECHNICAL DOCUMENT****PRODUCT: DIGOXIN INJECTION BP 0.25 mg/ml****Cardiac manifestations**

Cardiac manifestations are the most frequent and serious sign of both acute and chronic toxicity. Peak cardiac effects generally occur 3 to 6 hours following overdose and may persist for the ensuing 24 hours or longer. Digoxin toxicity may result in almost any type of arrhythmia. Multiple rhythm disturbances in the same patient are common. These include paroxysmal atrial tachycardia with variable atrioventricular (AV) block, accelerated junctional rhythm, slow atrial fibrillation (with very little variation in the ventricular rate) and bi directional ventricular tachycardia.

Premature ventricular contractions (PVCs) are often the earliest and most common arrhythmia. Bigeminy or trigeminy also occur frequently.

Sinus bradycardia and other bradyarrhythmias are very common.

First, second, third degree heart blocks and AV dissociation are also common.

Early toxicity may only be manifested by prolongation of the PR interval.

Ventricular tachycardia may also be a manifestation of toxicity.

Cardiac arrest from asystole or ventricular fibrillation due to digoxin toxicity is usually fatal.

Acute massive digoxin overdose can result in mild to pronounced hyperkalaemia due to inhibition of the sodium-potassium ($\text{Na}^+\text{-K}^+$) pump. Hypokalaemia may contribute to toxicity (see Section 4.4).

Non-cardiac manifestations

Gastrointestinal symptoms are very common in both acute and chronic toxicity. The symptoms precede cardiac manifestations in approximately half of the patients in most literature reports. Anorexia, nausea and vomiting have been reported with an incidence up to 80 %. These symptoms usually present early in the course of an overdose.

Neurologic and visual manifestations occur in both acute and chronic toxicity. Dizziness, various CNS disturbances, fatigue and malaise are very common. The most frequent visual disturbance is an aberration of colour vision (predominance of yellow green). These neurological and visual symptoms may persist even after other signs of toxicity have resolved.

In chronic toxicity, non-specific non-cardiac symptoms, such as malaise and weakness, may predominate.

Paediatric population

In children aged 1 to 3 years without heart disease, clinical observation suggests that an overdose of digoxin of 6 to 10 mg was the dose resulting in death in half of the patients.

If more than 10 mg of digoxin was ingested by a child aged 1 to 3 years without heart disease, the outcome was uniformly fatal when Fab fragment treatment was not given.

Most manifestations of chronic toxicity in children occur during or shortly after digoxin overdose.

**COMMON TECHNICAL DOCUMENT****PRODUCT: DIGOXIN INJECTION BP 0.25 mg/ml****Cardiac manifestations**

The same arrhythmias or combination of arrhythmias that occur in adults can occur in paediatrics. Sinus tachycardia, supraventricular tachycardia, and rapid atrial fibrillation are seen less frequently in the paediatric population.

Paediatric patients are more likely to present with an AV conduction disturbance or a sinus bradycardia.

Ventricular ectopy is less common, however in massive overdose, ventricular ectopy, ventricular tachycardia and ventricular fibrillation have been reported.

In neonates, sinus bradycardia or sinus arrest and/or prolonged PR intervals are frequent signs of toxicity. Sinus bradycardia is common in young infants and children. In older children, AV blocks are the most common conduction disorders.

Any arrhythmia or alteration in cardiac conduction that develops in a child taking digoxin should be assumed to be caused by digoxin, until further evaluation proves otherwise.

Non-cardiac manifestations

The frequent non-cardiac manifestations are similar to those seen in adults are gastrointestinal, CNS and visual. However, nausea and vomiting are not frequent in infants and small children.

In addition to the undesirable effects seen with recommended doses, weight loss in older age groups and failure to thrive in infants, abdominal pain due to mesenteric artery ischaemia, drowsiness and behavioural disturbances including psychotic manifestations have been reported in overdose.

Treatment

After recent ingestion, such as accidental or deliberate self-poisoning, the load available for absorption may be reduced by gastric lavage. Gastric lavage increases vagal tone and may precipitate or worsen arrhythmias. Consider pre-treatment with atropine if gastric lavage is performed. Treatment with digitalis Fab antibody usually renders gastric lavage unnecessary. In the rare instances in which gastric lavage is indicated, it should only be performed by individuals with proper training and expertise.

Patients with massive digitalis ingestion should receive large doses of activated charcoal to prevent absorption and bind digoxin in the gut during enteroenteric recirculation.

If hypokalaemia is present, it should be corrected with potassium supplements either orally or intravenously, depending on the urgency of the situation. In cases where a large amount of digoxin has been ingested hyperkalaemia may be present due to release of potassium from skeletal muscle. Before administering potassium in digoxin overdose the serum potassium level must be known.

Bradyarrhythmias may respond to atropine but temporary cardiac pacing may be required. Ventricular arrhythmias may respond to lignocaine or phenytoin.

**COMMON TECHNICAL DOCUMENT****PRODUCT: DIGOXIN INJECTION BP 0.25 mg/ml**

Dialysis is not particularly effective in removing digoxin from the body in potentially life-threatening toxicity.

Digoxin-specific antibody Fab is a specific treatment for digoxin toxicity and is very effective. Rapid reversal of the complications that are associated with serious poisoning by digoxin, digitoxin and related glycosides has followed I.V. administration of digoxin-specific (ovine) antibody fragments (Fab). For details, consult the literature supplied with antibody fragments.

5. PHARMACOLOGICAL PROPERTIES**5.1 Pharmacodynamics Properties:**

Pharmacotherapeutic group: Cardiac therapy, cardiac glycosides, digitalis glycosides.

ATC code: C01AA05

Mechanism of action

Digoxin increases contractility of the myocardium by direct activity. This effect is proportional to dose in the lower range and some effect is achieved with quite low dosing; it occurs even in normal myocardium although it is then entirely without physiological benefit. The primary action of digoxin is specifically to inhibit adenosine triphosphatase, and thus sodium-potassium ($\text{Na}^{\text{-}}\text{-K}^{\text{+}}$) exchange activity, the altered ionic distribution across the membrane resulting in an augmented calcium ion influx and thus an increase in the availability of calcium at the time of excitation-contraction coupling. The potency of digoxin may therefore appear considerably enhanced when the extracellular potassium concentration is low, with hyperkalemia having the opposite effect.

Digoxin exerts the same fundamental effect of inhibition of the $\text{Na}^{\text{+}}\text{-K}^{\text{+}}$ exchange mechanism on cells of the autonomic nervous system, stimulating them to exert indirect cardiac activity. Increases in efferent vagal impulses result in reduced sympathetic tone and diminished impulse conduction rate through the atria and atrio-ventricular node. Thus, the major beneficial effect of digoxin is reduction of ventricular rate.

Intravenous administration of a loading dose produces an appreciable pharmacological effect within 5 to 30 mins, while using the oral route the onset of effect occurs in 0.5 to 2 hours.

Pharmacodynamic effects

The PROVED trial designed to determine the effectiveness of digoxin in 88 patients with chronic, stable mild to moderate heart failure. Withdrawal of digoxin or its continuation was performed in a prospective, randomized, double-blind, placebo-controlled multicenter trial of patients with chronic, stable mild to moderate heart failure secondary to left ventricular systolic dysfunction who had normal sinus rhythm and were receiving long-term treatment with diuretic drugs and digoxin. Patients withdrawn from digoxin therapy showed worsened maximal exercise capacity ($p = 0.003$) an increased incidence of treatment failures ($p = 0.039$) and a decreased time to treatment failure ($p = 0.037$). Patients who continued to receive digoxin had a lower body weight ($p = 0.044$) and heart rate ($p = 0.003$) and a higher left ventricular ejection fraction ($p = 0.016$). The overall percentage of participants having one or more adverse event was similar in the two groups: 59 %

**COMMON TECHNICAL DOCUMENT****PRODUCT: DIGOXIN INJECTION BP 0.25 mg/ml**

in the placebo group and 69 % in the digoxin group. The types of adverse event were unspecified

The RADIANCE trial examined the effects of discontinuation of digoxin in stable NYHA class II and III patients who were receiving diuretics and ACE inhibitors. The 178 patients were initially stabilised on a combination of captopril or enalapril, diuretics and digoxin, then randomised to continue digoxin therapy or change to placebo. The relative risk of worsening disease in the placebo group was 5.9 compared to the digoxin group. Withdrawal of digoxin was accompanied by worsening symptoms, reduced exercise tolerance, and a deteriorating quality of life, indicating that patients with CHF were at considerable risk from discontinuation of the drug in spite of the continuation of therapy with diuretics and ACE inhibitors. Approximately 56 % in the placebo group and 49% in the digoxin group experienced unspecified side effects.

In the DIG trial, 6800 patients with heart failure were randomised to receive digoxin or placebo. No difference was found in all-cause mortality between patients who were treated with digoxin and those who were given placebo. In the digoxin group, there was a trend toward a decrease in the risk of death attributed to worsening heart failure (risk ratio, 0.88; 95% confidence interval, 0.77 to 1.01; $p = 0.06$). However, the patients who received digoxin had significantly ($p < 0.001$) fewer hospital admissions when the drug was given in addition to diuretics and ACE inhibitors. Digoxin therapy was most beneficial in patients with ejection fractions of $\leq 25\%$, patients with enlarged hearts (cardiothoracic ratio of > 0.55), and patients in NYHA functional class III or IV. In the DIG study, 11.9 % of patients in the digoxin arm and 7.9 % of patients in the placebo arm were suspected of having digoxin toxicity, the most common symptoms being new episodes of ventricular fibrillation, supraventricular arrhythmia, tachycardia, or advanced atrioventricular block.

The AFFIRM study involved a total of 4060 patients recruited to a randomised, multicentre comparison of two treatment strategies in patients with atrial fibrillation and a high risk of stroke or death. The primary end point was overall mortality. There were 356 deaths among the patients assigned to rhythm-control therapy (amiodarone, disopyramide, flecainide, moricizine, procainamide, propafenone, quinidine, sotalol, and combinations of these drugs) and 310 deaths among those assigned to rate-control [β -blockers, calcium-channel blockers (verapamil and diltiazem), digoxin, and combinations of these drugs) therapy (mortality at five years, 23.8% and 21.3%, respectively; hazard ratio, 1.15 [95% confidence interval, 0.99 to 1.34]; $p = 0.08$). More patients in the rhythm-control group than in the rate-control group were hospitalised, and there were more adverse drug effects in the rhythm-control group as well.

Indirect cardiac contractility changes also result from changes in venous compliance brought about by the altered autonomic activity and by direct venous stimulation. The interplay between direct and indirect activity governs the total circulatory response, which is not identical for all subjects. In the presence of certain supraventricular arrhythmias, the neurogenically mediated slowing of AV conduction is paramount.

The degree of neurohormonal activation occurring in patients with heart failure is associated with clinical deterioration and an increased risk of death. Digoxin reduces activation of both the

**COMMON TECHNICAL DOCUMENT****PRODUCT: DIGOXIN INJECTION BP 0.25 mg/ml**

sympathetic nervous system and the (renin-angiotensin) system independently of its inotropic actions, and may thus favourably influence survival. Whether this is achieved via direct sympathoinhibitory effects or by re-sensitising baroreflex mechanisms remains unclear.

5.2 Pharmacokinetic properties:**Absorption**

The T_{max} following IV administration is approximately 1 to 5 hours, while the T_{max} for oral administration is 2 to 6 hours. Upon oral administration, digoxin is absorbed from the stomach and upper part of the small intestine. When digoxin is taken after meals, the rate of absorption is slowed, but the total amount of digoxin absorbed is usually unchanged. When taken with meals high in fibre, however, the amount absorbed from an oral dose may be reduced.

The bioavailability of orally administered digoxin is approximately 63 % in tablet form and 75 % as oral solution.

Distribution

The initial distribution of digoxin from the central to the peripheral compartment generally lasts from 6 to 8 h. This is followed by a more gradual decline in serum digoxin concentration, which is dependent upon digoxin elimination from the body. The volume of distribution is large ($V_{d_{ss}} = 510$ litres in healthy volunteers), indicating digoxin to be extensively bound to body tissues. The highest digoxin concentrations are seen in the heart, liver and kidney, that in the heart averaging 30-fold that in the systemic circulation. Although the concentration in skeletal muscle is far lower, this store cannot be overlooked since skeletal muscle represents 40 % of total body weight. Of the small proportion of digoxin circulating in plasma, approximately 25 % is bound to protein.

Biotransformation and Elimination**Biotransformation**

The majority of digoxin is excreted by the kidneys as an intact drug, although a small fraction of the dose is metabolised to pharmacologically active and inactive metabolites. The main metabolites of digoxin are dihydrodigoxin and digoxigenin.

Elimination

The major route of elimination is renal excretion of the unchanged drug.

Digoxin is a substrate for P-glycoprotein. As an efflux protein on the apical membrane of enterocytes, P-glycoprotein may limit the absorption of digoxin. P-glycoprotein in renal proximal tubules appears to be an important factor in the renal elimination of digoxin (see Section 4.5).

Following I.V. administration to healthy volunteers, between 60 and 75 % of a digoxin dose is recovered unchanged in the urine over a six day follow-up period. Total body clearance of digoxin has been shown to be directly related to renal function, and percent daily loss is thus a function of creatinine clearance. The total and renal clearances of digoxin have been found to be 193 ± 25

**COMMON TECHNICAL DOCUMENT****PRODUCT: DIGOXIN INJECTION BP 0.25 mg/ml**

ml/min and 152 ± 24 ml/min in a healthy control population.

In a small percentage of individuals, orally administered digoxin is converted to cardioinactive reduction products (digoxin reduction products or DRPs) by colonic bacteria in the gastrointestinal tract. In these subjects over 40 % of the dose may be excreted as DRPs in the urine. Renal clearances of the two main metabolites, dihydrodigoxin and digoxigenin, have been found to be 79 ± 13 ml/min and 100 ± 26 ml/min, respectively.

In the majority of cases however, the major route of digoxin elimination is renal excretion of the unchanged drug.

The terminal elimination half-life of digoxin in patients with normal renal function is 30 to 40 h.

Since most of the drug is bound to the tissues rather than circulating in the blood, digoxin is not effectively removed from the body during cardiopulmonary by-pass. Furthermore, only about 3 % of a digoxin dose is removed from the body during 5 h of haemodialysis.

Special patient populations**Paediatric population**

In the newborn period, renal clearance of digoxin is diminished and suitable dosage adjustments must be observed. This is especially pronounced in the premature infant since renal clearance reflects maturation of renal function. Digoxin clearance has been found to be 65.6 ± 30 ml/min/ 1.73m^2 at three months, compared to only 32 ± 7 ml/min/ 1.73m^2 at one week. By 12 months digoxin clearance of 88 ± 43 ml / min / 1.73m^2 has been reported. Beyond the immediate newborn period, children generally require proportionally larger doses than adults on the basis of body weight and body surface area.

Renal impairment

The terminal elimination half-life of digoxin is prolonged in patients with impaired renal function, and in anuric patients may be of the order of 100 h.

Hepatic impairment

Hepatic impairment has little effect on digoxin clearance.

Elderly

Age-related declines in renal function in elderly patients can result in a lower rates of digoxin clearance than in younger subjects, with reported digoxin clearance rates in the elderly of 53 ml/min/ 1.73m^2 .

Gender

Digoxin clearance is 12% – 14% less in females than males and may need to be considered in dosing calculations.

**COMMON TECHNICAL DOCUMENT****PRODUCT: DIGOXIN INJECTION BP 0.25 mg/ml****5.3 Preclinical Safety Data:****Carcinogenesis, mutagenesis**

Digoxin showed no genotoxic potential in in vitro studies (Ames test and mouse lymphoma). No data are available on the carcinogenic potential of digoxin.

6. PHARMACEUTICAL PARTICULARS**6.1 List of excipients**

Ethanol

Propylene Glycol

Citric Acid Monohydrate

Sodium phosphate anhydrous

or Sodium phosphate

Water for Injection

6.2 Incompatibilities:

None known.

6.3 Shelf Life:

3 years.

When Digoxin injection is diluted at a ratio of 1 to 250, it is stable for up to 48 hours at room temperature (20 to 25 °C).

6.3 Special Precaution for Storage:

Store below 30 °C Protect from light, keep out of reach of children

6.4 Nature and contents of container:

5 x 2 ml Amber USP type I Ampoule.

6.5 Special precautions for disposal and other handling:**Digoxin injection:**

Digoxin injection can be administered undiluted or diluted with a 4-fold or greater volume of 0.9% Sodium Chloride Injection, 0.18 % Sodium Chloride/4% Glucose Injection or 5% Glucose Injection. A 4-fold volume of diluent equates to adding one 2 ml ampoule of digoxin to 6 ml of injection solution. The use of less than a 4-fold volume of diluent could lead to precipitation of digoxin.

Digoxin injection, when diluted in the ratio of 1 to 250 is known to be compatible with the following infusion solutions;

Sodium Chloride I.V. Infusion, B.P., 0.9 % w/v.



COMMON TECHNICAL DOCUMENT

PRODUCT: DIGOXIN INJECTION BP 0.25 mg/ml

Sodium Chloride (0.18 % w/v) and Glucose (4 % w/v) I.V. Infusion, B.P.

Glucose I.V. Infusion, B.P., 5 % w/v.

A ratio of 1 to 250 can be obtained for example by diluting one 2 ml ampoule with 500 ml of infusion solution.

Dilution should be carried out either under full aseptic conditions immediately before use. Any unused solution should be discarded.

7. MARKETING AUTHORISATION HOLDER

CIRON DRUGS & PHARMACEUTICALS PVT.LTD.

C- 1101/1102, Lotus Corporate Park,

Graham Firth Steel Compound, Jay Coach Junction,

Western Express Highway, Goregaon (East)

Mumbai, Maharashtra, India – 400063

Tel: +91-22-3359800

E-mail:mail@cironpharma.com

8. MARKETING AUTHORISATION NUMBER(S)

None

9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

Not applicable