



LEAFLET

AGODIC GEL (Diclofenac Gel BP 1% w/w)

COMPOSITION :
Diclofenac Diethylamine BP..... 1.16% w/w
Equivalent to Diclofenac Sodium BP..... 1.0 % w/w
Gel base q.s.

INDICATIONS :
• Agodic Gel is used to treat pain and reduce swelling.
• Used in the treatment of joint pain.
• Used in the treatment of muscle pain.
• Used in sprains.
• Used in soft tissue injury.
• Used in the treatment of Rheumatism and mild arthritis.

DIRECTION FOR USE :
• Apply the gel 3-4 times a day.
• Gently rub on the affected area.
• Agodic Gel quickly absorbed without the need to massage.

CONTRA-INDICATIONS :
• Do not use if patient is allergic to Diclofenac, aspirin, and any other similar medicines known as NSAIDs.
• If patient is taking other products containing Diclofenac, aspirin or other NSAID medicines by topical or oral route.

WARNINGS AND PRECAUTIONS :
• Take special care and check with your doctor before you use this medicine if you are pregnant or breastfeeding.
• Have asthma or allergic disease.
• If the patient have or ever had liver or kidney disease. There have been rare reports of hepatitis with topical use of Diclofenac.
• If the patient have or ever had a stomach ulcer, perforation or bleeding of the stomach, as there is a possibility of bleeding or perforation of ulcer.

- Keep away from eyes, nose and mouth.
- Avoid use on broken skin, the eyes, nose and lips.
- Avoid exposing the treated area to sunlight or sun lamps.
- Avoid exposure to sunlight or artificial UV rays.
- Do not cover the treated area with bandages or dressings.
- Do not use over large areas.
- Do not apply to the same area at the same time as other skin products (e.g. moisturizing cream).
- Wash hands after use.
- For External use only.

OVERDOSE :
If you swallow the gel, seek medical advice immediately.

PREGNANCY AND LACTATION :
• Consult a doctor prior to using this during pregnancy.
• Diclofenac gel is not recommended for use during lactation.

SIDE-EFFECTS :
When using this product the patient may experience Redness, swelling or skin irritation at the site of application. Stop using this medicine and tell your doctor immediately if allergic reactions such as generalized skin rashes and itching. Sometimes with breathing problems or swelling of the lips, throat or face seek medical advice.

STORAGE :
Store under normal storage conditions (15°C to 30°C).
Protect from light.

Keep all medicines out of reach of children.

PRESENTATION :
30 gm tube in a carton.



Manufactured in India by :

AGOG PHARMA LTD.

Plot No. 33, Sector II, The Vasai Taluka Indl. Co-op. Estate Ltd., Vasai (E), Dist. Thane. INDIA.



1.5.3 Patient Information Leaflet (PIL)

AGODIC GEL (Diclofenac Gel BP 1% w/w)

COMPOSITION :
Diclofenac Diethylamine BP..... 1.16% w/w
Equivalent to Diclofenac Sodium BP..... 1.0 % w/w
Gel base Q.S.

INDICATIONS :
• Agodic Gel is used to treat pain and reduce swelling.
• Used in the treatment of joint pain.
• Used in the treatment of muscle pain.
• Used in sprains.
• Used in soft tissue injury.
• Used in the treatment of Rheumatism and mild arthritis.

DIRECTION FOR USE :
• Apply the gel 3-4 times a day.
• Gently rub on the affected area.
• Agodic Gel quickly absorbed without the need to massage.

CONTRA-INDICATIONS :
• Do not use if patient is allergic to Diclofenac, aspirin, and any other similar medicines known as NSAIDs.
• If patient is taking other products containing Diclofenac, aspirin or other NSAID medicines by topical or oral route.

WARNINGS AND PRECAUTIONS :
• Take special care and check with your doctor before you use this medicine if you are pregnant or breastfeeding.
• Have asthma or allergic disease.
• If the patient have or ever had liver or kidney disease. There have been rare reports of hepatitis with topical use of Diclofenac.
• If the patient have or ever had a stomach ulcer, perforation or bleeding of the stomach, as there is a possibility of bleeding or perforation of ulcer.

- Keep away from eyes, nose and mouth.
- Avoid use on broken skin, the eyes, nose and lips.
- Avoid exposing the treated area to sunlight or sun lamps.
- Avoid exposure to sunlight or artificial UV rays.
- Do not cover the treated area with bandages or dressings.
- Do not use over large areas.
- Do not apply to the same area at the same time as other skin products (e.g. moisturizing cream).
- Wash hands after use.
- For External use only.

OVERDOSE :
If you swallow the gel, seek medical advice immediately.

PREGNANCY AND LACTATION :
• Consult a doctor prior to using this during pregnancy.
• Diclofenac gel is not recommended for use during lactation.

SIDE-EFFECTS :
When using this product the patient may experience Redness, swelling or skin irritation at the site of application. Stop using this medicine and tell your doctor immediately. If allergic reactions such as generalized skin rashes and itching, Sometimes with breathing problems or swelling of the lips, throat or face seek medical advice.

STORAGE :
Store under normal storage conditions (15°C to 30°C).
Protect from light.

Keep all medicines out of reach of children.

PRESENTATION :
30 gm tube in a carton.



Manufactured in India by :

AGOG PHARMA LTD.

Plot No. 33, Sector II, The Vasai Taluka Indl. Co-op. Estate Ltd., Vasai (E), Dist. Thane. INDIA.