

**Module 1****PIRICAM 20****(Piroxicam Capsules USP 20mg)****1.6 PRODUCT INFORMATION****MMC Healthcare Ltd.**No.: 34-B, SIDCO Industrial Estate,  
Thirumazhisai, Chennai-600 124, India.**1.6.1 Prescribing Information (Summary of Product Characteristics)****1. Name of the medicinal product****Trade Name : PIRICAM 20****Generic Name : Piroxicam Capsules USP 20 mg****2. Qualitative and quantitative composition:****A. Qualitative Composition**

Each hard gelatin capsules contains:

Piroxicam USP 20mg

**B. Quantitative Composition****Batch Size: 1, 50,000 Capsules**

Sr. No.	Material Name	Spec.	Quantity (mg)/ Capsule	% For- mula w/w	Theoretical Quantity / Batch (kg)	Ovg. in %/#	Standard Quantity / Batch (kg)
<b>Granulation / Mixing of Piroxicam</b>							
1	Piroxicam	USP	21.00	8.4	3.150	5%	3.150
2	Lactose Monohydrate	BP	146.81	58.72	22.021	-	22.021
3	Povidone K 30	BP	11.13	4.45	1.670	-	1.670
4	Purified water	BP	--	--	10.0Lt		10.0Lt
<b>Lubrication/Encapsulation</b>							
5	Microcrystalline cellulose	BP	40.00	16.00	6.000	-	6.000
6	Sodium lauryl sulfate	BP	4.50	1.80	0.675	-	0.675
7	Croscarmellose sodium	BP	11.13	4.45	1.670	-	1.670
8	Magnesium stearate	BP	2.25	0.90	0.338	-	0.338
9	Colloidal anhydrous silica	BP	2.25	0.90	0.338	-	0.338
10	Purified Talc	BP	2.25	0.90	0.338	-	0.338
11	Dark Red/Deep Red '2' size hard gelatin	IHS	--	--	155000 Capsules	-	155000 Capsules
12	Croscarmellose sodium	BP	8.67	3.47	1.300	-	1.300
<b>TOTAL</b>			<b>250.00mg /Capsule</b>		<b>37.50kg/ Batch</b>		<b>37.50kg/ Batch</b>

\* Maximum quantity of Croscarmellose sodium required to adjust weight of granules due to loss on drying.

\*Not present in the finished product

#5% overages used for Manufacturing Loss

**3. Pharmaceutical form**

Hard gelatin capsules

White colour powder filled in Dark red / Deep red colour '2' size Hard Gelatin capsules.

**4. Clinical particulars**

## Module 1

### PIRICAM 20

#### (Piroxicam Capsules USP 20mg)

Non steroidal Anti-inflammatory drug

ATC code: M01AC01.



MMC Healthcare Ltd.

No.: 34-B, SIDCO Industrial Estate,  
Thirumazhisai, Chennai-600 124, India.

#### 4.1 Indications

Piroxicam is indicated for symptomatic relief of osteoarthritis, rheumatoid arthritis, or ankylosing spondylitis.

Due to its safety profile, piroxicam is not a first line option should an NSAID be indicated. The decision to prescribe piroxicam should be based on an assessment of the individual patient's overall risks.

#### 4.2 Posology and method of administration

##### Posology

The prescription of piroxicam should be initiated by physicians with experience in the diagnostic evaluation and treatment of patients with inflammatory or degenerative rheumatic diseases.

The maximum recommended daily dose is 20mg.

Undesirable effects may be minimised by using the minimum effective dose for the shortest duration necessary to control symptoms. The benefit and tolerability of treatment should be reviewed within 14 days. If continued treatment is considered necessary, this should be accompanied by frequent review.

Given that piroxicam has been shown to be associated with an increased risk of gastrointestinal complications, the need for possible combination therapy with gastroprotective agents (e.g. misoprostol or proton pump inhibitors) should be carefully considered, in particular for elderly patients.

##### Elderly

Elderly, frail or debilitated patients may tolerate side-effects less well and such patients should be carefully supervised. As with other NSAIDs, caution should be used in the treatment of elderly patients who are more likely to be suffering from impaired renal, hepatic or cardiac function.

##### Method of Administration

For oral administration. To be taken preferably with or after food.

Undesirable effects may be minimised by using the lowest effective dose for the shortest duration necessary to control symptoms.

#### 4.3 Contraindications

Hypersensitivity to the active substance or to any of the excipients.

- History of gastro-intestinal ulceration, bleeding or perforation.
- Patient history of gastrointestinal disorders that predispose to bleeding disorders such as ulcerative colitis, Crohn's disease, gastrointestinal cancers or diverticulitis.
- Patients with active peptic ulcer, inflammatory gastrointestinal disorder or gastrointestinal bleeding.
- Concomitant use with other NSAIDs, including COX-2 selective NSAIDs and acetylsalicylic acid at analgesic doses.
- Concomitant use with anticoagulants.

## Module 1

### PIRICAM 20

#### (Piroxicam Capsules USP 20mg)



- History of previous serious allergic drug reaction of any type, especially cutaneous reactions such as erythema multiforme, Stevens-Johnson syndrome, toxic epidermal necrolysis.
- Previous skin reaction (regardless of severity) to piroxicam, other NSAIDs and other medications.
- Patients in whom aspirin and other non-steroidal anti-inflammatory drugs induce the symptoms of asthma, nasal polyps, angioedema or urticaria.
- Severe heart failure.
- During the last trimester of pregnancy.

#### 4.4 Special warnings and precautions for use

Undesirable effects may be minimised by using the minimum effective dose for the shortest duration necessary to control symptoms (see section 4.2, and GI and cardiovascular risks below).

The clinical benefit and tolerability should be re-evaluated periodically and treatment should be immediately discontinued at the first appearance of cutaneous reactions or relevant gastrointestinal events.

Gastrointestinal (GI) Effects, Risk of GI Ulceration, Bleeding, and Perforation NSAIDs, including piroxicam, can cause serious gastrointestinal events including bleeding, ulceration, and perforation of the stomach, small intestine or large intestine, which can be fatal. These serious adverse events can occur at any time, with or without warning symptoms, in patients treated with NSAIDs.

NSAID exposures of both short and long duration have an increased risk of serious GI event. Evidence from observational studies suggests that piroxicam may be associated with a high risk of serious gastrointestinal toxicity, relative to other NSAIDs.

Patients with significant risk factors for serious GI events should be treated with piroxicam only after careful consideration.

The possible need for combination therapy with gastro-protective agents (e.g. misoprostol or proton pump inhibitors) should be carefully considered.

#### Serious GI Complications

Identification of at-risk subjects: The risk for developing serious GI complications increases with age. Age over 70 years is associated with high risk of complications. The administration to patients over 80 years should be avoided.

Patients taking concomitant oral corticosteroids, selective serotonin reuptake inhibitors (SSRIs) or anti-platelet agents such as low-dose acetylsalicylic acid are at increased risk of serious GI complications (see below and section 4.5). As with other NSAIDs, the use of piroxicam in combination with protective agents (e.g. misoprostol or proton pump inhibitors) must be considered for these at-risk patients.

Patients and physicians should remain alerted for signs and symptoms of GI ulceration and/or bleeding during piroxicam treatment. Patients should be asked to report any new or unusual abdominal symptom during treatment. If a gastrointestinal complication is suspected during treatment, piroxicam should be discontinued immediately and additional clinical evaluation and treatment should be considered.

**Module 1**  
**PIRICAM 20**

**(Piroxicam Capsules USP 20mg)**



Appropriate monitoring and advice are required for patients with a history of hypertension and/or mild to moderate congestive heart failure as fluid retention and oedema have been reported in association with NSAID therapy.

Patients with uncontrolled hypertension, congestive heart failure, established ischaemic heart disease, peripheral arterial disease, and/or cerebrovascular disease should only be treated with piroxicam after careful consideration. Similar consideration should be made before initiating longer-term treatment of patients with risk factors for cardiovascular events (e.g. hypertension, hyperlipidaemia, diabetes mellitus, smoking).

Clinical trial and epidemiological data suggest that use of some NSAIDs (particularly at high doses and in long term treatment) may be associated with a small increased risk of arterial thrombotic events (for example myocardial infarction or stroke). There are insufficient data to exclude such a risk for Piroxicam.

Piroxicam should be used with caution in patients with a history of bronchial asthma.

Patients who are known or suspected to be poor CYP2C9 metabolizers based on previous history/experience with other CYP2C9 substrates should be administered piroxicam with caution as they may have abnormally high plasma levels due to reduced metabolic clearance

**Skin reactions**

Life-threatening cutaneous reactions Stevens-Johnson syndrome (SJS) and toxic epidermal necrolysis (TEN) have been reported with the use of piroxicam.

Patients should be advised of the signs and symptoms and monitored closely for skin reactions. The highest risk for occurrence of SJS or TEN is within the first weeks of treatment.

If symptoms or signs of SJS or TEN (e.g. progressive skin rash often with blisters or mucosal lesions) are present, piroxicam treatment should be discontinued. The best results in managing SJS and TEN come from early diagnosis and immediate discontinuation of any suspect drug. Early withdrawal is associated with a better prognosis.

If the patient has developed SJS or TEN with the use of piroxicam, piroxicam must not be re-started in this patient at any time.

Serious skin reactions, some of them fatal, including exfoliative dermatitis, Stevens-Johnson syndrome, and toxic epidermal necrolysis, have been reported very rarely in association with the use of NSAIDs (see section 4.8). Evidence from observational studies suggests that piroxicam may be associated with a higher risk of serious skin reaction than other non-oxicam NSAIDs. Patients appear to be at a highest risk of these reactions early in the course of therapy, the onset of the reaction occurring in the majority of cases within the first month of treatment. Piroxicam should be discontinued at the first appearance of skin rash, mucosal lesions, or any other sign of hypersensitivity.

Piroxicam should be used with caution in patients with renal, hepatic and cardiac impairment. In rare cases, non-steroidal anti-inflammatory drugs may cause interstitial nephritis, glomerulitis, papillary necrosis and the nephrotic syndrome. Such agents inhibit the synthesis of the prostaglandin which plays a supportive role in the maintenance of renal perfusion in patients whose renal blood flow and blood volume are decreased. In these patients, administration of a non-steroidal anti-inflammatory drug may precipitate overt renal decompensation, which is typically followed by recovery to pre-treatment state upon discontinuation of non-steroidal anti-inflammatory therapy. Patients at greatest risk of such a reaction are with congestive heart failure, liver cirrhosis, nephrotic syndrome and overt renal disease; such patients should be carefully monitored whilst receiving NSAID therapy.

**Module 1**  
**PIRICAM 20**

**(Piroxicam Capsules USP 20mg)**



Because of reports of adverse eye findings with non-steroidal anti-inflammatory drugs, it is recommended that patients who develop visual complaints during treatment with Piroxicam have ophthalmic evaluation.

**Impaired female fertility**

The use of piroxicam may impair female fertility and is not recommended in women attempting to conceive. In women who have difficulties conceiving or who are undergoing investigation of infertility, withdrawal of Piroxicam should be considered.

**Lactose**

Piroxicam contain lactose. Patients with rare hereditary problems of galactose intolerance, total lactase deficiency or glucose-galactose malabsorption should not take this medicine.

**Sodium**

This medicine contains less than 1 mmol sodium (23 mg) per capsule, that is to say essentially 'sodium-free'.

**4.5 Interaction with other medicinal products and other forms of interaction**

**Antacids:** Concomitant administration of antacids had no effect on piroxicam plasma levels.

**Anti-coagulants:** NSAIDs, including piroxicam, may enhance the effects of anticoagulants, such as warfarin. Therefore, the use of piroxicam with concomitant anticoagulant such as warfarin should be avoided.

**Anti-platelet agents and selective serotonin reuptake inhibitors (SSRIs):** Increased risk of gastrointestinal bleeding.

**Aspirin and other Non-Steroidal Anti-Inflammatory Drugs:** Piroxicam, like other non-steroidal anti-inflammatory drugs decreases platelet aggregation and prolongs bleeding time. This effect should be kept in mind when bleeding times are determined.

As with other NSAIDs, the use of piroxicam together with acetylsalicylic acid or concomitant use with other NSAIDs, including other piroxicam formulations, must be avoided, since data are inadequate to show that combinations produce greater improvement than that achieved with piroxicam alone; moreover, the potential for adverse reactions is enhanced. Human studies have shown that concomitant use of piroxicam and acetylsalicylic acid reduces the plasma piroxicam concentration to about 80% of the usual value.

**Cardiac glycosides:** NSAIDs may exacerbate cardiac failure, reduce GFR and increase plasma glycoside levels.

**Ciclosporin, Tacrolimus:** possible increased risk of nephrotoxicity when NSAIDs are given with ciclosporin or tacrolimus.

**Cimetidine:** Results of two separate studies indicate a slight but significant increase in absorption of piroxicam following cimetidine administration but no significant changes in elimination rate constants or half-life. The small increase in absorption is unlikely to be clinically significant.

**Corticosteroids:** increased risk of gastrointestinal ulceration or bleeding.

**Digoxin, Digitoxin:** Concurrent therapy with piroxicam and digoxin, or piroxicam and digitoxin, did not affect the plasma levels of either drug.

**Diuretics:** Non-steroidal anti-inflammatory drugs may cause sodium, potassium and fluid retention and may interfere with the natriuretic action of diuretic agents. These properties

## Module 1

### PIRICAM 20

#### (Piroxicam Capsules USP 20mg)

should be kept in mind when treating patients with compromised cardiac function or hypertension since they may be responsible for the worsening of those conditions.

**Highly protein-bound drugs:** Piroxicam is highly protein-bound and therefore might be expected to displace other protein-bound drugs. The physician should closely monitor patients for change when administering Piroxicam to patients on highly protein-bound drugs.

**Lithium:** Non-steroidal anti-inflammatory drugs, including Piroxicam, have been reported to increase steady state plasma lithium levels. It is recommended that these levels are monitored when initiating, adjusting and discontinuing Piroxicam.

Piroxicam, like other non-steroidal anti-inflammatory drugs, may interact with the following drugs / classes of therapeutic agents:

Antihypertensives - antagonism of the hypotensive effect

Methotrexate - Reduced excretion of methotrexate, possibly leading to acute toxicity

Quinolone antibiotics - possible increased risk of convulsions

Mifepristone - NSAIDs could interfere with mifepristone-mediated termination of pregnancy.

#### **4.6 Fertility, pregnancy and lactation**

##### Pregnancy

Although no teratogenic effects were seen in animal testing, the safety of Piroxicam during pregnancy or during lactation has not yet been established. Piroxicam inhibits prostaglandin synthesis and release through a reversible inhibition of the cyclo-oxygenase enzyme. This effect, as with other non-steroidal anti-inflammatory drugs (NSAIDs) has been associated with an increased incidence of dystocia and delayed parturition in pregnant animals when drug administration was continued in late pregnancy. In view of the known effects of NSAIDs on the foetal cardiovascular system (risk of closure of the ductus arteriosus), use in the last trimester of pregnancy is contraindicated. The onset of labour may be delayed and the duration increased with an increased bleeding tendency in both mother and child.

Inhibition of prostaglandin synthesis might adversely affect pregnancy. Data from epidemiological studies suggest an increased risk of spontaneous abortion after use of prostaglandin synthesis inhibitors in early pregnancy. In animals, administration of prostaglandin synthesis inhibitors has been shown to result in increased pre- and post-implantation loss.

NSAIDs should not be used during the first two trimesters of pregnancy or labour unless the potential benefit to the patient outweighs the potential risk to the foetus.

##### Breast-feeding

A study indicates that piroxicam appears in breast milk at about 1-3% of the maternal plasma concentrations. No accumulation of piroxicam occurred in milk relative to that in plasma during treatment for up to 52 days. Piroxicam is not recommended for use in nursing mothers as clinical safety has not been established.

##### Fertility

Based on the mechanism of action, the use of NSAIDs, including Piroxicam, may delay or prevent rupture of ovarian follicles, which has been associated with reversible infertility in some women. In women who have difficulties conceiving or who are undergoing investigation of infertility, withdrawal of NSAIDs, including Piroxicam, should be considered.



MMC Healthcare Ltd.

No.: 34-B, SIDCO Industrial Estate,  
Thirumazhisai, Chennai-600 124, India.

**Module 1  
PIRICAM 20**

**(Piroxicam Capsules USP 20mg)**



**MMC Healthcare Ltd.**  
No.: 34-B, SIDCO Industrial Estate,  
Thirumazhisai, Chennai-600 124, India.

**4.7 Effects on ability to drive and use machines**

Undesirable effects such as dizziness, drowsiness, fatigue and visual disturbances are possible after taking NSAIDs. If affected, patients should not drive or operate machinery.

**4.8 Undesirable effects**

<b>System Organ Class</b>	<b>Very Common ≥1/10</b>	<b>Common ≥1/100 to &lt;1/10</b>	<b>Uncommon ≥1/1000 to &lt;1/100</b>	<b>Rare ≥1/10 000 to &lt;1 000</b>	<b>Very Rare &lt;1/1000 0</b>	<b>Not Known (cannot be estimated from available data)</b>
<b>Blood and lymphatic system disorders</b>		Anaemia Eosinophilia Leucopenia Thrombocytopenia				Aplastic anaemia Haemolytic anaemia
<b>Immune system disorders</b>						Anaphylaxis Serum sickness
<b>Metabolism and nutrition disorders</b>		Anorexia Hyperglycaemia	Hypoglycaemia			Fluid retention
<b>Psychiatric disorders</b>						Depression Dream abnormalities Hallucinations Insomnia Mental confusion Mood alterations Nervousness
<b>Nervous system disorders</b>		Dizziness Headache Somnolence Vertigo				Paresthesia
<b>Eye disorders</b>			Blurred vision			Eye irritations Swollen eyes
<b>Ear and labyrinth disorders</b>		Tinnitus				Hearing impairment
<b>Cardiac disorders</b>			Palpitations			Cardiac failure Arterial thrombotic events
<b>Vascular disorders</b>						Vasculitis Hypertension
<b>Respiratory, thoracic and mediastinal disorders</b>						Bronchospasm Dyspnoea Epistaxis
<b>Gastrointestinal disorders</b>		Abdominal discomfort Abdominal pain Constipation	Stomatitis			Gastritis Gastrointestinal bleeding (including hematemesis and

**Module 1**  
**PIRICAM 20**  
**(Piroxicam Capsules USP 20mg)**

		Diarrhoea Epigastric distress Flatulence Nausea Vomiting Indigestion				melen) Pancreatitis Perforation Ulceration
<b>Hepatobiliary disorders</b>						Fatal hepatitis Jaundice
<b>Renal and urinary disorders</b>				Interstitial nephritis Nephrotic syndrome Renal failure Renal papillary necrosis		
<b>Skin and subcutaneous tissue disorders</b>		Pruritis Skin rash			Severe cutaneous adverse reactions (SCARs): Stevens-Johnson syndrome (SJS) and toxic epidermal necrolysis (TEN)	Alopecia Angioedema Dermatitis exfoliative Erythema multiforme Non-thrombocytopenic purpura (Henoch-Schoenlein) Onycholysis Photoallergic reactions Urticaria Vesiculo bullous reactions
<b>Reproductive system and breast disorders</b>						Female fertility decreased
<b>General disorders and administration site conditions</b>		Oedema (mainly of the ankle)				Malaise
<b>Investigations</b>		Increased serum				Positive ANA Weight decrease



		transaminase levels Weight increase				Decreases in hemoglobin and hematocrit unassociated with obvious gastrointestinal bleeding
--	--	--	--	--	--	--

Gastrointestinal: These are the most commonly encountered side-effects but in most instances do not interfere with the course of therapy.

Objective evaluations of gastric mucosa appearances and intestinal blood loss show that 20mg/day of Piroxicam administered either in single or divided doses is significantly less irritating to the gastrointestinal tract than aspirin.

Some epidemiological studies have suggested that piroxicam is associated with higher risk of gastrointestinal adverse reactions compared with some NSAIDs, but this has not been confirmed in all studies. Administration of doses exceeding 20mg daily (of more than several days duration) carries an increased risk of gastrointestinal side effects, but they may also occur with lower doses.

Cardiac: Oedema, hypertension, and cardiac failure, have been reported in association with NSAID treatment. The possibility of precipitating congestive heart failure in elderly patients or those with compromised cardiac function should therefore be borne in mind.

Clinical trial and epidemiological data suggest that use of some NSAIDs (particularly at high doses and in long term treatment) may be associated with a small increased risk of arterial thrombotic events (for example, myocardial infarction or stroke).

Liver function: Changes in various liver function parameters have been observed. Although such reactions are rare, if abnormal liver function tests persist or worsen, if clinical symptoms consistent with liver disease develop, or if systemic manifestations occur (e.g. eosinophilia, rash etc.), piroxicam should be discontinued.

Other:

Routine ophthalmoscopy and slit-lamp examination have revealed no evidence of ocular changes.

#### **4.9 Overdose**

In the event of overdosage with Piroxicam, supportive and symptomatic therapy is indicated. Studies indicate that administration of activated charcoal may result in reduced re-absorption of piroxicam, thus reducing the total amount of active drug available.

Although there are no studies to date, haemodialysis is probably not useful in enhancing elimination of piroxicam since the drug is highly protein-bound.

### **5. Pharmacological properties**

#### **5.1 Pharmacodynamic properties**

Piroxicam is a non-steroidal anti-inflammatory agent which also possesses analgesic and antipyretic properties. Oedema, erythema, tissue proliferation, fever and pain can all be inhibited in laboratory animals by the administration of piroxicam. It is effective regardless of the aetiology of the inflammation. While its mode of action is not fully understood, independent studies *in vitro* as well as *in vivo* have shown that piroxicam interacts at several steps in the immune and inflammation responses through:

**Module 1**  
**PIRICAM 20**

**(Piroxicam Capsules USP 20mg)**

Inhibition of prostanoid synthesis, including prostaglandins, through a reversible inhibition of the cyclo-oxygenase enzyme.

Inhibition of neutrophil aggregation.

Inhibition of polymorphonuclear cell and monocyte migration to the area of inflammation.

Inhibition of lysosomal enzyme release from stimulated leucocytes.

Reduction of both systemic and synovial fluid rheumatoid factor production in patients with seropositive rheumatoid arthritis.

It is established that piroxicam does not act by pituitary-adrenal axis stimulation. In-vitro studies have not revealed any negative effects on cartilage metabolism.

**5.2 Pharmacokinetic properties**

Absorption:

Piroxicam is well absorbed following oral or rectal administration. With food there is a slight delay in the rate but not the extent of absorption following administration. The plasma half-life is approximately 50 hours in man and stable plasma concentrations are maintained throughout the day on once-daily dosage. Continuous treatment with 20mg/day for periods of 1 year produces similar blood levels to those seen once steady state is first achieved.

Distribution:

Drug plasma concentrations are proportional for 10 and 20mg doses and generally peak within 3 to 5 hours after medication. A single 20mg dose generally produces peak piroxicam plasma levels of 1.5 to 2 mcg/ml while maximum plasma concentrations, after repeated daily ingestion of 20mg piroxicam, usually stabilise at 3 to 8 mcg/ml. Most patients approximate steady state plasma levels within 7 to 12 days.

Treatment with a loading dose regimen of 40mg daily for the first 2 days followed by 20mg daily thereafter allows a high percentage (approximately 76%) of steady state levels to be achieved immediately following the second dose. Steady state levels, area under the curves and elimination half-life are similar to that following a 20mg daily dose regimen.

A multiple dose comparative study of the bioavailability of the injectable forms with the oral capsule has shown that after intramuscular administration of piroxicam, plasma levels are significantly higher than those obtained after ingestion of capsules during the 45 minutes following administration the first day, during 30 minutes the second day and 15 minutes the seventh day. Bioequivalence exists between the two dosage forms.

A multiple dose comparative study of the pharmacokinetics and the bioavailability of Piroxicam FDDF with the oral capsule has shown that after once daily administration for 14 days, the mean plasma piroxicam concentration time profiles for capsules and Piroxicam FDDF were nearly superimposable. There were no significant differences between the mean steady state  $C_{max}$  values,  $C_{min}$  values,  $T_{1/2}$ , or  $T_{max}$  values. This study concluded that Piroxicam FDDF (Fast Dissolving Dosage Form) is bioequivalent to the capsule after once daily dosing. Single dose studies have demonstrated bioequivalence as well when the tablet is taken with or without water.

Biotransformation:

Piroxicam is extensively metabolised and less than 5% of the daily dose is excreted unchanged in urine and faeces. Piroxicam metabolism is predominantly mediated via cytochrome P450 CYP 2C9 in the liver. One important metabolic pathway is hydroxylation

## Module 1

### PIRICAM 20

#### (Piroxicam Capsules USP 20mg)

of the pyridyl ring of the piroxicam side-chain, followed by conjugation with glucuronic acid and urinary elimination.



MMC Healthcare Ltd.  
No.: 34-B, SIDCO Industrial Estate,  
Thirumazhisai, Chennai-600 124, India.

Patients who are known or suspected to be poor CYP2C9 metabolizers based on previous history/experience with other CYP2C9 substrates should be administered piroxicam with caution as they may have abnormally high plasma levels due to reduced metabolic clearance (see section 4.4).

#### Pharmacogenetics:

CYP2C9 activity is reduced in individuals with genetic polymorphisms, such as the CYP2C9\*2 and CYP2C9\*3 polymorphisms. Limited data from two published reports showed that subjects with heterozygous CYP2C9\*1/\*2 (n=9), heterozygous CYP2C9\*1/\*3 (n=9), and homozygous CYP2C9\*3/\*3 (n=1) genotypes showed 1.7-, 1.7-, and 5.3-fold higher piroxicam systemic levels, respectively, than the subjects with CYP2C9\*1/\*1 (n=17, normal metabolizer genotype) following administration of an oral single dose. The mean elimination half life values of piroxicam for subjects with CYP2C9\*1/\*3 (n=9) and CYP2C9\*3/\*3 (n=1) genotypes were 1.7- and 8.8-fold higher than subjects with CYP2C9\*1/\*1 (n=17). It is estimated that the frequency of the homozygous\*3/\*3 genotype is 0% to 5.7% in various ethnic groups.

### 5.3 Preclinical safety data

Not applicable.

## 6. Pharmaceutical particulars

### 6.1 List of excipients

1. Lactose Monohydrate USP
2. Povidone BP
3. Microcrystalline cellulose BP
4. Sodium lauryl sulfate BP
5. Croscarmellose sodium BP
6. Magnesium stearate BP
7. Colloidal anhydrous silica BP
8. Purified Talc BP

### 6.2 Incompatibilities

Not applicable

### 6.3 Shelf life

24 months from the date of Manufacturing

### 6.4 Special precautions for storage

Store below 30°C. Protect from light.

Keep out of reach of children

### 6.5 Nature and contents of container

White colour powder filled in Dark red / Deep red colour '2' size Hard Gelatin capsules. 10 Capsules packed in a Alu/ Alu blister pack and 10 such blisters are packed in a carton with pack insert. Such cartons packed in a 7 ply corrugated box.

### 6.6 Special precautions for disposal and other handling

No special requirements.

**Module 1**

**PIRICAM 20**

**(Piroxicam Capsules USP 20mg)**

**7. Marketing authorization holder**

**Cadila Healthcare Limited,**

Zydus Corporate Park, Scheme No.63,  
Survey No. 536, Khoraj (Gandhi Nagar)  
Nr. Vaishnodevi Circle,Ahmedabad,  
Gandhinagar GJ,382481, India



**MMC Healthcare Ltd.**

No.: 34-B, SIDCO Industrial Estate,  
Thirumazhisai, Chennai-600 124.India.

**8. Marketing authorization number(s)**

**9. Date of first authorization/renewal of the authorization**

**10. Date of revision of the text - Apr 2020**

**Reference:** electronic medicines compendium (emc)