CTD MODULE 1 ADMINISTRATIVE INFORMATION AND PRODUCT INFORMATION

Product Name: RECODIN SYRUP

(Chlorpheniramine Maleate 4mg and Codeine Phosphate 10mg)

Pack Insert

RECODIN

Each 5ml contains:

Codeine Phosphate BP 10mg

Chlorpheniramine Maleate BP 4mg

INDICATIONS

Recodin Syrup is specially formulated to provide relief from dry irritating cough and Flu associated with coughs (Painful coughs)

DOSE

Recodin syrup is to be taken as follows or as instructed by the Physician Adults: Take 5ml six hourly (Four times a day)

Children

Under 6 years: As directed by the Physician 6-12 years, 2.5ml three times a day (Eight hourly)

Contraindications & Caution.

Recodin syrup should not be used by patients with hypersensitivity to any of the ingredients. Do not drive or operate heavy machinery when dizziness occurs. Caffeine is an addictive drug hence should be taken under prescription only

Drug interaction

Do not take Codeine with other Narcotic pain medications, sedatives, tranquilizers, muscle relaxants or other medicines that can make you sleepy or slow your breathing. Avoid alcohol while taking Recodin syrup.

Side Effects

Dizziness, Nausea, Constipation, CNS stimulation, Euphoria

OVERDOSE

Overdose symptoms may include extreme drowsiness, pin point pupils, confusion, cold and clammy skin, weak pulse, shallow breathing, fainting or breathing that stops.

In case of this, perform Gastric lavage.

STORAGE

Store in a cool dry place bellow 30°C away from light Keep out of reach of children

MANUFACTURED BY

RENE INDUSTRIES LIMITED Plot 680 Kamuli, Kireka, P.O. Box 6034, Kampala, Uganda A Milestone to Golden Health