



**AGOG Pharma Ltd.**

(WHO - GMP CERTIFIED - GOVT RECOGNISED EXPORT HOUSE)



CIN: 11-06518

Regd. Office & Factory : Plot No. 33, Sector II, The Vasai Taluka Industrial Co-op. Estate Ltd. Gauripada, Vasai (E), Dist. Thane - 401 208. INDIA.  
Tel. : 95250 - 2455801 / 2452714 / 2453525 • Fax: 95250 - 2452074 (0091 - 250 - 2452074) • Email : agog@vsnl.net & agogpharma@rediffmail.com

<b>Brand Name</b> : CLOM TABLETS	2021
<b>Generic Name</b> : Clomipramine Tablets 25 mg	
<b>Module 1</b> Administrative Information and Product Information	<b>Confidential</b>
<b>1.5</b> Product Information	

## 1.5 PRODUCT INFORMATION

### 1.5.1 Prescribing information (Summary of products characteristics)

#### SUMMARY PRODUCT CHARACTERISTICS

#### 1. Name of drug product:

CLOM TABLETS (Clomipramine Tablets 25 mg)

#### 2. Qualitative and Quantitative Composition:

Each film coated tablet contains: Clomipramine Hydrochloride BP 25 mg

#### 3. Pharmaceutical form:

Yellow coloured, circular, biconvex, film coated tablet.

#### 4. Clinical particulars:

##### 4.1 Therapeutic Indications:

##### Adults

Clomipramine Tablets 25 mg are indicated for the treatment of:

- the symptoms of depressive illness especially where sedation is required
- obsessional and phobic states
- adjunctive treatment of cataplexy associated with narcolepsy.

##### Children and adolescents

In children and adolescents, there is not sufficient evidence of safety and efficacy of clomipramine in the treatment of depressive states, phobias and cataplexy associated with narcolepsy. The use of Clomipramine in children and adolescents (0-17 years of age) in these indications is therefore not recommended.



**AGOG Pharma Ltd.**

(WHO - GMP CERTIFIED - GOVT RECOGNISED EXPORT HOUSE)



QMS 11-005110

Regd. Office & Factory : Plot No. 33, Sector II, The Vasai Taluka Industrial Co-op. Estate Ltd. Gauraipada, Vasai (E), Dist. Thane - 401 208. INDIA.  
Tel. : 95250 - 2455801 / 2452714 / 2453525 • Fax : 95250 - 2452074 (0091 - 250 - 2452074) • Email : agog@vsni.net & agogpharma@rediffmail.com

## 4.2 Posology and Method of Administration:

### Posology

Before initiating treatment with Clomipramine, hypokalemia should be treated

After a response has been obtained, maintenance therapy should be continued at the optimum dose to avoid relapse. Patients with a history of recurrence require maintenance treatment for a longer duration. Duration of maintenance treatment and need for further treatment should be reviewed periodically.

As a precaution against possible QTc prolongation and serotonergic toxicity, adherence to the recommended doses of clomipramine is advised and any increase in dose should be made with caution if drugs that prolong QT interval or other serotonergic agents are co-administered.

Abrupt discontinuation of Clomipramine therapy should be avoided because of possible withdrawal symptoms. Therefore, dosage should be stopped gradually after regular use for long duration and the patient should be monitored carefully when Clomipramine therapy is discontinued.

### *Adults*

Oral – 10 mg/day initially, increasing gradually to 30-150 mg/day, if required, in divided doses throughout the day or as a single dose at bedtime. Many patients will be adequately maintained on 30-50 mg/day. Higher doses may be needed in some patients, particularly those suffering from obsessional or phobic disorders. In severe cases this dosage can be increased up to a maximum of 250 mg per day. Once a distinct improvement has set in, the daily dosage may be adjusted to a maintenance level of about 50-100 mg.

### *Elderly*

Elderly patients generally show a stronger response to Clomipramine than patients of intermediate age groups, Clomipramine should be used with caution in elderly patients and doses should be increased cautiously. The initial dose should be 10 mg/day, which may be increased with caution under close supervision to an optimum level of 30-75 mg daily which should be reached after about 10 days and then maintained until the end of treatment.

### *Paediatric population*

Not recommended.

### *Obsessional/phobic states*

The maintenance dosage of Clomipramine is generally higher than that used in depression. It is recommended that the dose be built up to 100-150 mg Clomipramine daily, according to the severity of the condition. This should be attained gradually over a period of 2 weeks starting with 1 x 25 mg Clomipramine daily. In elderly patients and those sensitive to tricyclic antidepressants a starting dose of 1 x 10 mg Clomipramine daily is recommended. Again where a higher dosage is required the sustained-release 75 mg formulation may be preferable.

### *Adjunctive treatment of cataplexy associated with narcolepsy*

(Oral treatment): 10-75 mg daily. It is suggested that treatment is commenced with 10 mg Clomipramine daily and gradually increased until a satisfactory response occurs. Control of cataplexy should be achieved within 24 hours of reaching the optimal dose. Where necessary, therapy may be combined with capsules up to the maximum dose of 75 mg per day.

### ***Treatment discontinuation***

Abrupt withdrawal should be avoided because of possible adverse reactions. If the decision is made to discontinue treatment, medication should be tapered, as rapidly as is feasible, but with recognition that abrupt discontinuation can be associated with certain symptoms.

### ***Renal impairment***

Clomipramine should be given with caution in patients with renal impairment.

### ***Hepatic impairment***

Clomipramine should be given with caution in patients with hepatic impairment

### **Method of administration**

For oral use

## **4.3 Contraindications:**

- Hypersensitivity to the active substance, or to any of the excipients listed in section 6.1 or cross-sensitivity to tricyclic antidepressants of the dibenzazepine group
- Recent myocardial infarction, any degree of heart block or other cardiac arrhythmias
- Severe liver disease
- Concurrent administration with monoamine oxidase inhibitors or within 3 weeks of start or cessation of therapy.
- Concomitant treatment with selective, reversible MAO-A inhibitors, such as moclobemide
- Narrow-angle glaucoma
- Retention of urine

## **4.4 Special warnings and precautions for use**

### **Suicide/suicidal thoughts or clinical worsening**

Depression is associated with an increased risk of suicidal thoughts, self-harm and suicide (suicide-related events). This risk persists until significant remission occurs. As improvement may not occur during the first few weeks or more of treatment, patients should be closely monitored until such improvement occurs. It is general clinical experience that the risk of suicide may increase in the early stages of recovery.

Other psychiatric conditions for which Clomipramine is prescribed can also be associated with an increased risk of suicide-related events. In addition, these conditions may be comorbid with major depressive disorder. The same precautions observed when treating patients with major depressive disorder should therefore be observed when treating patients with other psychiatric disorders.

Patients with a history of suicide-related events, or those exhibiting a significant degree of suicidal ideation prior to commencement of treatment are known to be at greater risk of suicidal thoughts or suicide attempts, and should receive careful monitoring during treatment. A meta-analysis of placebo-controlled clinical trials of antidepressant drugs in adult patients with psychiatric disorders showed an increased risk of suicidal behavior with antidepressants compared to placebo in patients less than 25 years old.



**AGOG Pharma Ltd.**

(WHO - GMP CERTIFIED - GOVT RECOGNISED EXPORT HOUSE)



CIN: 11-069118

Regd. Office & Factory : Plot No. 33, Sector II, The Vasai Taluka Industrial Co-op. Estate Ltd. Gauripada, Vasai (E), Dist. Thane - 401 208. INDIA.  
Tel. : 95250 - 2455801 / 2452714 / 2453525 • Fax : 95250 - 2452074 (0091 - 250 - 2452074) • Email : agog@vsni.net & agogpharma@rediffmail.com

Close supervision of patients and in particular those at high risk should accompany drug therapy especially in early treatment and following dose changes. Patients (and caregivers of patients) should be alerted about the need to monitor for any clinical worsening, suicidal behaviour or thoughts and unusual changes in behaviour and to seek medical advice immediately if these symptoms present.

Patients with depressive disorders, both adult and paediatric, may experience worsening of depression and/or suicidality or other psychiatric symptoms, whether or not they are taking antidepressant medication.

### **Paediatric population**

Clomipramine should not be used in the treatment of depressive states, phobias and cataplexy associated with narcolepsy in children and adolescents under the age of 18 years.

Antidepressants increase the risk of suicide-related behaviors (suicide attempt and suicidal thoughts), and hostility (predominately aggression, oppositional behavior and anger) were more frequently observed in clinical trials among children and adolescents treated with antidepressants compared to those treated with placebo. If based on clinical need, a decision to treat is nevertheless taken, the patient should be carefully monitored for the appearance of suicidal symptoms. In addition, long term safety data in children and adolescents concerning growth, maturation and cognitive behavioral development are lacking.

Families and care givers of both pediatric and adult patients being treated with antidepressants for both psychiatric and non-psychiatric indications, should be alerted about the need to monitor patients for the emergence of other psychiatric symptoms, as well as the emergence of suicidality, and to report such symptoms immediately to health care providers.

Prescriptions for Clomipramine should be written for the smallest quantity of capsules consistent with good patient management, in order to reduce the risk of overdose.

Modifying the therapeutic regimen, including possibly discontinuing the medication, should be considered in these patients, especially if these changes are severe, abrupt in onset, or were not part of the patient's presenting symptoms.

### **Other psychiatric effects**

Many patients with panic disorders experience intensified anxiety symptoms at the start of the treatment with antidepressants. This paradoxical initial increase in anxiety is most pronounced during the first few days of treatment and generally subsides within two weeks.

Activation of psychosis has occasionally been observed in patients with schizophrenia receiving tricyclic antidepressants.

Hypomanic or manic episodes have also been reported during a depressive phase in patients with cyclic affective disorders receiving treatment with a tricyclic antidepressant. In such cases it may be necessary to reduce the dosage of clomipramine or to withdraw it and administer an antipsychotic agent. After such episodes have subsided, low dose therapy with Clomipramine may be resumed if required.

In predisposed patients, tricyclic antidepressants may provoke pharmacogenic (delirious) psychoses, particularly at night. These disappear within a few days of withdrawing the drug.

As improvement in depression may not occur for the first two to four weeks treatment, patients should be closely monitored during this period.

Elderly patients are particularly liable to experience adverse effects, especially agitation, confusion, and postural hypotension.



**AGOG Pharma Ltd.**

(WHO - GMP CERTIFIED - GOVT RECOGNISED EXPORT HOUSE)



CPA 11-06118

Regd. Office & Factory : Plot No. 33, Sector II, The Vasai Taluka Industrial Co-op. Estate Ltd. Gauripada, Vasai (E), Dist. Thane - 401 208, INDIA.  
Tel. : 95250 - 2455801 / 2452714 / 2453525 • Fax : 95250 - 2452074 (0091 - 250 - 2452074) • Email : agog@vsni.net & agogpharma@rediffmail.com

### **Cardiac and vascular disorders**

Clomipramine should be administered with particular precaution in patients with cardiovascular disorders, especially those with cardiovascular insufficiency, conduction disorders, (*e.g.* atrioventricular block grades I to III), arrhythmias. Monitoring of cardiac function and the ECG is indicated in such patients.

There may be a risk of QTc prolongation and torsades de pointes, particularly at supra-therapeutic doses or supra-therapeutic plasma concentrations of Clomipramine, as occur in the case of co-medication with selective serotonin reuptake inhibitors (SSRIs). Therefore, concomitant administration of drugs that can cause accumulation of clomipramine Hcl should be avoided. Equally, concomitant administration of drugs that can prolong the QTc interval should be avoided. It is established that hypokalaemia is a risk-factor of QTc prolongation and torsades de pointes. Therefore, hypokalaemia should be treated before initiating treatment with Clomipramine.

Before initiating treatment it is advisable to check the patient's blood pressure, because individuals with hypotension or a labile circulation may react to the drug with a fall in blood pressure.

### **Serotonin syndrome**

Concomitant administration of Clomipramine and SSRIs, SNaRIs, tricyclic antidepressants, lithium and buprenorphine/ opioids may result in serotonin syndrome, a potentially life-threatening condition.

If concomitant treatment with other serotonergic agents is clinically warranted, careful observation of the patient is advised, particularly during treatment initiation and dose increases.

Symptoms of serotonin syndrome may include mental-status changes (delirium), autonomic instability (agitation, seizures, coma), neuromuscular abnormalities (myoclonus), and/or gastrointestinal symptoms.

If serotonin syndrome is suspected, a dose reduction or discontinuation of therapy should be considered depending on the severity of the symptoms.

### **Convulsions**

Tricyclic antidepressants are known to lower the convulsion threshold and Clomipramine should therefore, be used with extreme caution in patients with epilepsy and other predisposing factors, *e.g.* brain damage of varying aetiology, concomitant use of neuroleptics, withdrawal from alcohol or drugs with anticonvulsive properties (*e.g.* benzodiazepines). It appears that the occurrence of seizures is dose dependent. Therefore, the recommended total daily dose of Clomipramine should not be exceeded.

Concomitant treatment of Clomipramine and electroconvulsive therapy should only be resorted to under careful supervision.

### **Anticholinergic effects**

Because of its anticholinergic properties, Clomipramine should be used with caution in patients with a history of increased intra-ocular pressure, narrow-angle glaucoma, urinary retention or with symptoms of bladder neck obstruction, *e.g.* diseases of the prostate, such as prostatic hypertrophy.

Decreased lacrimation and accumulation of mucoid secretions due to the anticholinergic properties of tricyclic antidepressants may cause damage to the corneal epithelium in patients with contact lenses.



**AGOG Pharma Ltd.**

(WHO - GMP CERTIFIED - GOVT RECOGNISED EXPORT HOUSE)



CIN: 11-06918

Regd. Office & Factory : Plot No. 33, Sector II, The Vasai Taluka Industrial Co-op. Estate Ltd. Gauraipada, Vasai (E), Dist. Thane - 401 208, INDIA.  
Tel. : 95250 - 2455801 / 2452714 / 2453525 • Fax : 95250 - 2452074 (0091 - 250 - 2452074) • Email : agog@vsnl.net & agogpharma@rediffmail.com

### **Specific treatment populations**

Caution is called for when giving tricyclic antidepressants to patients with severe hepatic disease and tumours of the adrenal medulla (e.g. pheochromocytoma, neuroblastoma), in whom they may provoke hypertensive crises.

Caution is advised in patients with hyperthyroidism or during concomitant treatment with thyroid preparations since aggravation of unwanted cardiac effects may occur.

It is advisable to monitor cardiac and hepatic function during long-term therapy with clomipramine. In patients with hepatic and renal disease, periodic monitoring of hepatic enzyme levels and renal function is recommended.

An increase in dental caries has been reported during long-term treatment with tricyclic antidepressants. Regular dental check-ups are therefore advisable during long-term treatment.

Caution is called for in patients with chronic constipation. Tricyclic antidepressants may cause paralytic ileus, particularly in the elderly and in bed-ridden patients.

In elderly patients, tricyclic antidepressants may provoke pharmacogenic (delirious) psychoses, particularly at night. These disappear within a few days of withdrawing the drug.

Monitoring of cardiac function and the ECG is indicated in elderly patients.

### **White blood cell count**

Although changes in the white blood cell count have been reported with Clomipramine only in isolated cases, periodic blood cell counts and monitoring for symptoms such as fever and sore throat are called for, particularly during the first few months of therapy. They are also recommended during prolonged therapy.

### **Anticoagulants / Non-steroidal anti-inflammatory drugs**

Skin and mucous membrane bleeding has been reported with Clomipramine. The product should be used with caution among patients that simultaneously use medicines that increase the risk of bleeding, for example anticoagulants, salicylic acid derivatives and non-steroidal anti-inflammatory drugs (NSAIDs). Care should be taken in patients with an increased tendency to bleed.

### **Anaesthesia**

Anaesthetics given during tri/tetracyclic antidepressant therapy may increase the risk of arrhythmias and hypotension.

Before general or local anaesthesia, the anaesthetist should be aware that the patient has been receiving Clomipramine and of the possible interactions.

### **Treatment discontinuation**

Abrupt withdrawal should be avoided because of possible adverse reactions. If the decision is made to discontinue treatment, medication should be tapered, as rapidly as is feasible, but with recognition that abrupt discontinuation can be associated with certain symptoms.

Clomipramine capsules contain lactose. Patients with rare hereditary problems of galactose intolerance, the Lapp lactase deficiency or glucose-galactose malabsorption should not take this medicine.



**AGOG Pharma Ltd.**

(WHO - GMP CERTIFIED - GOVT RECOGNISED EXPORT HOUSE)



CPA: 11-06518

Regd. Office & Factory : Plot No. 33, Sector II, The Vasai Taluka Industrial Co-op. Estate Ltd. Gauripada, Vasai (E), Dist. Thane - 401 208, INDIA.  
Tel. : 95250 - 2455801 / 2452714 / 2453525 • Fax : 95250 - 2452074 (0091 - 250 - 2452074) • Email : agog@vsnl.net & agogpharma@rediffmail.com

## 4.5 Interaction with other medicinal products and other forms of interaction

### **Interactions resulting in a contraindication**

#### **MAO inhibitors**

Do not give Clomipramine for at least 3 weeks after discontinuation of treatment with MAO inhibitors (there is a risk of severe symptoms consistent with Serotonin Syndrome such as hypertensive crisis, hyperpyrexia, myoclonus, agitation, seizures, delirium and coma). The same applies when giving a MAO inhibitor after previous treatment with clomipramine. In both instances the treatment should initially be given in small gradually increasing doses and its effects monitored. There is evidence to suggest that Clomipramine may be given as little as 24 hours after a reversible MAO-A inhibitor such as moclobemide, but the 3-week wash-out period must be observed if the MAO-A inhibitor is used after Clomipramine.

### **Interactions resulting in a concomitant use not recommended**

#### **Diuretics**

Diuretics may lead to hypokalemia, which increases the risk of QTc prolongation and torsade's de pointes. Hypokalemia should therefore be treated prior to administration of clomipramine.

#### **Quinidine**

Tricyclic antidepressants should not be employed in combination with antiarrhythmic agents of the quinidine type.

#### **Selective serotonin reuptake inhibitors (SSRIs)**

SSRIs which are inhibitors of CYP2D6, such as fluoxetine, paroxetine, or sertraline, and of others including CYP1A2 and CYP2C19 (e.g. fluvoxamine), may also increase plasma concentrations of clomipramine, with corresponding adverse effects. Steady-state serum levels of clomipramine increased ~4-fold by co-administration of fluvoxamine (N-desmethylclomipramine decreased ~2-fold). In addition, co-medication with SSRIs may lead to additive effects on the serotonergic system.

#### **Serotonergic agents**

Serotonin Syndrome can possibly occur when Clomipramine is administered with other serotonergic co-medications such as selective serotonin reuptake inhibitors (SSRIs), serotonin and noradrenergic reuptake inhibitors (SNARIs), tricyclic antidepressants and lithium. For fluoxetine, a wash-out period of two to three weeks is advised before and after treatment with fluoxetine.

### **Interactions to be considered**

#### **Interactions resulting in increased effect of clomipramine**

##### **Oral antifungal, terbinafine**

Co-administration of Clomipramine with terbinafine, a strong inhibitor of CYP2D6, may result in increased exposure and accumulation of Clomipramine and its N-demethylated metabolite. Therefore, dose adjustments of Clomipramine may be necessary when co-administered with terbinafine.



**AGOG Pharma Ltd.**

(WHO - GMP CERTIFIED - GOVT RECOGNISED EXPORT HOUSE)



CPA 11-06918

Regd. Office & Factory : Plot No. 33, Sector II, The Vasai Taluka Industrial Co-op. Estate Ltd. Gauripada, Vasai (E), Dist. Thane - 401 208, INDIA.  
Tel. : 95250 - 2455801 / 2452714 / 2453525 • Fax : 95250 - 2452074 (0091 - 250 - 2452074) • Email : agog@vsnl.net & agogpharma@rediffmail.com

### **Cimetidine**

Co-administration with the histamine<sub>2</sub> (H<sub>2</sub>)-receptor antagonist, cimetidine (an inhibitor of several P450 enzymes, including CYP2D6 and CYP3A4), may increase plasma concentrations of tricyclic antidepressants, whose dosage should therefore be reduced.

### **Oral contraceptives**

No interaction between chronic oral contraceptive use (15 or 30 micrograms ethinylestradiol daily) and Clomipramine (25 mg daily) has been documented. Oestrogens are not known to be inhibitors of CYP2D6, the major enzyme involved in Clomipramine clearance and, therefore, no interaction is expected. Although, in a few cases with high dose oestrogen (50 micrograms daily) and the tricyclic antidepressant imipramine, increased side effects and therapeutic response were noted, it is unclear as to the relevance of these cases to Clomipramine and lower dose estrogen regimens. Monitoring therapeutic response of tricyclic antidepressants at high dose estrogen regimens (50 micrograms daily) is recommended and dose adjustments may be necessary.

### **Antipsychotics**

Co-medication of antipsychotic (*e.g.* phenothiazines) may result in increased plasma levels of tricyclic antidepressants, a lowered convulsion threshold, and seizures. Combination with thioridazine may produce severe cardiac arrhythmias.

### **Methylphenidate**

This drug may also increase plasma concentrations of tricyclic antidepressants, whose dosage should therefore be reduced.

### **Valproate**

Concomitant administration of valproate with Clomipramine may cause inhibition of CYP2C and/or UGT enzymes resulting in increased serum levels of Clomipramine and desmethylclomipramine.

### **Grapefruit, grapefruit juice, or cranberry juice**

Concomitant administration of Clomipramine with grapefruit, grapefruit juice, or cranberry juice may increase the plasma concentrations of clomipramine.

### **Interactions resulting in decreased effect of clomipramine**

#### **Rifampicin**

Rifampicin (CYP3A and CYP2C inducer), may decrease Clomipramine concentrations as concomitant administration of drugs known to induce cytochrome P450 enzymes, particularly CYP3A4, CYP2C19 may accelerate the metabolism and decrease the efficacy of Clomipramine.

#### **Anticonvulsants**

Anticonvulsants (CYP3A and CYP2C inducer) *e.g.* barbiturates, carbamazepine, phenobarbital and phenytoin, may decrease Clomipramine concentrations as concomitant administration of drugs known to induce cytochrome P450 enzymes, particularly CYP3A4, CYP2C19 may accelerate the metabolism and decrease the efficacy of Clomipramine.





**AGOG Pharma Ltd.**

(WHO - GMP CERTIFIED - GOVT RECOGNISED EXPORT HOUSE)



CPA 11-06918

Regd. Office & Factory : Plot No. 33, Sector II, The Vasai Taluka Industrial Co-op. Estate Ltd. Gauripada, Vasai (E), Dist. Thane - 401 208, INDIA.  
Tel. : 95250 - 2455801 / 2452714 / 2453525 • Fax : 95250 - 2452074 (0091 - 250 - 2452074) • Email : agog@vsnl.net & agogpharma@rediffmail.com

### **Cigarette smoking**

Known inducers of CYP1A2 (*e.g.* nicotine/components in cigarette smoke), decrease plasma concentrations of tricyclic drugs. In cigarette smokers, Clomipramine steady-state plasma concentrations were decreased 2-fold compared to non-smokers (no change in *N*-desmethyldomipramine).

### **Colestipol and cholestyramine**

Concomitant administration of ion exchange resins such as cholestyramine or colestipol may reduce the plasma levels of Clomipramine. Staggering the dosage of Clomipramine and resins, such that the drug is administered at least 2 h before or 4-6 h after the administration of resins, is recommended.

### **St. John's wort**

Concomitant administration of Clomipramine with St. John's wort during the treatment may decrease the plasma concentrations of clomipramine.

### **Interactions affecting other drugs**

#### **Anticholinergic agents**

Tricyclic antidepressants may potentiate the effects of these drugs (*e.g.* phenothiazines, antiparkinsonian agents, antihistamines, atropine, biperiden) on the eye, central nervous system, bowel and bladder).

#### **Antiadrenergic agents**

Clomipramine may diminish or abolish the antihypertensive effects of guanethidine, betanidine, reserpine, clonidine and alpha-methyldopa. Patients requiring co-medication for hypertension should therefore be given antihypertensives of a different type (*e.g.* vasodilators, or beta-blockers).

It would be advisable to review all antihypertensive therapy during treatment with tricyclic antidepressants.

#### **CNS depressants**

Tricyclic antidepressants may potentiate the effects of alcohol and other central depressant substances (*e.g.* barbiturates, benzodiazepines, or general anaesthetics).

#### **Sympathomimetic drugs**

Clomipramine may potentiate the cardiovascular effects of adrenaline, ephedrine, isoprenaline, noradrenaline, phenylephrine, and phenylpropanolamine (*e.g.* as contained in local and general anaesthetic preparations and nasal decongestants).

#### **Anticoagulants**

Tricyclic antidepressants may potentiate the anticoagulant effect of coumarin drugs by inhibiting their metabolism by the liver. Careful monitoring of plasma prothrombin is therefore advised.

#### **Drugs that can cause increase plasma clomipramine levels or which in themselves prolong the QTc interval**

The risk of QTc prolongation and torsades de pointes is likely to be increased if Clomipramine is co-administered with other drugs that can cause QTc prolongation. Therefore concomitant use of such agents with Clomipramine is not recommended. Examples include certain antiarrhythmic, such as those of Class 1A (such as quinidine, disopyramide



and procainamide) and Class III (such as amiodarone and sotalol), tricyclic antidepressants (such as amitriptyline), certain tetracyclic antidepressants (such as maprotiline), certain antipsychotic medications (such as phenothiazines and pimozide), certain antihistamines (such as terfenadine); lithium, quinine and pentamidine. This list is not exhaustive. The risk of QTc prolongation and torsades de pointes is likely to be increased if Clomipramine is co-administered with drugs that can cause increased plasma clomipramine levels. Clomipramine is metabolised by cytochrome P450 2D6 and the plasma concentration of Clomipramine may therefore be increased by drugs that are either substrates and/or inhibitors of this P450 isoform. Therefore, concurrent use of these drugs with Clomipramine is not recommended. Examples of drugs which are substrates or inhibitors of cytochrome P450 2D6 include antiarrhythmics, certain antidepressants including SSRIs, tricyclic antidepressants and moclobemide; certain antipsychotics;  $\beta$ -blockers; protease inhibitors, opiates, ecstasy (MDMA), cimetidine and terbinafine. This list is not exhaustive.

#### Calcium channel blockers

Diltiazem and verapamil may increase the plasma concentration of imipramine, and possibly other tricyclics.

#### Non-steroidal anti-inflammatory drugs

The potential pharmacodynamic interactions with drugs that increase the risk of bleeding, for example, salicylic acid derivatives, non-steroidal anti-inflammatory / antirheumatic drugs (NSAIDs) should be considered because of the increased risk of bleeding with concomitant clomipramine.

#### Cytotoxic

**Altretamine:** risk of severe postural hypotension.

#### Analgesics

Possible increased side effects with nefopam; possible increased risk of convulsions with tramadol; possible increased sedation with opioid analgesics; increased risk of ventricular arrhythmias with levacetylmethadol.

#### Antipsychotics

Increased risk of ventricular arrhythmias – avoid concomitant use with pimozide. Antipsychotic agents may increase the plasma concentration of tricyclic agents. No such effects are known to occur in combination with diazepam, but it might be necessary to reduce the dosage of Clomipramine if administered concomitantly with alprazolam or disulfiram. There may be increased antimuscarinic side-effects with phenothiazines, and possibly clozapine.

#### Dopaminergic

Concomitant use of tricyclic antidepressants and entacapone should be avoided; central nervous system toxicity has been reported with selegiline.

#### Muscle relaxants

Tricyclic antidepressants may enhance the muscle relaxant effect of baclofen.

Clomipramine should be used cautiously when co-administered with:

- **Buprenorphine/ opioids as the risk of serotonin syndrome, a potentially life-threatening condition is increased**



**AGOG Pharma Ltd.**

(WHO - GMP CERTIFIED - GOVT RECOGNISED EXPORT HOUSE)



Regd. Office & Factory : Plot No. 33, Sector II, The Vasai Taluka Industrial Co-op. Estate Ltd. Gauripada, Vasai (E), Dist. Thane - 401 208. INDIA.  
Tel. : 95250 - 2455801 / 2452714 / 2453525 • Fax : 95250 - 2452074 (0091 - 250 - 2452074) • Email : agog@vsnl.net & agogpharma@rediffmail.com

## 4.6 Fertility, pregnancy and lactation

### Women of child-bearing potential

There are no data supporting any special recommendations in women of child-bearing potential.

### Pregnancy

There is limited amount of data from the use of Clomipramine in pregnant women that indicates a potential to harm the foetus or cause congenital malformation.

Neonates whose mothers had taken tricyclic antidepressants up until delivery have developed dyspnoea, lethargy, colic, irritability, hypotension or hypertension, tremor or spasms, during the first few hours or days.

Studies in animals have shown reproductive toxicity. Clomipramine is not recommended during pregnancy and in women of childbearing potential not using contraception.

### Breast-feeding

Clomipramine passes into the breast milk in small quantities. Therefore nursing mothers should be advised to withdraw the medication or cease breast-feeding.

### Fertility

Clomipramine hydrochloride did not appear to have any significant effects on fertility and general reproductive performance.

## 4.7 Effects on ability to drive and use machines

Patients receiving clomipramine should be warned that blurred vision, drowsiness and other nervous system and psychiatric related disorders such as somnolence, disturbance in attention, confusion, disorientation, aggravation of depression, delirium etc have been observed. In the presence of such effects, patients should not drive, operate machinery or do anything else which may require alertness or quick actions. Patients should also be warned that alcohol or other drugs may potentiate these effects.

This medicine can impair cognitive function and can affect a patient's ability to drive safely. This class of medicine is in the list of drugs included in regulations under 5a of the Road Traffic Act 1988. When prescribing this medicine, patients should be told:

- The medicine is likely to affect your ability to drive
- Do not drive until you know how the medicine affects you
- It is an offence to drive while under the influence of this medicine
- However, you would not be committing an offence (called 'statutory defence') if:
  - The medicine has been prescribed to treat a medical or dental problem and
  - You have taken it according to the instructions given by the prescriber and in the information provided with the medicine and
  - It was not affecting your ability to drive safely



### 4.8 Undesirable effects

Unwanted effects are usually mild and transient, disappearing under continued treatment or with a reduction in the dosage. They do not always correlate with plasma drug levels or dose. It is often difficult to distinguish certain undesirable effects from symptoms of depression such as fatigue, sleep disturbances, agitation, anxiety, constipation, and dry mouth.

If severe neurological or psychiatric reactions occur, clomipramine should be withdrawn.

Adverse reactions are ranked under heading of frequency, the most frequent first, using the following convention: very common ( $\geq 1/10$ ); common ( $\geq 1/100$  to  $< 1/10$ ); uncommon ( $\geq 1/1,000$  to  $< 1/100$ ); rare ( $\geq 1/10,000$  to  $< 1/1,000$ ); very rare ( $< 1/10,000$ ); not known (cannot be estimated from the available data).

<b>Blood and lymphatic system disorders</b>	
Very rare	Leucopenia, agranulocytosis, thrombocytopenia, eosinophilia
<b>Immune system disorders</b>	
Very rare	Anaphylactic and anaphylactoid reactions including hypotension
<b>Endocrine disorders</b>	
Very rare	SIADH (inappropriate antidiuretic hormone secretion syndrome)
<b>Metabolism and nutrition disorders</b>	
Very common	Increased appetite
Common	Decreased appetite
<b>Psychiatric disorders</b>	
Very common	Restlessness
Common	Confusional state, disorientation, hallucinations (particularly in elderly patients and patients with Parkinson's disease), anxiety, agitation, sleep disorder, mania, hypomania, aggression, depersonalisation, aggravation of depression, insomnia, nightmares, delirium
Uncommon	Activation of psychotic symptoms
Not known	Suicidal ideation, suicidal behaviours <sup>2</sup>
<b>Nervous system disorders</b>	
Very common	Dizziness, tremor, headache, myoclonus, somnolence
Common	Speech disorder, paraesthesia, hypertonia, dysgeusia, memory impairment, disturbance in attention
Uncommon	Convulsions, ataxia
Very rare	Neuroleptic malignant syndrome <sup>1</sup>



Not known	Serotonin syndrome, extrapyramidal disorder (including akathisia and tardive dyskinesia) <sup>3</sup>
<b>Eye disorders</b>	
Very common	Accommodation disorder, vision blurred
Common	Mydriasis
Very rare	Glaucoma
<b>Ear and labyrinth disorders</b>	
Common	Tinnitus
<b>Cardiac disorders</b>	
Common	Sinus tachycardia, palpitation, orthostatic hypotension, clinically irrelevant ECG changes (e.g. ST and T changes) in patients of normal cardiac status
Uncommon	Arrhythmias, blood pressure increased
Very rare	Conduction disorder (e.g. widening of QRS complex, prolonged QT interval, PQ changes, bundle-branch block, torsades de pointes, particularly in patients with hypokalaemia)
<b>Vascular disorders</b>	
Common	Hot flush
<b>Respiratory, thoracic, and mediastinal disorders</b>	
Common	Yawning
Very rare	Alveolitis allergic (pneumonitis) with or without eosinophilia
<b>Gastrointestinal disorders</b>	
Very common	Nausea, dry mouth, constipation
Common	Vomiting, gastrointestinal disorders, diarrhoea
<b>Hepatobiliary disorders</b>	
Very rare	Hepatitis with or without jaundice
<b>Skin and subcutaneous tissue disorders</b>	
Very common	Hyperhidrosis
Common	Dermatitis allergic (skin rash, urticaria), photosensitivity reaction, pruritus
Very rare	Ecchymosis, purpura, alopecia
<b>Musculoskeletal and connective tissue disorders</b>	
Common	Muscular weakness
Not known	Rhabdomyolysis (as a complication of neuroleptic malignant syndrome) <sup>3</sup>



<b>Renal and urinary disorders</b>	
Very common	Micturition disorder
Common	Urinary retention
<b>Reproductive system and breast disorders</b>	
Very common	Libido disorder, erectile dysfunction
Common	Galactorrhoea, breast enlargement, women occasionally experience orgasmic impotence
Rare	Vaginal bleeding
Not known	Ejaculation failure, ejaculation delayed
<b>General disorders and administration site conditions</b>	
Very common	Fatigue
Very rare	Oedema (local or generalised), hyperpyrexia
<b>Investigations</b>	
Very common	Weight increased, blood sugar changes
Common	Transaminases increased
Very rare	Electroencephalogram abnormal
Not known	Blood prolactin increased <sup>3</sup>

- In post-marketing experience very rarely malignant neuroleptic syndrome has been reported although a causal relationship has not been confirmed.
- Cases of suicidal ideation and suicidal behaviors have been reported during clomipramine therapy or early after treatment discontinuation.
- These adverse events were reported in patients treated with clomipramine based on post marketing reports.

**Withdrawal symptoms**

The following symptoms commonly occur after abrupt withdrawal or reduction of the dose: nausea, vomiting, abdominal pain, diarrhoea, insomnia, headache, nervousness and anxiety (see section 4.4).

**Class effects**

Epidemiological studies, mainly conducted in patients 50 years of age and older, show an increased risk of bone fractures in patients receiving SSRIs and TCAs. The mechanism leading to this risk is unknown.

**Elderly population**

Elderly patients are particularly sensitive to anticholinergic, neurological, psychiatric, or cardiovascular effects. Their ability to metabolise and eliminate drugs may be reduced, leading to a risk of elevated plasma concentrations at therapeutic doses.



**AGOG Pharma Ltd.**

(WHO - GMP CERTIFIED - GOVT RECOGNISED EXPORT HOUSE)



024 11-86515

Regd. Office & Factory : Plot No. 33, Sector II, The Vasai Taluka Industrial Co-op. Estate Ltd. Gauripada, Vasai (E), Dist. Thane - 401 208. INDIA.  
Tel. : 95250 - 2455801 / 2452714 / 2453525 • Fax : 95250 - 2452074 (0091 - 250 - 2452074) • Email : agog@vsnl.net & agogpharma@rediffmail.com

### **Reporting of suspected adverse reactions**

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the Yellow Card Scheme Website: [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard)

### **4.9 Overdose**

The signs and symptoms of overdose with clomipramine are similar to those reported with other tricyclic antidepressants. Cardiac abnormalities and neurological disturbances are the main complications. In children accidental ingestion of any amount should be regarded as serious and potentially fatal.

#### **Signs and symptoms**

Symptoms generally appear within 4 hours of ingestion and reach maximum severity after 24 hours. Owing to delayed absorption (anticholinergic effect), long half-life, and enterohepatic recycling of the drug, the patient may be at risk for up to 4-6 days.

The following signs and symptoms may be seen:

#### **Central nervous system:**

Drowsiness, stupor, coma, ataxia, restlessness, agitation, enhanced reflexes, muscular rigidity, athetoid and choreoathetoid movements, convulsions, serotonin syndrome (e.g. hypertensive crisis, hyperpyrexia, myoclonus, delirium and coma).

#### **Cardiovascular system**

Hypotension, tachycardia, QTc prolongation and arrhythmia including torsades de pointes, conduction disorders, shock, heart failure; in very rare cases cardiac arrest.

Respiratory depression, cyanosis, vomiting, fever, mydriasis, sweating and oliguria or anuria may also occur.

#### **Treatment**

There is no specific antidote, and treatment is essentially symptomatic and supportive.

Anyone suspected of receiving an overdose of clomipramine, particularly children, should be hospitalized and kept under close surveillance for at least 72 hours.

Perform gastric lavage or induce vomiting as soon as possible if the patient is alert. If the patient has impaired consciousness, secure the airway with a cuffed endotracheal tube before beginning lavage, and do not induce vomiting. These measures are recommended for up to 12 hours or even longer after the overdose, since the anticholinergic effect of the drug may delay gastric emptying. Administration of activated charcoal may help to reduce drug absorption.

Treatment of symptoms is based on modern methods of intensive care, with continuous monitoring of cardiac function, blood gases, and electrolytes, and if necessary, emergency measures such as:



**AGOG Pharma Ltd.**

(WHO - GMP CERTIFIED - GOVT RECOGNISED EXPORT HOUSE)



Regd. Office & Factory : Plot No. 33, Sector II, The Vasai Taluka Industrial Co-op. Estate Ltd. Gauripada, Vasai (E), Dist. Thane - 401 208. INDIA.  
Tel. : 95250 - 2455801 / 2452714 / 2453525 • Fax : 95250 - 2452074 (0091 - 250 - 2452074) • Email : agog@vsnl.net & agogpharma@rediffmail.com

### **For respiratory failure:**

- Intubation and artificial respiration.

### **For cardiovascular symptoms:**

- In severe hypotension the patient should be placed in an appropriate position and be given a plasma expander, dopamine, or dobutamine by intravenous drip.
- Cardiac arrhythmias must be treated according to the requirements of the case.
- Implantation of a cardiac pacemaker should be considered.
- Low potassium values and acidosis should be corrected.

In all patients with ECG abnormalities, cardiac function should - even after the ECG tracings have reverted to normal - be kept under close observation for at least another 48 hours because relapses may occur.

### **Treatment of torsades de pointes:**

If torsades de pointes should occur during treatment with clomipramine, the drug should be discontinued and hypoxia, electrolyte abnormalities and acid base disturbances should be corrected. Persistent torsades de pointes may be treated with magnesium sulphate 2 g (20 ml of 10% solution) intravenously over 30-120 seconds, repeated twice at intervals of 5-15 minutes if necessary. Alternatively, if these measures fail, the arrhythmia may be abolished by increasing the underlying heart rate. This can be achieved by atrial and ventricular pacing or by isoprenaline (isoproterenol) infusion to achieve a heart rate of 90-110 beats per minute. Torsades de pointes is usually not helped by antiarrhythmic drugs and those which prolong the QTc interval (e.g. amiodarone, quinidine) may make it worse.

Since it has been reported that physostigmine may cause severe bradycardia, asystole and seizures, its use is not recommended in cases of overdose with clomipramine. Haemodialysis or peritoneal dialysis are ineffective because of the low plasma concentrations of clomipramine.

### **For convulsions:**

Diazepam should be given iv or other anticonvulsants such as phenobarbitone or paraldehyde (these substances may exacerbate existing respiratory failure, hypotension, or coma).

## **5. Pharmacological properties:**

### **5.1 Pharmacodynamic properties**

**Pharmacotherapeutic group:** Antidepressants, Non-selective monoamine reuptake inhibitors, ATC code: N06AA04.

### **Mechanism of action:**

The therapeutic activity of clomipramine is believed to be based on its ability to inhibit the neuronal re-uptake of noradrenaline (NA) and serotonin (5-HT) released in the synaptic cleft, with inhibition of 5-HT reuptake being the more important of these activities.

Clomipramine also has a wide pharmacological spectrum of action, which includes alpha1-adrenolytic, anticholinergic, antihistaminic, and antiserotonergic (5-HT-receptor blocking) properties.





**AGOG Pharma Ltd.**

(WHO - GMP CERTIFIED - GOVT RECOGNISED EXPORT HOUSE)



CIN: 11-06918

Regd. Office & Factory : Plot No. 33, Sector II, The Vasai Taluka Industrial Co-op. Estate Ltd. Gauripada, Vasai (E), Dist. Thane - 401 208. INDIA.  
Tel. : 95250 - 2455801 / 2452714 / 2453525 • Fax : 95250 - 2452074 (0091 - 250 - 2452074) • Email : agog@vsnl.net & agogpharma@rediffmail.com

## 5.2 Pharmacokinetic properties

### Absorption

The active substance is completely absorbed following oral administration and intramuscular injection.

The systemic bioavailability of unchanged clomipramine is reduced by 50% by "first-pass" metabolism to desmethylclomipramine (an active metabolite). The bioavailability of clomipramine is not markedly affected by the ingestion of food but the onset of absorption and therefore the time to peak may be delayed. Coated tablets and sustained release tablets are bioequivalent with respect to amount absorbed.

During oral administration of constant daily doses of clomipramine the steady state plasma concentrations of clomipramine and desmethylclomipramine (active metabolite) and the ratio between these concentrations show a high variability between patients, e.g. 75 mg clomipramine daily produces steady state concentrations of clomipramine ranging from about 20 to 175 ng/ml. Levels of desmethylclomipramine follow a similar pattern but are 40-85% higher.

Clomipramine slows gastro-intestinal transit time, absorption can, however, be delayed, particularly in overdose.

### Distribution

Clomipramine and desmethylclomipramine are widely distributed throughout the body and is 97.6% bound to plasma and tissue protein. The apparent volume of distribution is about 12-17 l/kg body weight. Concentrations in cerebrospinal fluid are about 2% of the plasma concentration.

It is reported to have a low and variable bioavailability following oral administration (48.2% of that after intravenous administration) and this has been related to extensive first-pass hepatic metabolism. Following single oral doses of 50 mg and 100 mg in healthy volunteers peak plasma concentrations of clomipramine of  $28.8 \pm 11.2$  ng/ml range 16.5 to 53 ng/ml (at 3 to 5 hours post-dose) and 70-140 ng/ml (at 1 to 2.5 hours post-dose) respectively are reported). Peak plasma concentrations of desmethylclomipramine of  $5.0 \pm 1.4$  ng/ml (range 2.9 to 7.8 ng/ml) have been reported to occur between 5 to 12 hours after a single oral dose of 50 mg.

After chronic administration in depressed patients steady state plasma concentrations of clomipramine have been noted to vary 20 to 30 fold. Vandell et al reported that following repeated doses of 75 mg a day for 1 month, steady state plasma concentrations of clomipramine and desmethylclomipramine were  $124.5 \pm 94$  ng/ml and  $144.8 \pm 113$  ng/ml respectively.

### Biotransformation

The major route of transformation of clomipramine is demethylation to desmethylclomipramine. In addition, clomipramine and desmethylclomipramine are hydroxylated to 8-hydroxy-clomipramine and 8-hydroxy-desmethyl-clomipramine but little is known about their activity *in vivo*. The hydroxylation of clomipramine and desmethylclomipramine is under genetic control similar to that of debrisoquine. In poor



**AGOG Pharma Ltd.**

(WHO - GMP CERTIFIED - GOVT RECOGNISED EXPORT HOUSE)



024 11-86518

Regd. Office & Factory : Plot No. 33, Sector II, The Vasai Taluka Industrial Co-op. Estate Ltd. Gauraipada, Vasai (E), Dist. Thane - 401 208, INDIA.  
Tel. : 95250 - 2455801 / 2452714 / 2453525 • Fax : 95250 - 2452074 (0091 - 250 - 2452074) • Email : agog@vsni.net & agogpharma@rediffmail.com

metabolisers of debrisoquine this may lead to high concentrations of desmethylclomipramine; concentrations of clomipramine are less significantly influenced.

### **Elimination**

Oral clomipramine is eliminated from the blood with a mean half-life of 21 hours (range 12-36 h), and desmethylclomipramine with a half-life of 36 hours.

About two-thirds of a single dose of clomipramine is excreted in the form of water-soluble conjugates in the urine, and approximately one-third in the faeces. The quantity of unchanged clomipramine and desmethylclomipramine excreted in the urine amounts to about 2% and 0.5% of the administered dose respectively.

### **Elderly**

In the elderly, plasma clomipramine concentrations may be higher for a given dose than would be expected in younger patients because of reduced metabolic clearance.

### **Hepatic and renal impairment**

The effects of hepatic and renal impairment on the pharmacokinetics of clomipramine have not been determined.

## **5.3 Preclinical safety data**

### **Repeat-dose toxicity**

Phospholipidosis and testicular changes considered to be secondary to the phospholipidosis, commonly associated with tricyclic compounds, have been observed with clomipramine hydrochloride at doses  $\geq 4$  fold greater than the maximum recommended human daily dose (MRHD). The clinical relevance of these findings is unknown.

### **Reproductive toxicity**

Clomipramine hydrochloride demonstrated evidence of embryotoxicity *e.g.* increased embryoletality and growth retardation, in the rat and mouse studies (at doses which are 5 to 10 times the estimated oral MRHD of 5 mg/kg on a mg/kg basis), but not in the rabbit study. The safety margin for increased embryoletality based on the administered dose is 2.5 times the oral MHRD.

No teratogenic effects were detected in mice, rats, and rabbits at doses up to 100, 50, and 60 mg/kg, respectively.

### **Mutagenicity**

Various *in vitro* and *in vivo* mutagenicity tests were performed and did not reveal any mutagenic activity of clomipramine hydrochloride.

### **Carcinogenicity**

The administration of clomipramine hydrochloride to mice and rats for 104 weeks did not show any evidence of carcinogenicity at dose levels representing 16 - 20 times the estimated oral MRHD of 5 mg/kg on a mg/kg basis



**AGOG Pharma Ltd.**

(WHO - GMP CERTIFIED - GOVT RECOGNISED EXPORT HOUSE)



CPA: 11-06518

Regd. Office & Factory : Plot No. 33, Sector II, The Vasai Taluka Industrial Co-op. Estate Ltd. Gauripada, Vasai (E), Dist. Thane - 401 208. INDIA.  
Tel. : 95250 - 2455801 / 2452714 / 2453525 • Fax : 95250 - 2452074 (0091 - 250 - 2452074) • Email : agog@vsnl.net & agogpharma@rediffmail.com

## 6. Pharmaceutical particulars:

### 6.1 List of Excipients:

Lactose	BP
Microcrystalline cellulose Powder	BP
Maize starch	BP
Methyl Paraben Sodium	BP
Propyl Paraben Sodium	BP
Iso Propyl alcohol	BP
Poly Vinyl Pyrrolidone K-30	BP
Talcum	BP
Magnesium stearate	BP
Maize Starch	BP
Cross Carmellose Sodium	BP
Colloidal silicon dioxide	BP
Polyplasdone XI-10 (Cross Povidone)	USP
Sodium Starch Glycolate	BP
Colour Instacoat Universal A05D01207 Yellow	INH
Iso Propyl alcohol	BP
Methylene Dichloride	BP

### 6.2 Incompatibilities:

None Reported

### 6.3 Shelf-Life:

36 months from the date of manufacture.

### 6.4 Special Precautions for Storage:

Store in a cool, dry and dark place. Protect from light.

### 6.5 Nature and Contents of Container:

10 tablets packed in one Blister. Such 1 blister packed in unit printed duplex board carton along with its package insert. Such cartons packed in export worthy shipper.

### 6.6 Special precautions for disposal:

None reported.

## 7. Registrant:

**AGOG PHARMA LTD.**

Plot No. 33, Sector II,  
The Vasai Taluka Industrial  
Co-Op. Estate Ltd., Gauripada,  
Vasai (E), Dist. Thane, India.



**AGOG Pharma Ltd.**

(WHO - GMP CERTIFIED - GOVT RECOGNISED EXPORT HOUSE)



CPA 11-00510

**Regd. Office & Factory :** Plot No. 33, Sector II, The Vasai Taluka Industrial Co-op. Estate Ltd. Gauripada, Vasai (E), Dist. Thane - 401 208. INDIA.  
Tel. : 95250 - 2455801 / 2452714 / 2453525 • Fax : 95250 - 2452074 (0091 - 250 - 2452074) • Email : agog@vsnl.net & agogpharma@rediffmail.com

- 8. Manufacturer:**  
**AGOG PHARMA LTD.**  
Plot No. 33, Sector II,  
The Vasai Taluka Industrial  
Co-Op. Estate Ltd., Gauripada,  
Vasai (E), Dist. Thane,  
India.
- 9. Date of revision of the text :**