



MYCOMIN

(Beclomethasone Dipropionate, Neomycin Sulfate & Miconazole Nitrate Cream)

INSERT:

114x150mm

<p style="text-align: center;"> MYCOMIN</p> <p style="text-align: center;">Beclomethasone Dipropionate, Neomycin Sulfate and Miconazole Nitrate Cream</p> <p>Composition: Beclomethasone Dipropionate USP.....0.025%w/w Neomycin Sulfate BP.....0.5%w/w Miconazole Nitrate BP.....2.0%w/w Cream Base.....q.s.</p> <p>Product description: A white colour semi solid mass filled in printed laminated tube.</p> <p>Mechanism of action: Beclomethasone is anti-inflammatory actions of corticosteroids are thought to involve lipocortins, phospholipase A2 inhibitory proteins which, through inhibition arachidonic acid, control the biosynthesis of prostaglandins and leukotrienes. Neomycin binds to specific 30S-subunit proteins and 16S rRNA of the bacteria for the treatment or prevention of bacterial infections. Miconazole interacts with 14-α demethylase, a cytochrome P-450 enzyme necessary to convert lanosterol to ergosterol. As ergosterol is an essential component of the fungal cell membrane, inhibition of its synthesis results in increased cellular permeability causing leakage of cellular contents. Miconazole may also inhibit endogenous respiration, interact with membrane phospholipids, inhibit the transformation of yeasts to mycelial forms, inhibit purine uptake, and impair triglyceride and/or phospholipid biosynthesis. The antifungal drug, miconazole nitrate, inhibits the growth of several species of <i>Candida</i>.</p> <p>Indications MYCOMIN Cream is used for the treatment, control, prevention, & improvement of the following diseases, conditions and symptoms:</p> <ul style="list-style-type: none"> • Cutaneous candidiasis • Corticosteroid-responsive dermatoses • Microbial infections • Tinea pedis • Tinea cruris • Tinea corporis • Tinea versicolor <p>Posology & route of administration Apply MYCOMIN Cream gently to the affected part 2 times a day for 2-4 days, depending upon severity of condition.</p> <p>Special warnings and precautions for use: Before using MYCOMIN Cream, inform your doctor about your current list of medications, over the counter products (e.g. vitamins, herbal supplements, etc.), allergies, pre-existing diseases, and current health conditions (e.g. pregnancy, upcoming surgery, etc.). Some health conditions may make you more susceptible to the side-effects of the drug. Take as directed by your doctor or follow the direction printed on the product insert. Dosage is based on your condition. Tell your doctor if your condition persists or worsens. Important counseling points are listed below.</p> <ul style="list-style-type: none"> • Avoid contact with the eyes • Avoid wearing plastic pants or tight-fitting diapers on a child being treated in the diaper area • Do not bandage, cover or wrap the treated skin area • Do not swallow • Do not take by mouth. Consult with your doctor before using this medicine on open wounds, dry, chapped, irritated, or sun-burned skin. Wash your hands before and after applying Beclomethasone Dipropionate, Neomycin Sulfate and Miconazole Nitrate Cream. • Clean and dry the skin area to be treated. • Do not wash the treated area after immediately applying Beclomethasone Dipropionate, Neomycin Sulfate and Miconazole Nitrate Cream. Also avoid the use of other products on the treated area unless directed by your doctor. • Applying an excessive amount may result in pilling. Use a thinner layer or lesser quantity of medicine to avoid pilling. • Avoid getting this medication in your eyes or nose or mouth. 	<p>Interaction with other medicinal products and other forms of interaction: If you use other drugs or over the counter products at the same time, the effects of Beclomethasone Dipropionate, Neomycin Sulfate & Miconazole Nitrate Cream may change. This may increase your risk for side-effects or cause your drug not to work properly. Tell your doctor about all the drugs, vitamins, and herbal supplements you are using, so that your doctor can help you prevent or manage drug interactions. Beclomethasone Dipropionate, Neomycin Sulfate & Miconazole Nitrate Cream may interact with the following drugs and products:</p> <ul style="list-style-type: none"> • Corticosteroids • Cyclosporine • Warfarin <p>Contraindication: Hypersensitivity to any of the ingredients Beclomethasone Dipropionate, Neomycin Sulfate and Miconazole Nitrate Cream is a contraindication. In addition, Beclomethasone Dipropionate, Neomycin Sulfate and Miconazole Nitrate Cream should not be used if you have the following conditions:</p> <ul style="list-style-type: none"> • History of hypersensitivity <p>Fertility, pregnancy and lactation: Certain medicines should not be used during pregnancy or breastfeeding. However, other medicines may be safely used in pregnancy or breastfeeding providing the benefits to the mother outweigh the risks to the unborn baby. Always inform your doctor if you are pregnant or planning a pregnancy, before using any medicine.</p> <ul style="list-style-type: none"> • The safety of this medicine during pregnancy and breastfeeding is not established. It is not recommended for use during pregnancy or breastfeeding. Consult your doctor for further advice. <p>Effects on ability to drive and use machines If you experience drowsiness, dizziness, hypotension or a headache as side-effects when using Beclomethasone Dipropionate, Neomycin Sulfate and Miconazole Nitrate Cream medicine then it may not be safe to drive a vehicle or operate heavy machinery. Pharmacists also advise patients not to drink alcohol with medicines as alcohol intensifies drowsiness side-effects. Please check for these effects on your body when using Beclomethasone Dipropionate, Neomycin Sulfate and Miconazole Nitrate Cream. Always consult with your doctor for recommendations specific to your body and health conditions.</p> <p>Overdose:</p> <ul style="list-style-type: none"> • Do not use more than prescribed dose. Taking more medication will not improve your symptoms; rather they may cause poisoning or serious side-effects. If you suspect that you or anyone else who may have overdosed of Beclomethasone Dipropionate, Neomycin Sulfate and Miconazole Nitrate Cream, please go to the emergency department of the closest hospital or nursing home. Bring a medicine box, container, or label with you to help doctors with necessary information. <p>Side effects: The following is a list of possible side-effects that may occur from all constituting ingredients of Beclomethasone Dipropionate, Neomycin Sulfate and Miconazole Nitrate Cream. This is not a comprehensive list. These side-effects are possible, but do not always occur. Some of the side-effects may be rare but serious. Consult your doctor if you observe any of the following side-effects, especially if they do not go away.</p> <ul style="list-style-type: none"> • Burning • Itching • Irritation • Dryness • Folliculitis • Hypertrichosis <p>Storage: Store at a temperature not exceeding 30°C. Protect from light. Do not freeze.</p> <p>For External Use Only</p> <p>Presentation : 15g filled in Printed lamitube & enclosed in carton</p> <p>Manufactured for :</p> <p style="text-align: center;"> Galaxy Pharmaceuticals</p> <p>B. No. : 36, Gala No. 2,3,4A, Bhiwandi, Thane (Mumbai) M.S. India</p> <p style="text-align: right;">Manufactured in India by : Curetech Skincare Plot No. 33 & 34, Phase-IV, Bhatoli Kalan, Baddi, Distt. Solan (HP)-173205</p>
--	---