	Artwork Details	Ca	arto
Product	SEAGRA ORAL JELLY	Size:	10
Exporting Country:	RWANDA	GSM:	6(
Artwork Code:		Paper:	М
Bar Code:		Colour:	В
Pack Type:		Side:	2
Date:	19.06.2022	Fo <b>l</b> d:	L:
		Folding Size:	50

Carton Packaging Details			
Size:	100 x 200 mm		
GSM:	60		
Paper:	Maplitho		
Colour:	Black		
Side:	2 side		
Fo <b>l</b> d:	L:2, H:2		
Folding Size:	50 x 50 mm		

# 100 mm

## SEAGRA ORAL JELLY

### COMPOSITION:

Each combi pack contains A. Sildenafil Oral Jelly (One Sachet)

Each 5 gm contains: Sildenafil Citrate 100 mg Ea. to Sildenafil Flavoured base Flavour : Banana

B. Sildenafil Oral Jelly (One Sachet) . Sildenafil Oral Jelly (Or Each 5 gm contains: Sildenafil Citrate BP Eq. to Sildenafil Flavoured base Flavour : Strawberry 100 mg

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you. Keep this leaflet. You may need to read it again. If you have any further questions, ask your doctor, pharmacist or nurse. This medicine has been prescribed for you only. You should not give it to others. It may harm them, even if they have the same signs of illness. If you get any side effects, talk to your doctor, pharmacist or nurse. See section 4.

 WHAT SEAGRA ORAL JELLY IS AND WHAT IT IS USED FOR SEAGRA ORAL JELLY contains the active substance sildenaff i citrate, which belongs to a group of medicines called phosphotesterase type 5 (PDES) inhibitors. This medicine works by relaxing the blood vessels of the penis, allowing the influx of blood to the penis when sexually stimulated. This medicine will only help you get an erection if you are sexually stimulated. This medicine is a treatment for adult men with erectile dysfunction, more commonly known as impotence. That is, when a man can not obtain, or maintain, a

# penile rigidity in erection, suitable for sexual activity. 2. WHAT YOU NEED TO KNOW BEFORE YOU TAKE SEAGRA ORAL JELLY

- What Trounced to Invitor and Traces are the American Control of the Ameri blood pressure. Tell your doctor if you are taking any of these medicines, which are usually used for relief of angina pectoris (or "chest pain"). If you have any further questions, ask your doctor or
- If you are using any of the medicines known as nitric oxide donors, such as amyl nitrite (poppers), as the
- If you are using any of the medicines known as nitric oxide donors, such as amyl nitrite (poppers), as the combination may lead to a potentially dangerous decrease in your blood pressure. If you are taking riociguat. This drug is used to treat pulmonary arterial hypertension (ie, high blood pressure in the lungs) and chronic thromboembolic pulmonary hypertension (ie, high blood pressure in the lungs due to the presence of clots). PDF inhibitors, such as SEAGRA ORAL JELU, have been shown to increase the hypotensive effects ofthis drug. If you are taking riociguat or if you have further questions, ask your dotor. If you have severe heart or liver problems.

  If you have ceretain are hereditary eye diseases (such as retinitis pigmentosa). If you have the advision loss due to non-arterial enterior ischemic optic neuropathy (NAION). rmings and Precautions

  If you have scrick in pharmacist or nurse before taking SEAGRA ORALJELLY

  If you have sickle cell anemia (a red blood cell anomaly), leukemia (blood cell cancer), multiple myeloma (bloom emrorov cancer).

- If you have sickle cell anemia (a red blood cell anomaly), leuxemia (blood cell anomaly), myeloma (bone marrow cancer).

  In have penile deformation or Peyronie's disease.

  If you have heart problems, in this case, your doctor should carefully evaluate whether your heart supports the extra effort associated with a sexual relationship.

  If you currently have a stomach ulcer orab leeding problem (such as hemophilia).

  If you have sudden or sudden loss of vision, stop taking this medicine and contact your doctor

mediately You should not take this medicine together with any other oral or local treatments for erectile dysfunction. You should not take SEAGRA ORAL JELLY if you do not have erectile dysfunction. You should not take SEAGRA ORAL JELLY if you are a woman. You should not use this medicine concurrently

You should not take ESAGRA ORAL JELLY if you are a woman. You should not use this medicine concurrently with pulmonary arterial hypertension [PAH] containing sildenally in or any other PDE5 inhibitors.

Special care to be taken in patients with kidney or liver problems

You should tell your doctor if you have kidney or liver problems. Your doctor may prescribe a lower dose.

Children and adolescents

This medicinal product must not be administered to individuals under the age of 1 8 years.

Other medicines and SEAGRA ORAL JELLY

Tell your doctor or pharmacist if you are taking or have recently taken any other medicines.

SEAGRA ORAL JELLY oral jelly may interfere with some medications, especially those used to treat "chest pain." In case of medical emergency, you should inform your doctor, pharmacist or nurse who is taking this medicine and when you did. Do not take this medicine with other medicines underso your doctor tells you to. medicine and when you did. Do not take this medicine with other medicines unless your doctor tells you to. You should not take this medicine with other medicines unless your doctor tells you to. cause a danger to take the your properties of the your doctor, pharmacist or nurse if you are taking any of these medicines can any of these medicines can you should not go any of the medicines known for the properties of your should not you should not go any of the medicines known for the properties of your doctor, such as

to us shown to take this mediume if you are using any on the mediumes in KNOWN as limit to white boilds, such as anyl nitrite (poppers), as the combination may also lead a diagerous drop in your blood pressure. If you are a liready taking riodiguat, town is your doctor or planmacist. If you are taking medicines known so you are to the planmacist. If you are taking medicines that treatment with the lowest dose (25 mg) of this medicine. Some patients taking alpha blockers for treatment of high blood pressure or enlarged prostate may

experience dizziness or a feeling of faintness that may be caused by low blood pressure when the subject sits or stands up quickly. Some patients had these symptoms when they took \$EAGRA DRAL JELLY with alpha blockers. These conditions are most likely to occur within 4 hours after taking this medicine. To reduce the likelihood of these symptoms occurring, you should be taking a regular daily dose of your alpha blocker before starting treatment with this drug. Your doctor may recommend that you start treatment with the

lowest dose (25 mg) of this medicine. SEAGRA ORAL IELLY with Food, drinks and alcohol. This medicine may be taken with or without food. However, you may find that this medication may take longer to work if you take it with a heavier meal.

Drinking alcohol can temporarily prevent the ability to get an erection. To get the maximum benefit from themedicine, it is advised not to drink large quantities of alcoholic beverages before taking this medicine. Pregnancy, breastfeeding and fertility

Fregians, briessteeding and returnity. This medicine is not intended for use by women. Driving and using machines. This medicine may cause dizziness and may affect your vision. You should be aware of how you respond to

Into medicine may cause dizzness and may arrect your vision, rou should be aware or now you respond to this medicine before driving or using machines.

3. HOW TO TAKE SEAGRA ORAL JELLY Take this medicine exactly as your doctor or pharmacist has told you. Talk to your doctor or pharmacist if you are not sure. The recommended starting dose is 50 mg. You should not use SEAGRA ORAL JELLY More than once a day. You should take this medicine about one hour before the scheduled time for sexual activity. Take the entire seadcass with shape for fuzions.

package with a glass of water. You should check with your doctor or pharmacist if you feel that the effect of this medicine is too strong or

too weak. This medicine will only help you get an erection if you are sexually stimulated. The length of time this drug The second of the seco takes to work varies from person to person, but usually it varies from half an hour to an hour. You may find that this medication takes longer to work if you take it with a heavier meal

that this medication also longer to work in you take it with a neavier meal. If this medicine so not help you got an erection or if your erection does not last long enough to complete the intercourse, you should tell your doctor. If you take more SEAGRA ORAL JELLY than you should You may experience an increase in side effects and their severity. Doses greater than 100 mg do not increase

You should not take more SEAGRA ORAL JELLY than your doctor has told you. If you take more tablets than you should, contact your doctor.

you should, contact your doctor.

If you have any further questions on the use of this product, ask your doctor, pharmacist or nurse.

4. POSSIBLE SIDE EFFECTS

Like all medicines, this medicine can cause side effects, although not everybody gets them. The reported side effects associated with the use of this medication are usually mild to moderate and of short duration. If you experience any of the following serious side effects, stop taking SEAGRA ORAL JELLY and get medical you experience any of the following serious side effects, stop taking schakha OKAL LELLY and get medical help right away. Allegic reaction - occurs infequently (can affect up to 1 in 100 people) symptoms include sudden wheezing, difficulty breathing or dizziness, swelling of the eyelids, face, lips, or throat.

(If to ccurs during or after the sexual act:

Sit in a semi-seated position and try to relax.

Do not use nitrates to treat your chest pain.

Prolonged and sometimes painful erections - occur rarely (may affect up to 1 in 1000 people).

If you have an erection that lasts more than 4 hours continuously, you should contact a doctor immediately.

- immediately.

  They of eal is sudden decrease or loss of vision occur rarely Serious skin reactions rarely occur.

  Symptoms may include peeling and severe skin swelling, blisters in the mouth, genitals and around the eyes,

rever.

Selizures or seizures rarely occur Other side effects:

Very common (can affect more than 1 in 10 people) is headache.

Common (may affect up to 1 in 10 people) is nause, a ficial redness, hot flashes (symptoms include a sudden feeling of warmth in the upper body), indigestion, colored vision, blurred vision, visual disturbance, stuffy

nose, and dizziness.

Uncommon (may affect up to 1 in 100 people): vomiting, skin rash, eye irritation, red eyes, somnolence, visual twitches, sharpness, light sensitivity, watery eyes, palpitations, rapid heartbeat, high blood pressure, low blood pressure, pain muscle weakness, drowsiness, tingling sensation, dizziness, ringing in the ears, dry mouth, clogged nasal sinuses, inflammation of the nose lining (symptoms include runny nose, sneezing and

mouth, clogged nasal sinuses, inflammation of the nose lining (symptoms include runny nose, sneezing and stuffy nose), high abdominal pain, gastroesophageal reflux disease (ossiomas include heartburn), blood in the urine, pain in the arms or legs, nosebleed, feeling hot and feeling tired Rare (may affect up or 1 in 1000 people): fainting, stroke, heart attack, irregular heartbeat, temporary decrease in blood flow to certain regions of the brain, sore throat, numb mouth, retinal hemorrhage, double vision decreased visual acuity, abnormal sensation in the eye, swelling of the eye or eyelid, small particles or spots in the eyes, vision of halos around the lights, dilation of the uppul of the eye, change in the color of the sclera, bleeding of the penis, presence of blood in semen, dry nose, swelling of the inside of the nose, in postmarketing experience, rare cases of unstable angina (cardiac problem), and sudden detath have been reported. It should be noted that most, but not all, men who had these side effects already suffered from heart problems before taking this medication. It is not possible to determine if these events were directly related to taking this medication.

related to taking this medication.

5. HOW TO KEEP SAGRA ORAL PELLY
Store in dry place, below 30°C, protect from light.
Keep medicines out of reach of children.
6. PRESENTATION: 2 Sachet packed in a nouter



Department	Design	QA	QA	QC	Production	Factory Manager	QA Manager	General Manager
Name:								
Sign:								
Date:								
. 1								l

200	mm