

**SUMMARY OF PRODUCT CHARACTERISTICS****1. Name of the finished pharmaceutical product****INN Name:** BINFIN 5 (Finasteride Tablets 5 mg)**Trade Name:** BINFIN 5**Strength:** 5 mg**Pharmaceutical form:** Tablets**2. Qualitative and quantitative composition**

Each film coated tablet contains 5 mg of Finasteride Ph.Eur.

3. Pharmaceutical form**Dosage form:** Tablets**Description:**

Film coated tablets

Blue colour, round film coated tablets, debossed with 'H' on one side and '37' on other side.

4. Clinical particulars**4.1 Therapeutic indications**

Finasteride 5 mg is indicated for the treatment and control of benign prostatic hyperplasia (BPH) in patients with an enlarged prostate to:

- cause regression of the enlarged prostate, improve urinary flow and improve the symptoms associated with BPH
- reduce the incidence of acute urinary retention and the need for surgery including transurethral resection of the prostate (TURP) and prostatectomy.

4.2 Posology and method of administrationPosology

The recommended adult dose is one 5 mg tablet daily, with or without food.

Finasteride 5 mg can be administered alone or in combination with the alpha-blocker doxazosin.



Although early improvement in symptoms may be seen, treatment for at least six months may be necessary to assess whether a beneficial response has been achieved. The risk of acute urinary retention is reduced within four months of treatment.

Dosage in the elderly

Dosage adjustments are not necessary although pharmacokinetic studies have shown that the elimination rate of finasteride is slightly decreased in patients over the age of 70.

Dosage in hepatic insufficiency

The effect of hepatic insufficiency on the pharmacokinetics of finasteride has not been studied

Dosage in renal insufficiency

No adjustment in dosage is required in patients with varying degrees of renal insufficiency (creatinine clearances as low as 9 mL/min), as pharmacokinetic studies did not indicate any change in the disposition of finasteride.

Pediatric population

Finasteride is contraindicated in children.

Method of administration

Finasteride is for oral use only

4.3 Contraindications

Finasteride 5 mg is not indicated for use in women or children.

Finasteride 5 mg is contraindicated in the following:

- Hypersensitivity to any component of this product
- Pregnancy - Use in women when they are or may potentially be pregnant.

4.4 Special warnings and precautions for use

General

Patients with large residual urine and/or severely diminished urinary flow should be carefully monitored for obstruction uropathy.

Consultation of an urologist should be considered in patients treated with finasteride.



Obstruction due to trilobular growth pattern of the prostate should be excluded before starting treatment with finasteride

To avoid obstructive complications it is important that patients with large residual urine and/or heavily decreased urinary flow are carefully controlled.

The possibility of surgery should be an option.

Women who are pregnant or may become pregnant should not handle crushed or broken finasteride tablets because of the possibility of absorption of finasteride and the subsequent potential risk to a male foetus. Finasteride tablets have a film coating which prevents contact with the active ingredient provided that the tablets have not been broken or crushed (see sections 4.6 and 6.6).

Effects on prostate-specific antigen (PSA) and prostate cancer detection

No clinical benefit has yet been demonstrated in patients with prostate cancer treated with finasteride. Patients with BPH and elevated PSA were monitored in controlled clinical studies with serial PSAs and prostate biopsies. In these BPH studies, finasteride did not appear to alter the rate of prostate cancer detection and the overall incidence of prostate cancer was not significantly different in patients treated with finasteride or placebo.

Digital rectal examination, as well as other evaluations for prostate cancer, should be performed on patients with BPH prior to initiating therapy with finasteride and periodically thereafter. Serum PSA is also used for prostate cancer detection. Generally a baseline PSA >10 ng/mL (Hybritech) prompts further evaluation and consideration of biopsy; for PSA levels between 4 and 10 ng/mL, further evaluation is advisable. There is considerable overlap in PSA levels among men with and without prostate cancer. Therefore, in men with BPH, PSA values within the normal reference range do not rule out prostate cancer regardless of treatment with finasteride. A baseline PSA <4 ng/mL does not exclude prostate cancer.



Finasteride causes a decrease in serum PSA concentrations by approximately 50% in patients with BPH, even in the presence of prostate cancer. This decrease in serum PSA levels in patients with BPH treated with finasteride should be considered when evaluating PSA data and does not rule out concomitant prostate cancer. This decrease is predictable over the entire range of PSA values, although it may vary in individual patients. In patients treated with finasteride for six months or more, PSA values should be doubled for comparison with normal ranges in untreated men. This adjustment preserves the sensitivity and specificity of the PSA assay and maintains its ability to detect prostate cancer.

Any sustained increase in PSA levels of patients treated with finasteride should be carefully evaluated, including consideration of non-compliance to therapy with finasteride

Percent free PSA (free to total PSA ratio) is not significantly decreased by finasteride and remains constant even under the influence of finasteride. When percent free PSA is used as an aid in the detection of prostate cancer, no adjustment to its value is necessary.

Drug/laboratory test interactions

Effect on levels of PSA

Serum PSA concentration is correlated with patient age and prostatic volume, and prostatic volume is correlated with patient age. When PSA laboratory determinations are evaluated, consideration should be given to the fact that PSA levels decrease in patients treated with finasteride. In most patients, a rapid decrease in PSA is seen within the first months of therapy, after which time PSA levels stabilize to a new baseline. The post-treatment baseline approximates half of the pre-treatment value. Therefore, in typical patients treated with finasteride for six months or more, PSA values should be doubled for comparison to normal ranges in untreated men. For clinical interpretation, see section 4.4 “Effects on prostate-specific antigen (PSA) and prostate cancer detection”.



Percent free PSA (free to total PSA ratio) is not significantly decreased by finasteride. The ratio of free to total PSA remains constant even under the influence of finasteride. When percent free PSA is used as an aid in the detection of prostate cancer, no adjustment to its value is necessary.

Breast cancer in men

Breast cancer has been reported in men taking finasteride 5 mg during clinical trials and the post-marketing period. Physicians should instruct their patients to promptly report any changes in their breast tissue such as lumps, pain, gynaecomastia or nipple discharge.

Mood alterations and depression

Mood alterations including depressed mood, depression and, less frequently, suicidal ideation have been reported in patients treated with finasteride 5 mg. Patients should be monitored for psychiatric symptoms and if these occur, the patient should be advised to seek medical advice.

Pediatric population

Finasteride is not indicated for use in children

Safety and effectiveness in children have not been established.

Lactose

The tablet contains lactose monohydrate. Patients with any of the following genetic deficiencies should not take this medicine: galactose intolerance, the Lapp lactase deficiency or glucose-galactose malabsorption.

Hepatic insufficiency

The effect of hepatic insufficiency on the pharmacokinetics of finasteride has not been studied.

4.5 Interaction with other medicinal products and other forms of interaction



No drug interactions of clinical importance have been identified. Finasteride is metabolized primarily via, but does not appear to affect significantly, the cytochrome P450 3A4 system. Although the risk for finasteride to affect the pharmacokinetics of other medicinal products is estimated to be small, it is probable that inhibitors and inducers of cytochrome P450 3A4 will affect the plasma concentration of finasteride. However, based on established safety margins, any increase due to concomitant use of such inhibitors is unlikely to be of clinical significance. Compounds which have been tested in man have included propranolol, digoxin, glibenclamide, warfarin, theophylline, and phenazone and no clinically meaningful interactions were found.

4.6 Fertility, pregnancy and lactation

Pregnancy: Finasteride 5 mg is contra-indicated in women when they are or may potentially be pregnant. Because of the ability of Type II 5 α -reductase inhibitors to inhibit conversion of testosterone to dihydrotestosterone, these drugs, including finasteride, may cause abnormalities of the external genitalia of a male foetus when administered to a pregnant woman.

In animal developmental studies, dose-dependent development of hypospadias were observed in the male offspring of pregnant rats given finasteride at doses ranging from 100 μ g/kg/day to 100 mg/kg/day, at an incidence of 3.6% to 100%. Additionally, pregnant rats produced male offspring with decreased prostatic and seminal vesicular weights, delayed preputial separation, transient nipple development and decreased anogenital distance, when given finasteride at doses below the recommended human dose. The critical period during which these effects can be induced has been defined in rats as days 16-17 of gestation.

The changes described above are expected pharmacological effects of Type II 5 α -reductase inhibitors. Many of the changes, such as hypospadias, observed in male rats exposed in utero to finasteride are similar to those reported in male infants with a genetic deficiency of Type II 5 α -reductase. It is for these reasons that Finasteride 5 mg is contra-indicated in women who are or may potentially be pregnant.

No effects were seen in female offspring exposed in utero to any dose of finasteride.

Exposure to finasteride - risk to male foetus



Women should not handle crushed or broken tablets of Finasteride 5 mg when they are or may potentially be pregnant because of the possibility of absorption of finasteride and the subsequent potential risk to a male foetus. Finasteride 5 mg tablets are coated and will prevent contact with the active ingredient during normal handling, provided that the tablets have not been broken or crushed.

Small amounts of finasteride have been recovered from the semen in subjects receiving finasteride 5 mg/day. It is not known whether a male foetus may be adversely affected if his mother is exposed to the semen of a patient being treated with finasteride. When the patient's sexual partner is or may potentially be pregnant, the patient is recommended to minimise exposure of his partner to semen.

Breast-feeding: Finasteride 5 mg is not indicated for use in women.

It is not known whether finasteride is excreted in human milk.

4.7 Effects on ability to drive and use machines

There are no data to suggest that Finasteride 5 mg affects the ability to drive or use machines.

4.8 Undesirable effects

The most frequent adverse reactions are impotence and decreased libido. These adverse reactions occur early in the course of therapy and resolve with continued treatment in the majority of patients.

The adverse reactions reported during clinical trials and/or post-marketing use are listed in the table below.

Frequency of adverse reactions is determined as follows:

Very common ($\geq 1/10$), Common ($\geq 1/100$ to $< 1/10$), Uncommon ($\geq 1/1,000$ to $< 1/100$), Rare ($\geq 1/10,000$ to $< 1/1,000$), Very rare ($< 1/10,000$), not known (cannot be estimated from the available data).




The frequency of adverse reactions reported during post-marketing use cannot be determined as they are derived from spontaneous reports.

System Organ Class	Frequency: adverse reaction
Immune system disorders	Unknown: hypersensitivity reactions including swelling of the lips, tongue, throat and face
Psychiatric disorders	Common: decreased libido Unknown: decreased libido that may continue after discontinuation of therapy, depression
Cardiac disorders	Unknown: palpitation
Hepatobiliary disorders	Unknown: increased hepatic enzymes
Skin and subcutaneous tissue disorders	Uncommon: rash Unknown: pruritus, urticaria
Reproductive system and breast disorders	Common: impotence Uncommon: ejaculation disorder, breast tenderness, breast enlargement. Unknown: testicular pain, sexual dysfunction (erectile dysfunction and ejaculation disorder) which may continue after discontinuation of treatment; male infertility and/or poor seminal quality. Normalization or improvement of seminal quality has been reported after discontinuation of finasteride.
Investigations	Common: decreased volume of ejaculate

In addition, the following has been reported in clinical trials and post-marketing use: male breast cancer.

Medical Therapy of Prostatic Symptoms (MTOPS)



The MTOPS study compared finasteride 5 mg/day (n=768), doxazosin 4 or 8 mg/day (n=756), combination therapy of finasteride 5 mg/day and doxazosin 4 or 8 mg/day (n=786), and placebo (n=737). In this study, the safety and tolerability profile of the combination therapy was.

generally consistent with the profiles of the individual components. The incidence of ejaculation disorder in patients receiving combination therapy was comparable to the sum of incidences of this adverse experience for the two monotherapies.


Other Long-Term Data

In a 7 year placebo-controlled trial that enrolled 18,882 healthy men, of whom 9060 had prostate needle biopsy data available for analysis, prostate cancer was detected in 803 (18.4%) men receiving Finasteride 5 mg and 1147 (24.4%) men receiving placebo. In the Finasteride 5 mg group, 280 (6.4%) men had prostate cancer with Gleason scores of 7-10 detected on needle biopsy vs 237 (5.1%) men in the placebo group. Additional analyses suggest that the increase in the prevalence of high-grade prostate cancer observed in the Finasteride 5 mg group may be explained by a detection bias due to the effect of Finasteride 5 mg on prostate volume. Of the total cases of prostate cancer diagnosed in this study, approximately 98% were classified as intracapsular (stage T1 or T2). The relationship between long-term use of Finasteride 5 mg and tumours with Gleason scores of 7-10 is unknown.

Laboratory Test Findings

Serum PSA concentration is correlated with patient age and prostatic volume, and prostatic volume is correlated with patient age. When PSA laboratory determinations are evaluated, consideration should be given to the fact that PSA levels are decreased in patients treated with finasteride (see section 4.4). In most patients, a rapid decrease in PSA is seen within the first months of therapy, after which time PSA levels stabilise to a new baseline. The post-treatment baseline approximates half of the pre-treatment value. Therefore, in typical patients treated with finasteride for six months or more, PSA values should be doubled for comparison to normal ranges in untreated men.

For clinical interpretation see 'Special warnings and precautions for use', Effects on prostate-specific antigen (PSA) and prostate cancer detection.



No other difference was observed in patients treated with placebo or Finasteride 5 mg in standard laboratory tests.

4.9 Overdose

No specific treatment of over dosage with Finasteride 5 mg is recommended. Patients have received single doses of Finasteride 5 mg up to 400 mg and multiple doses of Finasteride 5 mg up to 80 mg/day for up to three months without any adverse effects.

5. Pharmacological properties


5.1 Pharmacodynamic properties

Finasteride is a competitive inhibitor of human 5 α -reductase, an intracellular enzyme which metabolises testosterone into the more potent androgen, dihydrotestosterone (DHT). In benign prostatic hyperplasia (BPH), enlargement of the prostate gland is dependent upon the conversion of testosterone to DHT within the prostate. Finasteride 5 mg is highly effective in reducing circulating and intraprostatic DHT. Finasteride has no affinity for the androgen receptor.

In clinical studies of patients with moderate to severe symptoms of BPH, an enlarged prostate on digital rectal examination and low residual urinary volumes, Finasteride 5 mg reduced the incidence of acute retention of urine from 7/100 to 3/100 over four years and the need for surgery (TURP or prostatectomy) from 10/100 to 5/100. These reductions were associated with a 2-point improvement in QUASI-AUA symptom score (range 0-34), a sustained regression in prostate volume of approximately 20% and a sustained increase in urinary flow rate.

Medical therapy of prostatic symptoms

The Medical Therapy of Prostatic Symptoms (MTOPS) Trial was a 4- to 6-year study in 3047 men with symptomatic BPH who were randomised to receive finasteride 5 mg/day, doxazosin 4 or 8 mg/day*, the combination of finasteride 5 mg/day and doxazosin 4 or 8 mg/day*, or placebo. The primary endpoint was time to clinical progression of BPH, defined as a ≥ 4 point confirmed increase from baseline in symptom score, acute urinary retention, BPH-related renal insufficiency, recurrent urinary tract infections or urosepsis, or incontinence. Compared to placebo, treatment with finasteride, doxazosin, or combination therapy resulted in a significant reduction in the risk



of clinical progression of BPH by 34 (p=0.002), 39 (p<0.001), and 67% (p<0.001), respectively. The majority of the events (274 out of 351) that constituted BPH progression were confirmed ≥ 4 point increases in symptom score; the risk of symptom score progression was reduced by 30 (95% CI 6 to 48%), 46 (95% CI 25 to 60%), and 64% (95% CI 48 to 75%) in the finasteride, doxazosin, and combination groups, respectively, compared to placebo. Acute urinary retention accounted for 41 of the 351 events of BPH progression; the risk of developing acute urinary retention was reduced by 67 (p=0.011), 31 (p=0.296), and 79% (p=0.001) in the finasteride, doxazosin, and combination groups, respectively, compared to placebo. Only the finasteride and combination therapy groups were significantly different from placebo.

* Titrated from 1 mg to 4 or 8 mg as tolerated over a 3-week period


5.2 Pharmacokinetics properties

After an oral dose of ¹⁴C-finasteride in man, 39% of the dose was excreted in the urine in the form of metabolites (virtually no unchanged drug was excreted in the urine), and 57% of total dose was excreted in the faeces. Two metabolites have been identified which possess only a small fraction of the Type II 5 α -reductase activity of finasteride.

The oral bioavailability of finasteride is approximately 80%, relative to an intravenous reference dose, and is unaffected by food. Maximum plasma concentrations are reached approximately two hours after dosing and the absorption is complete within 6-8 hours. Protein binding is approximately 93%. Plasma clearance and the volume of distribution are approximately 165 ml/min and 76 l, respectively.

In the elderly, the elimination rate of finasteride is somewhat decreased. Half-life is prolonged from a mean half-life of approximately six hours in men aged 18-60 years to eight hours in men aged more than 70 years. This is of no clinical significance and does not warrant a reduction in dosage.

In patients with chronic renal impairment, whose creatinine clearance ranged from 9-55 ml/min, the disposition of a single dose of ¹⁴C-finasteride was not different from that in healthy volunteers. Protein binding also did not differ in patients with renal impairment. A portion of the



metabolites which normally is excreted renally was excreted in the faeces. It therefore appears that faecal excretion increases commensurate to the decrease in urinary excretion of metabolites. Dosage adjustment in non-dialysed patients with renal impairment is not necessary.

There are no data available in patients with hepatic insufficiency.

Finasteride has been found to cross the blood-brain barrier. Small amounts of finasteride have been recovered in the seminal fluid of treated patients.

5.3 Preclinical safety data

Non-clinical data reveal no special hazard for humans based on conventional studies of repeated dose toxicity, genotoxicity, and carcinogenic potential. Reproduction toxicology studies in male rats have demonstrated reduced prostate and seminal vesicular weights, reduced secretion from accessory genital glands and reduced fertility index (caused by the primary pharmacological effect of finasteride). The clinical relevance of these findings is unclear.

As with other 5-alpha-reductase inhibitors, feminisation of male rat foetuses has been seen with administration of finasteride in the gestation period. Intravenous administration of finasteride to pregnant rhesus monkeys at doses up to 800 ng/day during the entire period of embryonic and foetal development resulted in no abnormalities in male foetuses. This dose is about 60-120 times higher than the estimated amount in semen of a man who have taken 5 mg finasteride, and to which a woman could be exposed via semen. In confirmation of the relevance of the Rhesus model for human foetal development, oral administration of finasteride 2 mg/kg/day (the systemic exposure (AUC) of monkeys was slightly higher (3x) than that of men who have taken 5 mg finasteride, or approximately 1-2 million times the estimated amount of finasteride in semen) to pregnant monkeys resulted in external genital abnormalities in male foetuses. No other abnormalities were observed in male foetuses and no finasteride-related abnormalities were observed in female foetuses at any dose.”

6. Pharmaceutical particulars

6.1 List of excipients



Lactose monohydrate (Pharmatose 200M), Cellulose Microcrystalline (Avicel PH 102), Starch Pregelatinized (Starch 1500), Sodium starch glycolate (Primojel), Docusate sodium, Purified water, Magnesium Stearate and Opadry Blue 03A80928

6.2 Incompatibilities

Not applicable.

6.3 Shelf life

36 months

6.4 Special precautions for storage

Store below 30°C

6.5 Nature and contents of container

Blister pack: 15's pack

6.6 Special precautions for disposal and other handling

Women should not handle crushed or broken finasteride tablets when they are or may potentially be pregnant

7. Marketing Authorisation Holder and Manufacturing Site Addresses**7.1 Name and Address of Manufacturer**

Name: Hetero Labs Limited (Unit-III)

Business Address: Survey No.: 51, 22-110,
Industrial Development Area (IDA),
Jeedimetla, Hyderabad-500 055
Telangana

Country: INDIA

Phone: 91-40-23096171/172/173/174

Fax: 91-40-23095105

7.2 Name and Address of Principal

Name : Hetero Labs Limited

Business Address : 7-2-A2, Hetero Corporate,
Industrial Estates, Sanath Nagar,
Hyderabad-500 018



Telangana.
Country : INDIA
Telephone : 0091-040-23704923/24

8. Registration Number

Not Applicable

9. Category for Distribution

Prescription only medicine – List I

10. Date of Publication of This Package Insert



LABELLING

PARTICULARS TO APPEAR ON THE OUTER PACKAGING OR, WHERE THERE IS NO OUTER PACKAGING, ON THE IMMEDIATE PACKAGING

(a). NAME OF THE FPP

BINFIN 5 (Finasteride Tablets 5 mg)

(b). METHOD OF ADMINISTRATION

Oral

(c). A list of API(s) (using INNs if applicable) showing the amount of each present in a dosage unit, and a statement of the net contents of the container, e.g. number of dosage units, weight or volume.

Each film coated tablet contains 5 mg of Finasteride Ph.Eur.

(d). List of excipients known to be a safety concern for some patients, e.g. lactose, gluten, metabisulfites, parabens, ethanol, or tartrazine.

NA

(e). Indication(s) and recommended dosage, where practicable

Anti-Alopecia Agent, Testosterone-5-alpha reductase inhibitors

(f). The batch number assigned by the manufacturer.

< Batch > <Lot> <BN> {number}

(g). The manufacturing and expiry dates in an uncoded form.

MFG: MM/YYYY

EXP: MM/YYYY

(h) Storage conditions or handling precautions that may be necessary.

Store below 30°C

(i) Directions for use and any warnings or precautions that may be necessary.