



SCOTT-EDIL PHARMACIA LTD.

56, EPIP, Phase-I, Jharmajri, Baddi-173 205, (HP), India

1.6 Product information:

1.6.1 Prescribing Information (Summary of Product Characteristics)

1. NAME OF THE FINISHED PHARMACEUTICAL PRODUCT

IBUPLUS (Ibuprofen and Paracetamol Tablets)

1.1 Strength

400 mg/325mg

1.2 Pharmaceutical form

Tablet for Oral administration

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

2.1 Qualitative

Each uncoated tablet contains:

Ibuprofen BP 400mg

Paracetamol BP 325mg

Excipients q.s.

2.2 Quantitative declaration

S. No.	Ingredients	Specification	Qty. Req. per batch in Kg	Over ages	Actual Qty per batch	Actual Qty per Tablet (mg)	Function
ACTIVE PHARMACEUTICAL INGREDIENT							
1.	*Ibuprofen	BP	40.000	NIL	40.000	400.0	Active Ingredient (Anti-inflammatory, Analgesic)
2.	*Paracetamol	BP	32.500	NIL	32.500	325.0	Active Ingredient (Analgesic, Antipyretic)
EXCIPIENTS FOR GRANULATION							
3.	Maize starch	BP	6.000	NIL	6.000	60.0	Diluent



SCOTT-EDIL PHARMACIA LTD.

56, EPIP, Phase-I, Jharmajri, Baddi-173 205, (HP), India

4.	Microcrystalline Cellulose	BP	8.000	NIL	8.000	80.0	Diluent
5.	Povidone K- 30	BP	0.375	NIL	0.375	3.75	Disintegrant
6.	Maize starch (for paste)	BP	4.500	NIL	4.500	45.0	Binder
7.	Povidone K- 30 (for paste)	BP	0.375	NIL	0.375	3.75	Binder
8.	Methyl Hydroxybenzoate	BP	0.125	NIL	0.125	1.25	Antimicrobial preservative.
9.	Propyl Hydroxybenzoate	BP	0.025	NIL	0.025	0.25	Antimicrobial preservative.
EXCIPIENTS FOR LUBRICATION							
10.	Purified Talc	BP	2.500	NIL	2.500	25.0	Lubricant
11.	Magnesium Stearate	BP	3.000	NIL	3.000	30.0	Lubricant
12.	Sodium Starch Glycolate	BP	1.000	NIL	1.000	10.0	Disintegrant
13.	Croscarmellose Sodium	BP	0.800	NIL	0.800	8.0	Disintegrant
14.	Colloidal Anhydrous silica	BP	0.800	NIL	0.800	8.0	Glidant

* To be dispensed on 100% assay on as such basis and Tablet weight to be adjusted by reducing the quantity of Microcrystalline Cellulose.

2.3 Salts and hydrates

Not Applicable

2.4 Esters and pro-drugs

Not Applicable

2.5 Oral powders for solution or suspension

Not Applicable

2.6 Parenterals excluding powders for reconstitution

Not Applicable



SCOTT-EDIL PHARMACIA LTD.

56, EPIP, Phase-I, Jharmajri, Baddi-173 205, (HP), India

2.7 Powders for reconstitution prior to Parenteral administration

Not Applicable

2.8 Concentrates

Not Applicable

2.9 Transdermal patches

Not Applicable

2.10 Multidose solid or semi-solid products

Tablet for oral administration

2.11 Biological medicinal products

Not Applicable

3. PHARMACEUTICAL FORM

Tablets

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

Ibuplus tablets are indicated for the temporary relief of mild to moderate pain associated with migraine, headache, backache, period pain, dental pain, rheumatic and muscular pain, pain of non-serious arthritis, cold and flu symptoms, sore throat and fever. This product is especially suitable for pain which requires stronger analgesia than ibuprofen or paracetamol alone.

4.2 Posology and method of administration

Posology

For short term-use only.

Undesirable effects may be minimised by using the lowest effective dose for the shortest duration necessary to control symptoms.

The patient should consult a doctor if the symptoms persist or worsen or if the product is required for more than 3 days.

Adults: One tablet to be taken up to three times per day with water. Leave at least six hours between doses.



SCOTT-EDIL PHARMACIA LTD.

56, EPIP, Phase-I, Jharmajri, Baddi-173 205, (HP), India

If the one tablet dose does not control symptoms, a maximum of two tablets may be taken up to three times a day. Leave at least six hours between doses.

Do not take more than six tablets (3000mg Paracetamol, 1200mg Ibuprofen) in any 24 hours period.

To minimise side effects, it is recommended that patients take Nuromol with food.

Elderly: No special dosage modifications are required.

The elderly are at increased risk of the serious consequences of adverse reactions. If an NSAID is considered necessary, the lowest effective dose should be used for the shortest possible duration. The patient should be monitored regularly for gastrointestinal bleeding during NSAID therapy.

Not for use by children under 18 years.

4.3 Method of administration

For oral administration.

4.4 Contraindications

This product is contraindicated:

- In patients with a known hypersensitivity to ibuprofen, paracetamol or any other excipients in the product.
- In concomitant use with other Paracetamol-containing products – increased risk of serious adverse effects.
- In patients with a history of hypersensitivity reactions (e.g. bronchospasm, angioedema, asthma, rhinitis, or urticaria) associated with acetylsalicylic acid or other non-steroidal anti-inflammatory drugs (NSAIDs).
- In patients with Active, or a history of recurrent peptic ulcer/haemorrhage (two or more distinct episodes of proven ulceration or bleeding).
- In patients with a history of, or an existing gastrointestinal ulceration/perforation or bleeding, including that associated with NSAIDs.
- Patients with defects in coagulation.
- In patients with severe hepatic failure, severe renal failure or severe heart failure (NYHA Class IV) .
- In concomitant use with other NSAID containing products, including cyclo-oxygenase-2 (COX-2) specific inhibitors and doses of acetylsalicylic acid above 75 mg daily – increased risk of adverse reactions.



SCOTT-EDIL PHARMACIA LTD.

56, EPIP, Phase-I, Jharmajri, Baddi-173 205, (HP), India

- During the last trimester of pregnancy due to risk of premature closure of the foetal ductus arteriosus with possible pulmonary hypertension.

4.5 Special warnings and precautions for use

Do not exceed the recommended dose.

If symptoms persist consult your doctor.

Keep out of the sight and reach of children.

Paracetamol:

The hazards of paracetamol overdose are greater in patients with non-cirrhotic alcoholic liver disease. Immediate medical advice should be sought in the event of an overdose, even if the patient feels well, because of the risk of delayed, serious liver damage.

Do not take with any other paracetamol containing products. Immediate medical advice should be sought if this occurs, even if you feel well as this can result in an overdose.

Ibuprofen:

Undesirable effects may be minimised by using the lowest effective dose for the shortest duration necessary to control symptoms (gastrointestinal and cardiovascular risks below) and by patients taking the dose with food.

Elderly:

The elderly have an increased frequency of adverse reactions to NSAIDs especially gastrointestinal bleeding and perforation which may be fatal.

Caution is required in patients with certain conditions:

• Respiratory disorders:

In patients suffering from, or with a history of, bronchial asthma or allergic disease NSAIDs have been reported to precipitate bronchospasm.

• SLE and mixed connective tissue disease:

In patients with systemic lupus erythematosus (SLE) and mixed connective tissue disease disorders there may be an increased risk of aseptic meningitis.

• Cardiovascular and cerebrovascular effects

Appropriate monitoring and medical advice are required for patients with a history of hypertension and/or mild to moderate congestive heart failure as fluid retention, hypertension and oedema have been reported in association with NSAID therapy.

Clinical studies suggest that use of ibuprofen, particularly at a high dose (2400 mg/day) may be associated with a small increased risk of arterial thrombotic events (e.g. myocardial infarction or stroke). Overall, epidemiological studies do not suggest that low dose



SCOTT-EDIL PHARMACIA LTD.

56, EPIP, Phase-I, Jharmajri, Baddi-173 205, (HP), India

ibuprofen (e.g. ≤ 1200 mg/day) is associated with an increased risk of arterial thrombotic events.

Patients with uncontrolled hypertension, congestive heart failure (NYHA II-III), established ischaemic heart disease, peripheral arterial disease, and/or cerebrovascular disease should only be treated with ibuprofen after careful consideration and high doses (2400 mg/day) should be avoided. Careful consideration should be exercised before initiating long-term treatment for patients with risk factors for cardiovascular events (e.g. hypertension, hyperlipidaemia, diabetes mellitus, smoking) particularly if high doses of ibuprofen (2400 mg/day) are required.

- *Cardiovascular, renal and hepatic impairment:*

The administration of NSAIDs may cause a dose dependent reduction in prostaglandin formation and precipitate renal failure. Patients at greatest risk of this reaction are those with impaired renal function, cardiac impairment, liver dysfunction, those taking diuretics and the elderly. Renal function should be monitored in these patients.

- *Gastrointestinal effects:*

NSAIDs should be given with care to patients with a history of gastrointestinal disease (ulcerative colitis, Crohn's disease) as these conditions may be exacerbated.

Gastrointestinal (GI) bleeding, ulceration and perforation, which can be fatal, has been reported with all NSAIDs at any time during treatment, with or without warning symptoms or a previous history of serious GI events.

The risk of GI bleeding, ulceration or perforation is higher with increasing NSAID doses, in patients with a history of ulcer, particularly if complicated with haemorrhage or perforation and in the elderly. These patients should commence treatment on the lowest dose available. Combination therapy with protective agents (e.g. misoprostol or proton pump inhibitors) should be considered for these patients, and also for patients requiring concomitant low dose acetylsalicylic acid, or other drugs likely to increase gastrointestinal risk.

Patients with a history of GI toxicity, particularly the elderly, should report any unusual abdominal symptoms (especially GI bleeding) particularly in the initial stages of treatment.

Caution should be advised in patients receiving concomitant medications which could increase the risk of ulceration or bleeding, such as oral corticosteroids, anticoagulants such as warfarin selective serotonin-reuptake inhibitors or antiplatelet agents such as acetylsalicylic acid.

When GI bleeding or ulceration occurs in patients receiving ibuprofen containing products, the treatment should be withdrawn.

- *Dermatological effects:*



SCOTT-EDIL PHARMACIA LTD.

56, EPIP, Phase-I, Jharmajri, Baddi-173 205, (HP), India

Serious skin reactions, some of them fatal, including exfoliative dermatitis, Stevens-Johnson syndrome, and toxic epidermal necrolysis, have been reported very rarely in association with the use of NSAIDs. Patients appear to be at highest risk of these reactions early in the course of therapy, the onset of the reaction occurring in the majority of cases within the first month of treatment. Use of this product should be discontinued at the first appearance of skin rash, mucosal lesions, or any other sign of hypersensitivity.

• *Impaired female fertility:*

There is limited evidence that drugs which inhibit cyclo-oxygenase/prostaglandin synthesis may impair female fertility by an effect on ovulation and is not recommended in women attempting to conceive. This is reversible on withdrawal of treatment. In women who have difficulties conceiving or who are undergoing investigation of infertility, withdrawal of the product should be considered.

4.6 Paediatric population

Ibuplus tablet is contraindicated in adolescents under 40 kg body weight and children below 12 years of age because of the high active substance content. If in adolescents this medicinal product is required for more than 3 days, or if symptoms worsen a physician should be consulted.

4.7 Interaction with other medicinal products and other forms of Interaction

This product (like any other paracetamol containing products) is contraindicated in combination with other paracetamol containing products – increased risk of serious adverse effects.

This product (like any other ibuprofen containing products and NSAIDs) is contraindicated in combination with:

- Acetylsalicylic acid: Concomitant administration of ibuprofen and acetylsalicylic acid is not generally recommended because of the potential of increased adverse effects, unless low-dose acetylsalicylic acid (not above 75 mg daily) has been advised by a doctor.
- Experimental data suggest that Ibuprofen may competitively inhibit the effect of low dose acetylsalicylic acid on platelet aggregation when they are dosed concomitantly. Although there are uncertainties regarding extrapolation of these data to the clinical situation, the possibility that regular, long-term use of ibuprofen may reduce the cardioprotective effect of low-dose acetylsalicylic acid cannot be excluded. No clinically relevant effect is considered to be likely for occasional ibuprofen use.
- Other NSAIDs including cyclo-oxygenase-2 selective inhibitors as these may increase the risk of adverse effects.



SCOTT-EDIL PHARMACIA LTD.

56, EPIP, Phase-I, Jharmajri, Baddi-173 205, (HP), India

This product (like any other paracetamol containing products) should be used with caution in combination with:

- Cholestyramine: The speed of absorption of paracetamol is reduced by cholestyramine. Therefore, cholestyramine should not be taken within one hour if maximal analgesia is required.
- Metoclopramide and Domperidone: The absorption of paracetamol is increased by metoclopramide and domperidone. However, concurrent use need not be avoided.
- Warfarin: The anticoagulant effect of warfarin and other coumarins may be enhanced by prolonged regular use of paracetamol with increased risk of bleeding; occasional doses have no significant effect.

This product (like any other ibuprofen containing products and NSAIDs) should be used with caution in combination with:

- Anticoagulants: NSAIDs may enhance the effects of anticoagulants, i.e. warfarin .
- Antihypertensives (ACE inhibitors and Angiotensin II Antagonists) and diuretics: NSAIDs may reduce the effects of these drugs. In some patients with compromised renal function (e.g. dehydrated patients or elderly patients with compromised renal function) the co-administration of an ACE inhibitor or Angiotensin II antagonist and agents that inhibit cyclo-oxygenase may result in further deterioration of renal function, including possible acute renal failure, which is usually reversible. These interactions should be considered in patients taking a coxib concomitantly with ACE inhibitors or angiotensin II antagonists. Therefore, the combination should be administered with caution, especially in the elderly. Patients should be adequately hydrated and consideration should be given to monitoring of renal function after initiation of concomitant therapy, and periodically thereafter. Diuretics may increase the risk of nephrotoxicity of NSAIDs.
- Antiplatelet agents and selective serotonin reuptake inhibitors (SSRIs): Increased risk of gastrointestinal bleeding.
- Cardiac glycosides: NSAIDs may exacerbate cardiac failure, reduce GFR and increase plasma glycoside levels.
- Ciclosporin: Increased risk of nephrotoxicity.
- Corticosteroids: Increased risk of gastrointestinal ulceration or bleeding.
- Lithium: Decreased elimination of lithium.
- Methotrexate: Decreased elimination of methotrexate.
- Mifepristone: NSAIDs should not be used for 8-12 days after mifepristone administration as NSAIDs can reduce the effect of mifepristone.



SCOTT-EDIL PHARMACIA LTD.

56, EPIP, Phase-I, Jharmajri, Baddi-173 205, (HP), India

- Quinolone antibiotics: Animal data indicate that NSAIDs can increase the risk of convulsions associated with quinolone antibiotics. Patients taking NSAIDs and quinolones may have an increased risk of developing convulsions.
- Tacrolimus: Possible increased risk of nephrotoxicity when NSAIDs are given with tacrolimus.
- Zidovudine: Increased risk of haematological toxicity with NSAIDs are given with zidovudine. There is evidence of an increased risk of haemarthroses and haematoma in HIV (+) haemophiliacs receiving concurrent treatment with zidovudine and ibuprofen.

4.8 Additional information on special populations

Pediatric patients

Ibuplus tablet is contraindicated in adolescents under 40 kg body weight and children below 12 years of age because of the high active substance content. If in adolescents this medicinal product is required for more than 3 days, or if symptoms worsen a physician should be consulted.

Elderly patients

No special dosage adjustments are required. Due to possible undesirable effect profile, it is recommended to monitor the elderly particularly carefully.

Hepatic impairment

No dose adjustments are required in patients with mild to moderate impairment to hepatic function (patients with severe hepatic dysfunction).

Renal impairment

No dose adjustments is required in patients with mild to moderate impairment to renal function (patients with severe renal insufficiency).

4.9 Paediatric population

Ibuplus tablet is contraindicated in adolescents under 40 kg body weight and children below 12 years of age because of the high active substance content. If in adolescents this medicinal product is required for more than 3 days, or if symptoms worsen a physician should be consulted.

4.10 Fertility, pregnancy and lactation

Pregnancy:



SCOTT-EDIL PHARMACIA LTD.

56, EPIP, Phase-I, Jharmajri, Baddi-173 205, (HP), India

There is no experience of use of this product in humans during pregnancy.

A large amount of data on pregnant women indicate neither malformative, nor fetoneonatal toxicity. Epidemiological studies on neurodevelopment in children exposed to paracetamol in utero show inconclusive results. If clinically needed, paracetamol can be used during pregnancy however it should be used at the lowest effective dose for the shortest possible time and at the lowest possible frequency.

Congenital abnormalities have been reported in association with NSAID administration in man; however, these are low in frequency and do not appear to follow any discernible pattern. In view of the known effects of NSAIDs on the foetal cardiovascular system (risk of closure of ductus arteriosus), use in the last trimester is contraindicated. The onset of labour may be delayed, and duration increased with an increased bleeding tendency in both mother and child. NSAIDs should not be used during the first two trimesters of pregnancy or labour unless the potential benefit to the patient outweighs the potential risk to the foetus.

Therefore if possible, the use of this product should be avoided in the first six months of pregnancy and contraindicated in the last three months of pregnancy.

Lactation:

Ibuprofen and its metabolites can pass in very small amounts (0.0008% of the maternal dose) into the breast milk. No harmful effects to infants are known.

Paracetamol is excreted in breast milk but not in a clinically significant amount. Available published data do not contraindicate breastfeeding.

Therefore it is not necessary to interrupt breastfeeding for short-term treatment with the recommended dose of this product.

4.11 Undesirable effects

Clinical trials with this product have not indicated any other undesirable effects other than those for ibuprofen or paracetamol alone.

The following table lists adverse effects from pharmacovigilance data experienced by patients taking ibuprofen alone or paracetamol alone in short-term and long-term use.

Adverse events which have been associated with Ibuprofen alone or Paracetamol alone are given below, tabulated by system organ class and frequency. Frequencies are defined as: very common ($\geq 1/10$), common ($\geq 1/100$ to $< 1/10$), uncommon ($\geq 1/1,000$ to $< 1/100$), rare ($\geq 1/10,000$ to $< 1/1,000$), very rare ($< 1/10,000$) and not known (cannot be estimated from the available data). Within each frequency grouping, adverse events are presented in order of decreasing seriousness.

System Organ Class	Frequency	Adverse Event
--------------------	-----------	---------------



SCOTT-EDIL PHARMACIA LTD.

56, EPIP, Phase-I, Jharmajri, Baddi-173 205, (HP), India

Blood and Lymphatic System Disorders	Very rare	Haematopoietic disorders
Immune System Disorders	Uncommon	Hypersensitivity with urticaria and pruritus
	Very rare	Severe hypersensitivity reactions. Symptoms can include facial, tongue and throat swelling, dyspnoea, tachycardia, hypotension (anaphylaxis, angioedema or severe shock)
Psychiatric Disorders	Very rare	Confusion, depression and hallucinations
Nervous System Disorders	Uncommon	Headache and dizziness
	Very rare	Aseptic meningitis, paraesthesia, optic neuritis and somnolence
Eye Disorders	Very rare	Visual disturbance
Ear and Labyrinth Disorders	Very rare	Tinnitus and vertigo
Cardiac Disorders	Very rare	Cardiac failure and oedema ⁴
Vascular Disorders	Very rare	Hypertension
Respiratory and thoracic and mediastinal disorders	Very rare	Respiratory reactivity including: asthma, exacerbation of asthma, bronchospasm and dyspnoea
Gastrointestinal Disorders	Common	Abdominal pain, vomiting, diarrhoea, nausea, dyspepsia and abdominal discomfort
	Uncommon	peptic ulcer, gastrointestinal perforation or gastrointestinal haemorrhage, melaena, haematemesis, mouth ulceration, exacerbation of colitis and Crohn's disease gastritis, pancreatitis, flatulence and constipation
Hepatobiliary Disorders	Very rare	Abnormal liver function, hepatitis and jaundice
Skin and Subcutaneous Tissue Disorders	Common	Hyperhidrosis
	Uncommon	Various skin rashes

	Very rare	Bullous reactions including Stevens-Johnson syndrome, erythema multiforme and toxic epidermal necrolysis.Exfoliative dermatoses, purpura, photosensitivity
	Not known	Drug reaction with eosinophilia and systemic symptoms (DRESS syndrome)
Renal and Urinary Disorders	Very rare	Nephrotoxicity in various forms, including interstitial nephritis, nephrotic syndrome, and acute and chronic renal failure
General Disorders and Administration Site Conditions	Very rare	Fatigue and malaise
Investigations	Common	Alanine aminotransferase increased, gamma-glutamyltransferase increased and liver function tests abnormal with paracetamol. Blood creatinine increased, blood urea increased.
	Uncommon	Aspartate aminotransferase increased, blood alkaline phosphatase increased, blood creatine phosphokinase increased, haemoglobin decreased and platelet count increased.

4.12 Overdose

Paracetamol

Liver damage is possible in adults who have taken 10 g (equivalent to 20 tablets) or more of paracetamol. Ingestion of 5 g (equivalent to 10 tablets) or more of paracetamol may lead to liver damage if the patient has one or more of the risk factors below:

- a) Is on long term treatment with carbamazepine, phenobarbitone, phenytoin, primidone, rifampicin, St John's Wort or other drugs that induce liver enzymes.
- b) Regularly consumes alcohol in excess of recommended amounts.
- c) Is likely to be glutathione depleted e.g. eating disorders, cystic fibrosis, HIV infection, starvation, cachexia

Symptoms



SCOTT-EDIL PHARMACIA LTD.

56, EPIP, Phase-I, Jharmajri, Baddi-173 205, (HP), India

Symptoms of paracetamol overdose in the first 24 hours include pallor, nausea, vomiting, anorexia and abdominal pain. Liver damage may become apparent 12 to 48 hours after ingestion as liver function tests become abnormal. Abnormalities of glucose metabolism and metabolic acidosis may occur. In severe poisoning, hepatic failure may progress to encephalopathy, haemorrhage, hypoglycaemia, cerebral oedema and death. Acute renal failure with acute tubular necrosis, strongly suggested by loin pain, haematuria and proteinuria, may develop even in the absence of severe liver damage. Cardiac arrhythmias and pancreatitis have been reported.

Management

Immediate treatment is essential in the management of paracetamol overdose. Despite a lack of significant early symptoms, patients should be referred to hospital urgently for immediate medical attention. Symptoms may be limited to nausea or vomiting and may not reflect the severity of overdose or the risk of organ damage. Management should be in accordance with established treatment guidelines.

Treatment with activated charcoal should be considered if the overdose has been taken within 1 hour. Plasma paracetamol concentration should be measured at 4 hours or later after ingestion (earlier concentrations are unreliable).

Treatment with N-acetylcysteine may be used up to 24 hours after ingestion of paracetamol however; the maximum protective effect is obtained up to 8 hours post ingestion. The effectiveness of the antidote declines sharply after this time.

If required the patient should be given intravenous-N-acetylcysteine, in line with the established dosage schedule. If vomiting is not a problem, oral methionine may be a suitable alternative for remote areas, outside hospital.

Patients who present with serious hepatic dysfunction beyond 24 hours from ingestion should be managed in accordance with established guidelines.

Ibuprofen

In children ingestion of more than 400 mg/kg of Ibuprofen may cause symptoms. In adults the dose response effect is less clear cut.

The half-life in overdose is 1.5-3 hours.

Symptoms

Most patients who have ingested clinically important amounts of NSAIDs will develop no more than nausea, vomiting, epigastric pain, or more rarely diarrhoea. Tinnitus, headache and gastrointestinal bleeding are also possible. In more serious poisoning, toxicity is seen in the central nervous system, manifesting as drowsiness, occasionally excitation and disorientation or coma. Occasionally patients develop convulsions. In serious poisoning metabolic acidosis may occur and the prothrombin time / INR may be prolonged, probably due to interference



SCOTT-EDIL PHARMACIA LTD.

56, EPIP, Phase-I, Jharmajri, Baddi-173 205, (HP), India

with the actions of circulating clotting factors. Acute renal failure and liver damage may occur if there is a co-incident of dehydration. Exacerbation of asthma is possible in asthmatics.

Management

Management should be symptomatic and supportive and include the maintenance of a clear airway and monitoring of cardiac and vital signs until stable. Consider oral administration of activated charcoal if the patient presents within 1 hour of ingestion of a potentially toxic amount. If frequent or prolonged, convulsions should be treated with intravenous diazepam or lorazepam. Give bronchodilators for asthma.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Musculoskeletal system, anti-inflammatory and antirheumatic products, non-steroids, propionic acid derivatives. Ibuprofen combinations

ATC code: M01AE51

The pharmacological actions of ibuprofen and paracetamol differ in their site and mode of action. These complementary modes of action are synergistic which results in greater antinociception and antipyresis than the single actives alone.

Ibuprofen is an NSAID that has demonstrated its efficacy in the common animal experimental inflammation models by inhibition of prostaglandin synthesis. Prostaglandins sensitise nociceptive afferent nerve terminals to mediators such as bradykinin. Ibuprofen therefore elicits an analgesic effect through peripheral inhibition of the cyclooxygenase-2 (COX-2) isoenzyme with a subsequent reduction in sensitisation of nociceptive nerve terminals. Ibuprofen has also been shown to inhibit induced-leucocyte migration into inflamed areas. Ibuprofen has a pronounced action within the spinal cord due, in part, to the inhibition of COX. Ibuprofen's antipyretic effects are produced by the central inhibition of prostaglandins in the hypothalamus. Ibuprofen reversibly inhibits platelet aggregation. In humans, ibuprofen reduces inflammatory pain, swellings and fever.

Experimental data suggest that ibuprofen may competitively inhibit the effect of low dose acetylsalicylic acid on platelet aggregation when they are dosed concomitantly. Some pharmacodynamic studies show that when single doses of ibuprofen 400mg was taken within 8 h before or within 30 min after immediate release acetylsalicylic acid dosing (81mg), a decreased effect of acetylsalicylic acid on the formation of thromboxane or platelet aggregation occurred. Although there are uncertainties regarding extrapolation of these data to the clinical situation, the possibility that regular, long-term use of ibuprofen may reduce the cardioprotective effect of low-dose acetylsalicylic acid cannot be excluded. No clinically relevant effect is considered to be likely for occasional ibuprofen use.

Paracetamol's exact mechanism of action is still not completely defined; however there is considerable evidence to support the hypothesis of a central antinociceptive effect. Various



SCOTT-EDIL PHARMACIA LTD.

56, EPIP, Phase-I, Jharmajri, Baddi-173 205, (HP), India

biochemical studies point to inhibition of central COX-2 activity. Paracetamol may also stimulate the activity of descending 5-hydroxytryptamine (serotonin) pathways that inhibit nociceptive signal transmission in the spinal cord. Evidence has shown that paracetamol is a very weak inhibitor of peripheral COX-1 and 2 isoenzymes.

The clinical efficacy of ibuprofen and paracetamol has been demonstrated in pain associated with headache, toothache and dysmenorrhoea, and fever; furthermore efficacy has been shown in patients with pain and fever associated with cold and influenza and in pain models such as sore throat, muscular pain or soft tissue injury and backache.

This product is especially suitable for pain which requires stronger pain relief than ibuprofen 400 mg or paracetamol 1000 mg alone, and faster pain relief than ibuprofen.

Summary of 2 tablet clinical data

A randomised, double-blind placebo-controlled studies were conducted with the combination using the acute pain model of post-operative dental pain. The studies show that:

- This product provides more effective pain relief than paracetamol 1000 mg ($p < 0.0001$) and ibuprofen 400 mg ($p < 0.05$) which are clinically and statistically significant.
- This product has a fast onset of action with 'confirmed perceptible pain relief' achieved in a median of 18.3 minutes. The onset of action was significantly more rapid than for ibuprofen 400 mg (23.8 minutes, $p = 0.0015$). 'Meaningful pain relief' for this product was achieved in a median of 44.6 minutes, which was significantly faster than for ibuprofen 400 mg (70.5 minutes, $p < 0.0001$).
- Duration of analgesia was significantly longer for this product (9.1 hours) compared to paracetamol 500 mg (4 hours) or 1000 mg (5 hours).
- The global evaluation of the study medication by the subjects showed high levels of satisfaction with 93.2% rating the product as 'good', 'very good' or 'excellent' in achieving pain relief. The fixed combination product performed significantly better than paracetamol 1000 mg ($p < 0.0001$).

A randomised, double-blind controlled clinical study was conducted with the product in the treatment of chronic knee pain. The study showed that:

- The product provides more effective pain relief than paracetamol 1000 mg in short-term treatment ($p < 0.01$) and long term treatment ($p < 0.01$).
- The global evaluation of the product by the subjects showed high levels of satisfaction with 60.2% rating the product as 'good' or 'excellent' as a long term treatment for a painful knee. The product performed significantly better than paracetamol 1000 mg ($p < 0.001$).

5.2 Pharmacokinetic properties



SCOTT-EDIL PHARMACIA LTD.

56, EPIP, Phase-I, Jharmajri, Baddi-173 205, (HP), India

Ibuprofen is well absorbed from the gastrointestinal tract and is extensively bound to plasma proteins. Ibuprofen diffuses into the synovial fluid. Plasma levels of ibuprofen from this product are detected from 5 minutes with peak plasma concentrations achieved within 1-2 hours after ingestion on an empty stomach. When this product was taken with food peak ibuprofen plasma levels were lower and delayed by a median of 25 minutes, but overall extent of absorption was equivalent.

Ibuprofen is metabolised in the liver to two major metabolites with primary excretion via the kidneys, either as such or as major conjugates, together with a negligible amount of unchanged ibuprofen. Excretion by the kidney is both rapid and complete. The elimination half-life is approximately 2 hours.

In limited studies, ibuprofen appears in the breast milk in very low concentrations.

No significant differences in ibuprofen pharmacokinetic profile are observed in the elderly.

Paracetamol is readily absorbed from the gastrointestinal tract. Plasma protein binding is negligible at usual therapeutic concentrations, although this is dose-dependent. Plasma levels of paracetamol from this product are detected from 5 minutes with peak plasma concentrations occurring at 0.5-0.67 hours after ingestion on an empty stomach. When this product was taken with food peak paracetamol plasma levels were lower and delayed by a median of 55 minutes, but overall extent of absorption was equivalent.

Paracetamol is metabolised in the liver and excreted in the urine mainly as the glucuronide and sulphate conjugates, with about 10% as glutathione conjugates. Less than 5% is excreted as unchanged paracetamol. The elimination half-life is approximately 3 hours.

A minor hydroxylated metabolite, which is usually produced in very small amounts by mixed function oxidases in the liver and detoxified by conjugation with liver glutathione, may accumulate following paracetamol overdose and cause liver damage.

No significant differences in the paracetamol pharmacokinetic profile are observed in the elderly.

The bioavailability and pharmacokinetic profiles of ibuprofen and paracetamol taken as this product are not altered when taken in combination as a single or repeat dose.

This product is formulated using a technology which releases both Ibuprofen and Paracetamol simultaneously, so that the active ingredients deliver a combination effect.

5.3 Preclinical safety data

The toxicological safety profile of ibuprofen and paracetamol has been established in animal experiments and in humans from extensive clinical experience. There are no new preclinical data of relevance to the prescriber which are additional to the data already presented in this Summary of Product Characteristics.



SCOTT-EDIL PHARMACIA LTD.

56,EPIP,Phase-I,Jharmajri,Baddi-173 205, (HP), India

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Maize starch	BP
Microcrystalline Cellulose	BP
Povidone K- 30	BP
Methyl Hydroxybenzoate	BP
Propyl Hydroxybenzoate	BP
Purified Talc	BP
Magnesium Stearate	BP
Sodium Starch Glycolate	BP
Croscarmellose Sodium	BP
Colloidal Anhydrous silica	BP

6.2 Incompatibilities

Not Applicable.

6.3 Shelf life

36 Months

6.4 Special precautions for storage

Store protected from light at a temperature not exceeding 30°C

6.5 Nature and contents of container

10 PVC Blister of 10 tablets are packed in a carton along with a leaflet.

6.6 Special precautions for disposal and other handling

No Special requirements.

7. Marketing Authorisation Holder And Manufacturing Site Addresses

Scott-Edil Pharmacia Ltd.,
56,EPIP,Phase-I, Jharmajri, Baddi-173205, (HP), INDIA



SCOTT-EDIL PHARMACIA LTD.

56,EPIP,Phase-I,Jharmajri,Baddi-173 205, (HP), India

MANUFACTURING SITE ADDRESS

Scott-Edil Pharmacia Ltd.,
56,EPIP,Phase-I, Jharmajri, Baddi-173205, (HP), INDIA

8. MARKETING AUTHORISATION NUMBER

Not Applicable

9. DATE OF FIRST REGISTRATION/RENEWAL OF THE REGISTRATION

Not Applicable.

10. DATE OF REVISION OF THE TEXT

Not Applicable

11. DOSIMETRY

Not Applicable