

105mm

150mm

Stopkof P

Pediatric Cough Syrup

Composition:

Each 5 ml contains Diphenhydramine Hydrochloride 7 mg

Introduction:

Stopkof P is a cough preparation, containing Diphenhydramine Hydrochloride having antihistaminic, antitussive and anticholinergic properties.

Indications:

Stopkof P is indicated for relieving of cough symptoms due to cold or allergy respiratory tract congestion.

Dosage:

For Children:

6-12 years : 10 ml of Stopkof P syrup 4 times a day or as directed by the physician

Not recommended for children under 6 years.

Precautions:

Care should be taken while administrating Stopkof P syrup to patients with hypersensitivity to Diphenhydramine Hydrochloride.

Concomitant use of Anti-depressants or Atropine will increase the chances of side effects such as drowsiness and or dry mouth.

Contraindication:

Stopkof P syrup should not be given to patients receiving MAO inhibitors or CNS depressants and in patients with reduced liver function and metabolic alkalosis.

Side Effects:

Adverse effects are usually due to

Diphenhydramine that enhances sedative effects, which can cause drowsiness or difficulty in concentration. Occasionally upset of stomach, constipation, nausea, dry mouth, nose or throat, difficulty in passing urine blurred vision may also occur.

Interaction with other drugs:

Stopkof P syrup should not be given to patients receiving MAO inhibitors or CNS depressants.

Patient Instructions:

Stopkof P syrup may cause sedation and drowsiness. Hence caution to be exercised while performing tasks requiring mental alertness.

Storage: Store below 30° C, protected from light.

Presentation: 60 & 100 ml syrup in bottle.

THIS IS A MEDICAMENT

- Medicament is a product which affects your health and its' consumption contrary to instruction is dangerous for you.
- Follow strictly the doctor's prescription, the method of use and the instruction of the pharmacist who sold the medicament.
- The doctor and the pharmacist are experts in medicine, its benefits and risks.
- Do not by yourself interrupt the period of treatment pre-scribed for you.
- Do not repeat the same prescription without consulting your doctor.

Keep medicament out of the reach of children.
Council of Arab Health Ministers
Union of Arab Pharmacists

Manufactured by :

National Pharmaceutical Industries Co. (SAOG)

Street No 15, Rusayl Industrial Area,

Muscat, Sultanate of Oman.

Rev. No: 00

Rev. Date 011/16

NCLA 004

Pharma Code 2503